Enhancement of the Benjamin-Feir instability with dissipation

T. J. Bridges

Department of Mathematics, University of Surrey, Guildford, Surrey GU2 7XH, England

F. Dias

CMLA, ENS Cachan, CNRS, PRES UniverSud, 61, avenue du President Wilson, 94230 Cachan cedex, France

(Received 11 May 2007; accepted 16 August 2007; published online 17 October 2007)

It is shown that there is an overlooked mechanism whereby some kinds of dissipation can enhance the Benjamin-Feir instability of water waves. This observation is new, and although it is counterintuitive, it is due to the fact that the Benjamin-Feir instability involves the collision of modes with opposite energy sign (relative to the carrier wave), and it is the *negative energy perturbations* that are enhanced. © 2007 American Institute of Physics. [DOI: 10.1063/1.2780793]

The discovery of the Benjamin-Feir (BF) instability of traveling waves was a milestone in the history of water waves. Before 1960, the idea that a Stokes wave could be unstable does not appear to be given much thought. The possibility that the Stokes wave could be unstable was pointed out in the early 1960s,^{1–4} but it was the seminal work of Benjamin and Feir^{5,6} that combined experimental evidence with a weakly nonlinear theory that convinced the scientific community.

Indeed, Benjamin and Feir started their experiments in 1963 *assuming that the Stokes wave was stable*. After several frustrating years watching their waves disintegrate—in spite of equipment and laboratory changes and improvements—they finally came to the conclusion that they were witnessing a new kind of instability. The appearance of "sidebands" in the experiments suggested the form that the perturbations should take. A history of these experiments and the outcome are reported in Ref. 7.

The theory of the BF instability is based on inviscid fluid mechanics, and the assumption that the system is conservative. Therefore, it is natural to study the implication of perturbations on the system. The implications of a range of perturbations on the BF instability have been studied in the literature: for example, the effect of wind^{8,9} and the effect of viscosity.^{8,10-13} Some perturbations have been shown to stabilize and others to destabilize the BF instability.

However, there is a fundamental overlooked mechanism in all this work. Mathematically, the BF instability can be characterized as a collision of two pairs of purely imaginary eigenvalues of opposite energy sign, as shown in Fig. 1. In Ref. 14, this observation is implicit, but the demonstration and implications have not been given heretofore. This characterization of the BF instability also appears in the nonlinear Schrödinger (NLS) model for modulation of dispersive traveling waves.^{15,16} The eigenvalue with a smaller positive imaginary part in the figure—just before collision—has negative energy, whereas the eigenvalue with larger imaginary part has positive energy. This energy is relative to the energy of the carrier wave E^{Stokes} : $E_- < E^{\text{Stokes}} < E_+$, where E_+ are the energies of the modes associated with the respective purely imaginary eigenvalues in Fig. 1. Hence, "negative energy" means that $E_{-}-E^{\text{Stokes}} < 0$.

Once these facts are established, we can appeal to the result that dissipation can *destabilize* negative energy modes.^{17–19} There are many examples in fluid mechanics where negative energy modes—which are stable in the inviscid limit—are destabilized by the addition of dissipation:²⁰ Kelvin-Helmholtz instability,^{17,18,21} interaction of a fluid with a flexible boundary,^{22,23} stability of a fluid-loaded elastic plate,²⁴ and Euler modes perturbed by the Navier-Stokes equations.²⁵

The book of Fabrikant and Stepanyants²¹ reports on experimental results for interfacial waves near the Kelvin-Helmholtz instability illustrating the enhancement of the instability of negative energy waves due to dissipation (see Fig. 3.5 on p. 83, and the discussion on pp. 82–83 in Ref. 21).

In this brief communication, we sketch the basic result for water waves and then use a nonlinear Schrödinger model perturbed by dissipation for illustration. The NLS equation has shortcomings (e.g., symmetry that enables the phase to be factored out, lack of validity for all time²⁶), but it provides a simple example of the phenomenon.

Davey¹⁰ gives a general argument for the form of a dissipation-perturbed NLS model, and Blennerhassett⁸ starts with the full Navier-Stokes equations for a free-surface flow with viscous free-surface boundary conditions and derives a similar perturbed NLS equation. For the dissipatively perturbed Stokes wave in deep water, these NLS models take the form

$$iA_t + (\alpha - ia)A_{xx} + ibA + (\gamma + ic)|A|^2A = 0,$$
 (1)

where A is the envelope of the wave carrier, and the modulations are restricted to one space dimension x. When a=b=c=0, Eq. (1) reduces to the NLS equation for the modulations of Stokes waves in deep water; hence, α and γ are positive real numbers. This NLS model has a BF instability, and one can show explicitly that it involves a collision of eigenvalues of the form shown in Fig. 1. We show below that when a>0, there is always dissipation induced instability (before the BF instability), no matter how small a is. The

19, 104104-1



FIG. 1. Schematic of the eigenvalue movement associated with the BF instability, as a function of the amplitude of the basic carrier wave (Amp) and the sideband wavenumber σ . For a fixed σ (vertical dashed line), there is a threshold amplitude. Below the threshold, the eigenvalues are purely imaginary. At the threshold a collision occurs and above the threshold, the eigenvalues are complex.

parameter *a* is the perturbation of the rate of change of the group velocity dc_{e}/dk due to dissipation.

First, consider the linear stability problem for gravity waves in deep water. As the wave amplitude increases, we show that there is a threshold value at which two eigenvalues of the linear stability problem collide, and these two modes have negative and positive energy, respectively.

With $\theta = x - ct$, the speed *c* and amplitude η of the basic gravity wave of wavelength $2\pi/k$, to leading order, are $c = c_0(1 + k^2\varepsilon^2 + \cdots), c_0^2 = gk^{-1}$,

$$\eta(\theta) = \varepsilon \,\eta_1(\theta) + \varepsilon^2 \,\eta_2(\theta) + \mathcal{O}(\varepsilon^3),$$

where ε is a measure of the amplitude,

$$\eta_1(\theta) = \sqrt{2}\cos(k\theta - \theta_0), \quad \eta_2(\theta) = k\cos(2k\theta - 2\theta_0),$$

with θ_0 an arbitrary phase shift. Using standard results on integral properties of Stokes waves, the total energy *relative* to the moving frame is

$$E^{\text{Stokes}} = T + V - cI = V - T, \quad \text{using } 2T = cI,$$

where *T* and *V* are the kinetic and potential energies, respectively, and *I* is the momentum.²⁷ Substitution of the Stokes expansion shows that $E^{\text{Stokes}}=0+\varepsilon^3 E_3+\mathcal{O}(\varepsilon^4)$. Although the actual value of E_3 is not important for the argument below, it is noteworthy that it is negative, and, using Table 2 of Ref. 27, one can confirm that E^{Stokes} is negative at finite amplitude as well.

To formulate the linear stability for gravity waves take

$$\eta(\theta, x, t) \mapsto \hat{\eta}(\theta, \varepsilon) + \eta(\theta, x, t),$$

where $\hat{\eta}(\theta, \varepsilon)$ is the basic carrier wave. Take a similar expression for the velocity potential $\phi(\theta, x, y, t)$, where y denotes the vertical space dimension. Next one substitutes this form into the water wave equations, linearizes about the carrier wave, and takes $\eta(\theta, x, t)$ of the form

$$\eta(\theta, x, t) = \operatorname{Re}[\Sigma(\theta, \sigma) e^{i\sigma x + \lambda t}],$$

where σ is real (the modulation wavenumber), and $\Sigma(\theta, \sigma)$ is periodic of the same period as the Stokes wave. The result is an eigenvalue problem for the eigenfunction Σ and eigenvalue λ .

The BF instability corresponds to a solution of this eigenvalue problem with $0 < \sigma \ll 1$ and $\text{Re}(\lambda) > 0$. When σ is fixed—but nonzero and small—and the amplitude of the Stokes wave is increased, there is a threshold amplitude where the BF instability occurs, and it corresponds to a collision of two eigenvalues as shown in Fig. 1. To leading order, the eigenvalues collide at $\lambda = \pm i c_g \sigma$, where $c_g = \frac{1}{2} \sqrt{g/k}$ is the group velocity.

To show that the colliding modes have opposite energy sign, we need a definition of the energy of the perturbation. This definition requires some consideration because the perturbation is quasiperiodic in space: $2\pi/k$ -periodic in θ , and $2\pi/\sigma$ -periodic in x. The total energy relative to the moving frame is

$$E^{\text{total}} = \frac{\sigma}{2\pi} \int_0^{2\pi/\sigma} \frac{k}{2\pi} \int_0^{2\pi/k} \hat{E} \, \mathrm{d}\theta \, \mathrm{d}x,$$

where $\hat{E} = \hat{T} - \hat{V} - c\hat{I}$, $\hat{V} = 1/2g \eta^2$,

$$\hat{T} = \int_{-\infty}^{\eta} \frac{1}{2} (\phi_{\theta}^2 + 2\phi_{\theta}\phi_x + \phi_x^2 + \phi_y^2) dy, \text{ and}$$

$$\hat{I} = \int_{-\infty}^{\eta} (\phi_{\theta} + \phi_x) \mathrm{d}y.$$

Evaluating the perturbation energy for the two modes that collide leads to $E^{\text{total}} = E^{\text{Stokes}} + \varepsilon^2 E_2^{\pm} + \cdots$, with

$$E_2^{\pm} = 2(k \pm \sigma) \left(1 - \sqrt{1 \mp \frac{\sigma}{k}}\right) |C_{\pm}|^2.$$

Here, C_{\pm} are scale factors associated with the eigenfunctions. Clearly, $sign(E_2^+E_2^-) < 0$ for $0 < \sigma < 1$.

Having shown that the colliding modes have opposite energy signs, we consider a simple example that illustrates the mechanism for destabilization of negative energy modes by damping. A prototype for a conservative system, where the linearization has a collision of eigenvalues of opposite energy sign, which is perturbed by Rayleigh damping, is

$$\mathbf{q}_{tt} + 2b\mathbf{J}\mathbf{q}_t + (\chi - \tau^2)\mathbf{q} + 2\delta\mathbf{q}_t = \mathbf{0}, \quad \mathbf{q} \in \mathbb{R}^2,$$

$$\mathbf{J} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$
(2)

where $\tau > 0$ is the "gyroscopic coefficient," χ a real parameter with $|\chi| \ll \tau^2$, and $\delta \ge 0$.

The energy of the system (2) is strictly decreasing when $\delta > 0$ and $\|\mathbf{q}_t\| > 0$. Let $\mathbf{q}(t) = \hat{\mathbf{q}} e^{\lambda t}$; then substitution into (2) leads to the roots

Downloaded 27 Mar 2008 to 128.95.176.16. Redistribution subject to AIP license or copyright; see http://pof.aip.org/pof/copyright.jsp



FIG. 2. Schematic of the effect of dissipation on the eigenvalues associated with (2).

$$\lambda = i\tau - \delta \pm i\sqrt{\chi + 2i\tau\delta - \delta^2} \quad \text{and}$$

$$\lambda = -i\tau - \delta \pm i\sqrt{\chi - 2i\tau\delta - \delta^2}.$$
(3)

When $\delta=0$, there are four roots: $\lambda = \pm i(\tau \pm \sqrt{\chi})$. The eigenvalue movement shown in Fig. 1 is realized as χ decreases from a positive value to a negative value, the collision occurring at $\chi=0$. Suppose that χ is small and positive (just before the collision) and look at the effect of dissipation on the two modes $\lambda_0 = i\tau \pm i\sqrt{\chi}$. Substitution of the eigenfunctions for these two eigenvalues into the energy shows that the mode associated with $i\tau - i\sqrt{\chi}$ has negative energy, while the mode associated with $i\tau + i\sqrt{\chi}$ has positive energy.

With δ small, expand the first pair of roots in (3) in a Taylor series

$$\lambda(\delta) = i\tau \pm i\sqrt{\chi} \mp \frac{\delta}{\sqrt{\chi}}(\tau \pm \sqrt{\chi}) + \mathcal{O}(\delta^2).$$

With $0 < \delta \le 1$, the eigenvalues are perturbed, as shown to the right in Fig. 2. The negative energy mode, i.e., $\lambda_0 = i(\tau - \sqrt{\chi})$, has a positive real part when dissipatively perturbed, and the positive energy mode, i.e., $\lambda_0 = i(\tau + \sqrt{\chi})$, has a negative real part under perturbation. Consequently, when small dissipation is added to the otherwise *stable* system (that is, $0 < \chi \le \tau^2$), the mode with negative energy will destabilize. After the collision (when $\chi < 0$), the growth rate of the instability is enhanced.

It should be noted that other mathematically consistent forms of damping can be used. For example, the uniform damping

$$\mathbf{q}_t = \frac{\partial H}{\partial \mathbf{p}} - \delta \mathbf{q}, \quad \mathbf{p}_t = -\frac{\partial H}{\partial \mathbf{q}} - \delta \mathbf{p},$$
 (4)

makes mathematical sense. However, it leads to uniform contraction of the phase space, and does not destabilize negative energy modes. In order to study the effect of dissipation on water waves, one could start with the Navier-Stokes equations and perturb about the Stokes wave solution (see Ref. 8, for instance, for the case of wind forcing). Another approach is to add viscous perturbations to the potential flow in various forms.²⁸ From the modified equations one can derive a dissipative NLS equation. There are two issues to highlight: negative energy modes can be destabilized, so the BF instability can be enhanced by dissipation, and secondly, the form of the damping is important. It is known that negative energy modes of the Euler equations can be destabilized by the form of damping found in the Navier-Stokes equations.²⁵

Following Refs. 8 and 10, a general perturbed NLS equation for various types of physical situations can be written in the form (1). The parameters a, b, and c are taken to be non-negative. When they are positive, they represent dissipative perturbations, since the norm of the solution is strictly decreasing in time when $a^2+b^2+c^2>0$.

When a=b=c=0, the resulting NLS equation is a Hamiltonian partial differential equation; with $A=u_1+iu_2$ and $\mathbf{u}=(u_1,u_2)$,

$$\mathbf{J}\mathbf{u}_{t} = \nabla H(\mathbf{u}) + a\mathbf{J}\mathbf{u}_{xx} - b\mathbf{J}\mathbf{u} - c\|\mathbf{u}\|^{2}\mathbf{J}\mathbf{u},$$
(5)

where \mathbf{J} was defined in (2), and

$$H(\mathbf{u}) = \int_{\mathbb{R}} \left[\frac{1}{2} \alpha \|\mathbf{u}_x\|^2 - \frac{1}{4} \gamma \|\mathbf{u}\|^4 \right] \mathrm{d}x.$$
 (6)

Let $\theta(x,t) = kx - \omega t + \theta_0$, and consider the basic traveling wave solution to (5) when dissipation is neglected,

$$\hat{\mathbf{u}}(x,t) = \mathbf{R}_{\theta(x,t)}\mathbf{u}_0, \quad \mathbf{R}_{\theta} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$
(7)

Then, \mathbf{u}_0 , ω , k satisfy $-\omega + \alpha k^2 = \gamma \|\mathbf{u}_0\|^2$.

It is assumed that the Stokes traveling wave exists for a sufficiently long time before any dissipation can affect it: dissipation is taken to be a second-order effect.

Next we check the energetics of the BF stability problem in NLS. Linearize the partial differential equation (5) with a=b=c=0 about the basic traveling wave (7). Letting $\mathbf{u}(x,t)=\mathbf{R}_{\theta(x,t)}[\mathbf{u}_0+\mathbf{v}(x,t)]$, substituting into the conservative version of (5), linearizing about \mathbf{u}_0 , and simplifying, leads to

$$\mathbf{J}\mathbf{v}_t + 2\alpha k \mathbf{J}\mathbf{v}_x + \alpha \mathbf{v}_{xx} + 2\gamma \langle \mathbf{u}_0, \mathbf{v} \rangle \mathbf{u}_0 = \mathbf{0},$$
(8)

where $\langle \cdot, \cdot \rangle$ is the standard scalar product on \mathbb{R}^2 .

The class of solutions of interest are solutions that are periodic in *x* with wavenumber σ . The parameter σ represents the sideband. The BF instability will be associated with the limit $|\sigma| \leq 1$. Therefore, let

$$\mathbf{v}(x,t) = \frac{1}{2}\mathbf{v}_0(t) + \sum_{n=1}^{\infty} \left[\mathbf{v}_n(t)\cos n\sigma x + \mathbf{w}_n(t)\sin n\sigma x\right].$$

Neglecting the σ -independent modes (superharmonic instability), the σ -dependent modes decouple into fourdimensional subspaces for each *n*, and satisfy

Downloaded 27 Mar 2008 to 128.95.176.16. Redistribution subject to AIP license or copyright; see http://pof.aip.org/pof/copyright.jsp

$$\mathbf{J}\dot{\mathbf{v}}_{n} + 2\alpha kn\sigma \mathbf{J}\mathbf{w}_{n} - \alpha(n\sigma)^{2}\mathbf{v}_{n} + 2\gamma \mathbf{u}_{0}\mathbf{u}_{0}^{T}\mathbf{v}_{n} = \mathbf{0},$$

$$\mathbf{J}\dot{\mathbf{w}}_{n} - 2\alpha kn\sigma \mathbf{J}\mathbf{v}_{n} - \alpha(n\sigma)^{2}\mathbf{w}_{n} + 2\gamma \mathbf{u}_{0}\mathbf{u}_{0}^{T}\mathbf{w}_{n} = \mathbf{0}.$$
(9)

When the amplitude $\|\mathbf{u}_0\|=0$, it is easy to show that all eigenvalues of the above system (i.e., taking solutions of the form $e^{\lambda t}$ and computing λ) are purely imaginary. Considering all other parameters fixed, and increasing $\|\mathbf{u}_0\|$, we find that there is a critical amplitude where the n=1 mode becomes unstable first through a collision of eigenvalues of opposite signature.

To analyze this instability, take n=1 and study the reduced four-dimensional system

$$\mathbf{J}\dot{\mathbf{v}}_{1} + 2\alpha k\sigma \mathbf{J}\mathbf{w}_{1} - \alpha\sigma^{2}\mathbf{v}_{1} + 2\gamma \mathbf{u}_{0}\mathbf{u}_{0}^{T}\mathbf{v}_{1} = \mathbf{0},$$

$$\mathbf{J}\dot{\mathbf{w}}_{1} - 2\alpha k\sigma \mathbf{J}\mathbf{v}_{1} - \alpha\sigma^{2}\mathbf{w}_{1} + 2\gamma \mathbf{u}_{0}\mathbf{u}_{0}^{T}\mathbf{w}_{1} = \mathbf{0}.$$
(10)

To determine the spectrum, let $(\mathbf{v}_1, \mathbf{w}_1) = (\mathbf{q}, \mathbf{p})e^{\lambda t}$. Then, (λ, σ) are determined by roots of

$$\Delta(\lambda,\sigma) = \lambda^4 + 2(p^2 + 4k^2\alpha^2\sigma^2)\lambda^2 + (p^2 - 4k^2\alpha^2\sigma^2)^2,$$

where $p^2 = \alpha^2 \sigma^4 - 2\alpha \gamma \|\mathbf{u}_0\|^2 \sigma^2$. Suppose $p^2 > 0$; then all four roots are purely imaginary (see Fig. 1) and are given by

 $\lambda = i2\alpha k\sigma \pm ip$ and $\lambda = -i2\alpha k\sigma \pm ip$.

These modes are purely imaginary as long as $p^2 > 0$; equivalently $2\gamma \alpha \|\mathbf{u}_0\|^2 < \alpha^2 \sigma^2$. Since $\alpha \gamma > 0$, the instability threshold is achieved when the amplitude reaches

$$\|\mathbf{u}_0\| = \frac{|\alpha\sigma|}{\sqrt{2\alpha\gamma}}.$$
(11)

At this threshold, a collision of eigenvalues occurs at the points $\lambda = \pm 2ik\alpha\sigma$; see Fig. 1 for a schematic of this collision.

It will be assumed henceforth that $k \neq 0$. Instability is, then, through a collision of eigenvalues of opposite energy signs, which reproduces the instability mechanism for the full water-wave problem.

Purely imaginary eigenvalues of a Hamiltonian system have a signature associated with them, and this signature is related to the sign of the energy.^{17,29,30} Collision of eigenvalues of opposite signatures is a necessary condition for the collision resulting in instability.

It is straightforward to compute the signature of the modes in the NLS model. Suppose that the amplitude $\|\mathbf{u}_0\|$ of the basic state is smaller than the critical value (11) for instability. There are, then, two pairs of purely imaginary eigenvalues, and they each have a signature. Let us concentrate on the eigenvalues on the positive imaginary axis:

$$\lambda = i\Omega_{\pm}$$
 with $\Omega_{\pm} = c_g \sigma \pm p$, $c_g = 2\alpha k$. (12)

Then,

$$\operatorname{sign}(\Omega_{\pm}) = \mathrm{i} \langle \overline{\mathbf{q}}, \mathbf{J} \mathbf{q} \rangle + \mathrm{i} \langle \overline{\mathbf{p}}, \mathbf{J} \mathbf{p} \rangle,$$

where the inner product is real in order to make the conjugation explicit. One can also show that this signature has the same sign as the energy perturbation restricted to this mode. A straightforward calculation shows that $sign(\Omega_+)=\pm 4$, when $\|\mathbf{u}_0\| = 0$. Since p^2 decreases as the amplitude increases, the two modes will have opposite signatures for all $\|\mathbf{u}_0\|$ between $\|\mathbf{u}_0\|=0$ and the point of collision.

Now consider the effect of the damping terms. Consider the reduced system (10) for the BF instability with the *abc*-damping terms included:

$$\mathbf{J}\dot{\mathbf{v}}_{1} + 2\alpha k\sigma \mathbf{J}\mathbf{w}_{1} - \alpha\sigma^{2}\mathbf{v}_{1} + 2\gamma \mathbf{u}_{0}\mathbf{u}_{0}^{T}\mathbf{v}_{1} + \mathcal{D}_{1} = \mathbf{0},$$

$$\mathbf{J}\dot{\mathbf{w}}_{1} - 2\alpha k\sigma \mathbf{J}\mathbf{v}_{1} - \alpha\sigma^{2}\mathbf{w}_{1} + 2\gamma \mathbf{u}_{0}\mathbf{u}_{0}^{T}\mathbf{w}_{1} + \mathcal{D}_{2} = \mathbf{0},$$
(13)

with

$$\mathcal{D}_{1} = 2ka\sigma\mathbf{w}_{1} + a\sigma^{2}\mathbf{J}\mathbf{v}_{1} + b\mathbf{J}\mathbf{v}_{1} + 2c\langle\mathbf{u}_{0},\mathbf{v}_{1}\rangle\mathbf{J}\mathbf{u}_{0}$$

$$(14)$$

$$\mathcal{D}_{2} = -2ka\sigma\mathbf{v}_{1} + a\sigma^{2}\mathbf{J}\mathbf{w}_{1} + b\mathbf{J}\mathbf{w}_{1} + 2c\langle\mathbf{u}_{0},\mathbf{w}_{1}\rangle\mathbf{J}\mathbf{u}_{0}.$$

Now, let $[\mathbf{v}_1(t), \mathbf{w}_1(t)] = (\tilde{\mathbf{v}}_1, \tilde{\mathbf{w}}_1)e^{\lambda t}$. The eigenvalue problem for the stability exponent then reduces to studying the roots of a determinant showing (with the help of MAPLE) that the two roots in the upper half-plane are given by

$$\lambda_{\pm} = 2\mathbf{i}k\sigma\alpha - (b + a\sigma^2 + c\|\mathbf{u}_0\|^2) \pm \mathbf{i}\sqrt{S},\tag{15}$$

with

$$\begin{split} S &= 4\mathbf{i}a\sigma^3k\alpha - c^2 \|\mathbf{u}_0\|^4 - 4k^2\sigma^2a^2 - 4\mathbf{i}a\sigma k\gamma \|\mathbf{u}_0\|^2 \\ &- 2\alpha\gamma\sigma^2 \|\mathbf{u}_0\|^2 + \alpha^2\sigma^4. \end{split}$$

When a=b=c=0, these stability exponents reduce to

 $\lambda_{\pm} = 2ik\sigma\alpha \pm i\sqrt{\alpha^2\sigma^4 - 2\alpha\gamma\sigma^2 \|\mathbf{u}_0\|^2}.$

Now suppose these two eigenvalues are purely imaginary: the amplitude $\|\mathbf{u}_0\|$ is below the critical value (11). To determine the leading order effect of dissipation, expand (15) in a Taylor series with respect to *a*, *b*, and *c* and take the real part

$$\operatorname{Re}(\lambda_{\pm}) = - (a\sigma^{2} + b + c \|\mathbf{u}_{0}\|^{2}) \mp \frac{2ak\sigma(\alpha\sigma^{2} - \gamma \|\mathbf{u}_{0}\|^{2})}{\sqrt{\alpha^{2}\sigma^{4} - 2\alpha\gamma\sigma^{2}\|\mathbf{u}_{0}\|^{2}}} + \cdots$$
(16)

For any a > 0, there is an open region of parameter space where these two real parts have opposite signs since their product to leading order is

$$\operatorname{Re}(\lambda_{-})\operatorname{Re}(\lambda_{+}) = (a\sigma^{2} + b + c \|\mathbf{u}_{0}\|^{2})^{2}$$
$$- \frac{4a^{2}k^{2}\sigma^{2}(\alpha\sigma^{2} - \gamma \|\mathbf{u}_{0}\|^{2})^{2}}{\alpha^{2}\sigma^{4} - 2\alpha\gamma\sigma^{2}\|\mathbf{u}_{0}\|^{2}} + \cdots$$

For any a, b, c with $a \neq 0$, there is an open set of values of $||\mathbf{u}_0||$, where this expression is strictly negative, showing that $\operatorname{Re}(\lambda_{-})$ and $\operatorname{Re}(\lambda_{+})$ perturb in opposite directions. In this parameter regime the dissipation perturbs the negative energy mode, as shown schematically in Fig. 2.

It is clear that when only the *b*-term is present, all eigenvalues shift to the left. Therefore, the *b*-term does not produce any enhancement of the instability, in agreement with Ref. 12. This damping is analogous to the *uniform damping* in (4). It is the *a*-term that leads to enhancement. However, the NLS is a simplified model for water waves.

In summary, the fundamental observation is that BF instability is associated with a collision of eigenvalues of posi-

Downloaded 27 Mar 2008 to 128.95.176.16. Redistribution subject to AIP license or copyright; see http://pof.aip.org/pof/copyright.jsp

tive and negative energy, and there are physically realizable forms of damping which enhance this instability. It remains to be seen how this effect can be revealed in laboratory experiments, in numerical experiments based on the full waterwave equations, and in the open ocean.

This work was enhanced by a grant of a CNRS Fellowship to the first author, and by support from the CMLA at Ecole Normale Supérieure de Cachan. Helpful discussions with Gianne Derks are gratefully acknowledged.

- ¹O. M. Phillips, "On the dynamics of unsteady gravity waves of finite amplitude. I. The elementary interactions," J. Fluid Mech. 9, 193 (1960).
 ²K. Hasselmann, "On the non-linear energy transfer in a gravity-wave spec-
- trum. Part 1. General theory," J. Fluid Mech. **12**, 481 (1962). ³M. J. Lighthill, "Contributions to the theory of waves in nonlinear disper-
- sive systems," J. Inst. Math. Appl. 1, 269 (1965). ⁴G. B. Whitham, "A general approach to linear and nonlinear dispersive
- waves using a Lagrangian," J. Fluid Mech. **22**, 273 (1965). ⁵T. B. Benjamin and J. E. Feir, "The disintegration of wavetrains in deep
- water. Part 1," J. Fluid Mech. 27, 417 (1967).
- ⁶T. B. Benjamin, "Instability of periodic wavetrains in nonlinear dispersive systems," Proc. R. Soc. London, Ser. A **299**, 59 (1967).
- ⁷J. C. R. Hunt, "Thomas Brooke Benjamin 15 April 1929–16 August 1995," Biogr. Mem. Fellows R. Soc. **49**, 39 (2003).
- ⁸P. J. Blennerhassett, "On the generation of waves by wind," Philos. Trans. R. Soc. London, Ser. A **298**, 451 (1980).
- ⁹L. F. Bliven, N. E. Huang, and S. R. Long, "Experimental study of the influence of wind on Benjamin-Feir sideband instability," J. Fluid Mech. 162, 237 (1986).
- ¹⁰A. Davey, "The propagation of a weak nonlinear wave," J. Fluid Mech. **53**, 769 (1972).
- ¹¹A. L. Fabrikant, "Nonlinear dynamics of wave packets in a dissipative medium," Sov. Phys. JETP **59**, 274 (1984).
- ¹²H. Segur, D. Henderson, J. Carter, J. Hammack, C.-M. Li, D. Pheiff, and K. Socha, "Stabilizing the Benjamin-Feir instability," J. Fluid Mech. **539**, 229 (2005).

- ¹³G. Wu, Y. Liu, and D. K. P. Yue, "A note on stabilizing the Benjamin-Feir instability," J. Fluid Mech. 556, 45 (2006).
- ¹⁴T. J. Bridges and A. Mielke, "A proof of the Benjamin-Feir instability," Arch. Ration. Mech. Anal. 133, 145 (1995).
- ¹⁵L. A. Ostrovskiĭ, "Propagation of wave packets and space-time selffocusing in a nonlinear medium," Sov. Phys. JETP 24, 797 (1967).
- ¹⁶V. E. Zakharov, "Stability of periodic waves of finite amplitude on the surface of a deep fluid," J. Appl. Mech. Tech. Phys. 9, 190 (1968).
- ¹⁷R. A. Cairns, "The role of negative energy waves in some instabilities of parallel flows," J. Fluid Mech. **92**, 1 (1979).
- ¹⁸A. D. D. Craik, *Wave Interactions and Fluid Flows* (Cambridge University Press, Cambridge, 1988).
- ¹⁹R. S. MacKay, "Movement of eigenvalues of Hamiltonian equilibria under non-Hamiltonian perturbation," Phys. Lett. A **155**, 266 (1991).
- ²⁰L. A. Ostrovskiĭ, S. A. Rybak, and L. Sh. Tsimring, "Negative energy waves in hydrodynamics," Sov. Phys. Usp. **29**, 1040 (1986).
- ²¹A. L. Fabrikant and Yu. A. Stepanyants, *Propagation of Waves in Shear Flows* (World Scientific, Singapore, 1998).
- ²²T. B. Benjamin, "Classification of unstable disturbances in flexible surfaces bounding inviscid flows," J. Fluid Mech. **16**, 436 (1963).
- ²³M. T. Landahl, "On the stability of a laminar incompressible boundary layer over a flexible surface," J. Fluid Mech. **13**, 609 (1962).
- ²⁴N. Peake, "Nonlinear stability of fluid-loaded elastic plate with mean flow," J. Fluid Mech. **434**, 101 (2001).
- ²⁵G. Derks and T. Ratiu, "Attracting curves on families of stationary solutions in two-dimensional Navier-Stokes and reduced magnetohydrodynamics," Proc. R. Soc. London, Ser. A **454**, 1407 (2002); "Unstable manifolds of relative equilibria in Hamiltonian systems with dissipation," Nonlinearity **15**, 531 (1998).
- ²⁶W. Craig, C. Sulem, and P. L. Sulem, "Nonlinear modulation of gravity waves: a rigorous approach," Nonlinearity 5, 497 (1992).
- ²⁷M. C. Longuet-Higgins, "Integral properties of periodic gravity waves of finite amplitude," Proc. R. Soc. London, Ser. A **342**, 157 (1975).
- ²⁸D. Dutykh and F. Dias, "Viscous potential free-surface flows in a fluid layer of finite depth," C. R. Acad. Sci., Ser. I: Math. **345**, 113 (2007).
- ²⁹R. S. MacKay and P. G. Saffman, "Stability of water waves," Proc. R. Soc. London, Ser. A **406**, 115 (1986).
- ³⁰T. J. Bridges, "A geometric formulation of the conservation of wave action and its implications for signature and the classification of instabilities," Proc. R. Soc. London, Ser. A **453**, 1365 (1997).