North Atlantic winter cyclones starve seabirds

Highlights

- Five key North Atlantic seabird species are exposed to winter cyclones
- High exposure is in the Labrador Sea, the Davis Strait, the Barents Sea, and off Iceland
- Seabird energy requirements do not seem to increase under cyclonic conditions
- Seabird mortality during high-intensity cyclones is likely caused by starvation

Authors

Manon Clairbaux, Paul Mathewson, Warren Porter, ..., Anders Mosbech, Aevar Petersen, David Grémillet

Correspondence

clairbauxm@gmail.com (M.C.), david.gremillet@cebc.cnrs.fr (D.G.)

In brief

Using tracking data for >1,500 migration seabirds from five key North Atlantic species, cyclone locations, and bioenergetics modeling, Clairbaux et al. demonstrate that high-intensity winter cyclones impact birds from all studied species and breeding colonies and suggest that cyclonic conditions starve seabirds by preventing them from feeding.





Report

North Atlantic winter cyclones starve seabirds

Manon Clairbaux,^{1,31,*} Paul Mathewson,² Warren Porter,² Jérôme Fort,³ Hallvard Strøm,⁴ Børge Moe,⁵ Per Fauchald,⁵ Sebastien Descamps,⁴ Hálfdán H. Helgason,⁴ Vegard S. Bråthen,⁵ Benjamin Merkel,⁶ Tycho Anker-Nilssen,⁵ Ingar S. Bringsvor,⁷ Olivier Chastel,⁸ Signe Christensen-Dalsgaard,⁵ Jóhannis Danielsen,⁹ Francis Daunt,¹⁰ Nina Dehnhard,⁵ Kjell Einar Erikstad,^{11,12} Alexey Ezhov,¹³ Maria Gavrilo,^{14,15} Yuri Krasnov,¹³ Magdalene Langset,⁵ Svein-H. Lorentsen,⁵ Mark Newell,¹⁰ Bergur Olsen,⁹ Tone K. Reiertsen,¹¹ Geir Helge Systad,⁵ Thorkell L. Thórarinsson,¹⁶ Mark Baran,¹⁷ Tony Diamond,¹⁷ Annette L. Fayet,¹⁸ Michelle G. Fitzsimmons,¹⁹ Morten Frederiksen,²⁰ Hugh G. Gilchrist,²¹

(Author list continued on next page)

¹CEFE, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul Valéry Montpellier 3, Montpellier, France

- ²Department of Integrative Biology, University of Wisconsin, Madison, WI, USA
- ³Littoral, Environnement et Sociétés (LIENSs), UMR7266 CNRS La Rochelle Université, 2 rue Olympe de Gouges, 17000 La Rochelle, France ⁴Norwegian Polar Institute, Fram Centre, PO Box 6606 Langnes, 9296 Tromsø, Norway
- ⁵Norwegian Institute for Nature Research NINA, PO Box 5685 Torgarden, 7485 Trondheim, Norway
- ⁶Akvaplan-niva AS, Fram Centre, PO Box 6606 Langnes, 9296 Tromsø, Norway
- ⁷Norwegian Ornithological Society, Sandgata 30 B, 7012 Trondheim, Norway
- ⁸Centre d'Etudes Biologiques de Chizé (CEBC), UMR 7372 CNRS/La Rochelle Univ, La Rochelle, France
- ⁹Faroe Marine Research Institute, PO Box 3051, Nóatún 1, 110 Tórshavn, Faroe Islands
- ¹⁰UK Centre for Ecology & Hydrology, Bush Estate, Penicuik EH26 0QB, UK
- ¹¹Norwegian Institute for Nature Research, Fram Centre, 9296 Tromsø, Norway
- ¹²Centre for Biodiversity Dynamics, Department of Biology, Norwegian University of Science and Technology, 7491 Trondheim, Norway

¹³Murmansk Marine Biological Institute, 17 Vladimirskaya Street, 183010 Murmansk, Russia

- ¹⁴Association Maritime Heritage, Saint Petersburg, Russia
- ¹⁵National Park Russian Arctic, 57 Sovetskikh Kosmonavtove Avenue, Archangelsk, Russia
- ¹⁶Northeast Iceland Nature Research Centre, Hafnarstétt 3, 640 Húsavík, Iceland
- ¹⁷Atlantic Laboratory for Avian Research, University of New Brunswick, PO Box 4400, Fredericton, NB E3B 5A3, Canada
- ¹⁸Department of Zoology, University of Oxford, 11a Mansfield Road, Oxford OX1 3SZ, UK
- ¹⁹Wildlife Research Division, Environment and Climate Change Canada, 6 Bruce Street, Mount Pearl, NL A1N 4T3, Canada

(Affiliations continued on next page)

SUMMARY

Each winter, the North Atlantic Ocean is the stage for numerous cyclones, the most severe ones leading to seabird mass-mortality events called "winter wrecks."¹⁻³ During these, thousands of emaciated seabird carcasses are washed ashore along European and North American coasts. Winter cyclones can therefore shape seabird population dynamics^{4,5} by affecting survival rates as well as the body condition of surviving individuals and thus their future reproduction. However, most often the geographic origins of impacted seabirds and the causes of their deaths remain unclear.⁶ We performed the first ocean-basin scale assessment of cyclone exposure in a seabird community by coupling winter tracking data for \sim 1,500 individuals of five key North Atlantic seabird species (Alle alle, Fratercula arctica, Uria aalge, Uria lomvia, and Rissa tridactyla) and cyclone locations. We then explored the energetic consequences of different cyclonic conditions using a mechanistic bioenergetics model⁴ and tested the hypothesis that cyclones dramatically increase seabird energy requirements. We demonstrated that cyclones of high intensity impacted birds from all studied species and breeding colonies during winter but especially those aggregating in the Labrador Sea, the Davis Strait, the surroundings of Iceland, and the Barents Sea. Our broad-scale analyses suggested that cyclonic conditions do not increase seabird energy requirements, implying that they die because of the unavailability of their prey and/or their inability to feed during cyclones. Our study provides essential information on seabird cyclone exposure in a context of marked cyclone regime changes due to global warming.⁸

RESULTS

Cyclone-induced seabird mortality adds to other threats, such as habitat loss, invasive species, or interactions with fisheries, contributing to the general downward trend of the global seabird community⁹ and making them one of the most threatened bird groups.¹⁰ It is therefore essential to understand the causes of seabird mortality during cyclones and to identify factors



Report

Tim Guilford,¹⁸ Nicholas P. Huffeldt,^{20,22} Mark Jessopp,^{23,24} Kasper L. Johansen,²⁰ Amy-Lee Kouwenberg,²⁵ Jannie F. Linnebjerg,²⁰ Heather L. Major,²⁶ Laura McFarlane Tranquilla,²⁵ Mark Mallory,²⁷ Flemming R. Merkel,²⁰ William Montevecchi,²⁸ Anders Mosbech,²⁰ Aevar Petersen,²⁹ and David Grémillet^{8,30,*} ²⁰Aarhus University, Department of Bioscience, Frederiksborgvej 399, 4000 Roskilde, Denmark ²¹National Wildlife Research Centre, Environment and Climate Change Canada, Ottawa, ON, Canada

²²Greenland Institute of Natural Resources, Kivioq 2, 3900 Nuuk, Greenland

²³School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Field, North Mall, Cork, Ireland

²⁴MaREI Centre, Environmental Research Inst., Univ. College Cork, Cork, Ireland

²⁵Bird Studies Canada, Sackville, NB, Canada

²⁶Department of Biological Sciences, University of New Brunswick, PO Box 5050, Saint John, NB E2L 4L5, Canada

²⁷Biology, Acadia University, 15 University Avenue, Wolfville, NS B4P 2R6, Canada

²⁸Psychology and Biology Departments, Memorial University of Newfoundland, St. John's, NL A1C 5S7, Canada
²⁹Brautarland 2, 108 Reykjavik, Iceland

³⁰Percy FitzPatrick Institute, DST/NRF Centre of Excellence, University of Cape Town, Rondebosch, South Africa ³¹Lead contact

*Correspondence: clairbauxm@gmail.com (M.C.), david.gremillet@cebc.cnrs.fr (D.G.) https://doi.org/10.1016/j.cub.2021.06.059

contributing to contrasting spatiotemporal mortality patterns at the seabird-community level. Multiple seabird species from different breeding colonies may inhabit different wintering areas,^{11,12} and this may strongly affect the composition of seabird winter wrecks. In this context, community-wide analyses are lacking due to the technical difficulties of studying seabirds wintering far out at sea under harsh conditions.

Winter cyclone exposure of the North Atlantic seabird community

The North Atlantic Ocean and adjacent seas (100°W-100°E, 30°N-90°N; see Figure S1) is the stage for numerous winter cyclones (October to February; see Figure S2), which are defined by the National Hurricane Center as large-scale air masses that rotate around a center of low atmospheric pressure. Their locations between 2000 and 2016 were obtained from climatic reanalysis data (see STAR Methods). Using the Dvorak storm classification.¹³ we sorted them across their duration into one of the four classes of cyclone intensity defined (see STAR Methods). Calculating, for each month, the average number of cyclones of each category for each 250-km grid cell, we demonstrated that a limited number of localized low-intensity cyclones (class 1 and 2) occurred mainly in the Baffin Bay, the Gulf of Maine, and the Mediterranean Sea regions (see Data S1A). Class 3 cyclones were more numerous and widespread, particularly in October and November, and mainly occurred in the Baffin and Hudson bays, the Davis Strait, the Labrador Sea, east off Newfoundland, around Iceland, and in the Barents Sea. Finally, class 4 cyclones were observed in all winter months off west Iceland, the Norwegian coast, and in the Barents Sea, whereas the Davis Strait and the Labrador Sea were only affected by class 4 cyclones between December and February.

Winter bird locations were obtained from global location sensors (GLSs) deployed and retrieved during the breeding season on 1,532 individuals from 39 breeding colonies across the North Atlantic Ocean (see STAR Methods and Table S1). We approximated core use areas for each colony and species during the winter months, calculating monthly utilization distributions (UDs) with the BRB-MKDE software (see STAR Methods).¹⁴ To conclude on seabird cyclone exposure, we overlapped cyclone locations with core wintering areas of different seabird species and populations (see Figure 1 and Data S1A–S1F) before calculating an average cyclone exposure index (see Figure 2 and STAR Methods).

Low-intensity cyclones (class 1 and 2) showed limited overlap with the seabird community. However, birds from all species and colonies were likely to encounter class 3 and 4 cyclones during winter, with substantial inter-annual variability in the mean number of such events occurring within seabird core use areas (Figure 1; Data S1B-S1F). Notably, seabirds from Canadian, Greenlandic, and Icelandic colonies were exposed to class 3 and 4 cyclones in the Davis Strait, the Labrador Sea, and off Newfoundland, although Norwegian and Russian seabirds were likely impacted by those occurring in the Barents Sea. In contrast, class 3 cyclones were fewer in the mid-west Atlantic Ocean and were more likely to spare seabirds wintering there (see Data S1A). Due to major seabird aggregation and the marked presence of class 3 and 4 cyclones, areas such as the Labrador Sea, the Davis Strait, the vicinity of Iceland, and the Barents Sea are predicted as sources of major winter wrecks (Figure 2).

Seabird energy requirements under cyclonic and noncyclonic conditions

Necropsies suggest that seabirds are lean after being exposed to high-intensity cyclones^{6,15} and that the resulting mortality can be aggravated by mercury contamination.¹⁶ Yet we do not know whether they starve due to abnormally high energy expenditure linked to harsh climatic conditions,¹⁷ because of reduced foraging profitability and energy inputs, or because both these constraints synoptically affect their energy balance. Recent research has shown that seabirds wintering in the North Atlantic Ocean track environmental gradients to remain in an energetic steady state,^{18–20} but how winter cyclones affect this delicate balance remains mysterious.

To test the hypothesis that cyclones dramatically increase seabird energy requirements, we modeled species-specific, monthly energy requirements for each winter between 2000 and 2016 on a 1,000 km × 1,000 km area off North Newfound-land (see Figure S1), using the mechanistic model Niche Mapper (see STAR Methods)⁷ under four intensities of cyclones and under non-cyclonic conditions (see STAR Methods). Following the protocol used by Grémillet and colleagues,²¹ we calculated how

CellPress



Figure 1. Average number of cyclones within the core use areas of common guillemot from different colonies between 2000 and 2016 in October

BJ, Bjørnøya; CG, Cape Gorodetskiy; FI, Faroe Islands; GR, Grimsey; HJ, Hjelmsøya; HO, Hornøya; JM, Jan Mayen; LaS, Langanes and Skjalfandi; LA, Latrabjarg; SK, Sklinna. Error bars correspond to standard deviations capturing the variation between years. Results for other species, colonies, and months are provided in Data S1B–S1F.

many days each of the five studied species could fast before dying, when exposed to class 2, class 3, and class 4 cyclones in the studied area (see STAR Methods). Statistical analyses revealed several significant differences (Kruskal-Wallis, p < 0.05) in seabird energy requirements (see Table 1) between the six categories of conditions tested (class 1-4 cyclones, non-cyclonic conditions with usual seabird flight and diving activities, and non-cyclonic conditions with cyclonic seabird flight and diving activities). In a very limited number of cases, cyclonic conditions led to increases (up to 36%) in seabird energy requirements in comparison with non-cyclonic conditions with cyclonic flight and diving activities (post hoc Dunn's test, p < 0.05). Surprisingly, we found that seabird energy requirements were generally similar or lower during cyclones, compared to non-cyclonic conditions (see Figure 3 and Data S1G-S1K). In particular, for little auks and black-legged kittiwakes, energy requirements were generally significantly lower during class 2, 3, and 4 cyclones than during non-cyclonic conditions experienced with usual flight and diving activities (post hoc Dunn's test, p < 0.05). Those differences were not significant for deep divers (except for Atlantic puffins in October and December under specific cyclonic conditions; see Table 1). Further, seabird energy requirements were not significantly different during class 1 cyclones than for non-cyclonic conditions experienced with usual flight and diving activities (post hoc Dunn's test, p < 0.05).

Energy requirements under class 2, class 3, and class 4 cyclones were not significantly different (see Data S1G–S1K), and these classes were therefore pooled when calculating fasting endurance. Between October and December, average fasting endurance was 2.2 days (\pm 0.1) for little auk and 6.5 days (\pm 2.5) for Atlantic puffins, although black-legged kittiwakes, Brünnich's guillemots, and common guillemots could fast 8.4 days (\pm 0.5), 7.3 days (\pm 0.06), and 8.1 days (\pm 0.1), respectively. With winter's advance, these durations dropped, respectively, to 1.6 days

(±0.1), 4.6 days (±0.6), 6.4 days (±0.5), 5.9 days (±0.6), and

6.3 days (±0.7) between January and February.

Current Biology

Report

DISCUSSION

We demonstrated that seabirds wintering in areas such as the Labrador Sea, the Davis Strait, the vicinity of Iceland, and the Barents Sea are particularly exposed to cyclones. Crucially, our broad-scale analyses led us to reject our hypothesis: climatic conditions experienced by seabirds during cyclones generally did not enhance their energy requirements. These results suggest that seabird mortality during winter high-intensity cyclones is likely caused by starvation due to the unavailability of prey and/ or their incapacity to feed.

Mechanistic insights into storms' impacts on seabirds

Our surprising conclusion regarding death causes in seabirds exposed to cyclones is in agreement with a study conducted on greater flamingos (Phoenicopterus roseus) during cold spells in the Camargue, France:22 thousands of flamingos died, not because of hypothermia but due to inaccessible food in frozen salt pans. Conversely, our results contrast with former modeling work that pointed to weather-induced energetic bottlenecks in seabirds wintering in the North Atlantic Ocean.¹⁷ Birds were predicted to experience markedly higher thermoregulatory costs in December, because of lower air and water temperatures and higher wind speeds. Yet this previous study investigated seabird energetics at the scale of months, whereas cyclones affect them at the scale of days. In this context, little auks appear as particularly vulnerable to fasting induced by cyclones, because of low lipid reserves and high mass-specific energy expenditure, likely explaining extensive and recurrent winter wrecks on North American coasts near their main wintering grounds.^{23,24} Other species were predicted to cope with relatively longer fasts but remain vulnerable to prolonged cyclonic conditions, especially at the end of winter. Therefore, recurrent cyclones affecting a specific area and seabird population are predicted as strong drivers of winter wrecks.

What exactly prevents seabirds from feeding during cyclonic conditions remains unclear. One possibility is that cyclones may enhance water turbidity, decrease underwater light intensity, and perturb prey patches and vertical migration. Potentially disrupting water stratification, cyclones may modify prey aggregation and negatively impact seabird foraging efficiency.²⁵ Therefore, this "washing-machine effect" could strongly affect seabirds feeding within the 50 upper meters of the water column, such as black-legged kittiwakes and little auks in our study. Further, cyclones and associated disturbances alter underwater soundscapes and thus constrain the detection of acoustic cues by seabirds, being from conspecifics, prey, or predators.^{26,27} Prey could also occur deeper in the water column during cyclones as already observed for some elasmobranchs or

Report





teleosts,^{28,29} but how this impacts pursuit-diving species, such as puffins and guillemots, is not known. By continuing to dive during cyclones, deep divers, such as guillemots and puffins, potentially take the risk of using energy reserves in an attempt to track inaccessible prey. However, common and Brünnich's guillemots have been observed switching their diet during strong winds, from schooling fish to amphipods or smaller sandeels, potentially targeting prey that required less underwater searching.^{30,31} Alternatively, cyclones may shift alcids away from their preferred prey patches and into unprofitable foraging habitats.³² Overall, even if starvation may be the main driver of seabird winter wrecks, we cannot exclude other causes of mortality, such as drowning,⁶ collision with reefs and rocky coastlines (M. Baran, personal communication), or inland stranding.³²

Potential limitations

Despite these advances, we recommend caution in interpreting our results, as they are based on reanalyzed environmental data and on thermodynamic modeling. Notably, measuring empirical data is nearly impossible during cyclones. This calls for the use of modeling approaches to characterize these climatic events and their energy consequences for seabirds but also constrains model outputs. This lack of in situ measurements influenced NicheMapper simulations in two main ways. First, the behavior of pelagic seabirds is hard to assess under cyclonic conditions, and the animal module mainly relies on the assumption that North Atlantic seabirds significantly reduce flying and/or diving during cyclones. On a worldwide scale, although many seabird species exhibit avoidance behavior, seeking coastal shelter and/or reducing activity levels during high-intensity cyclones,³³ others, such as great frigatebirds (Fregata minor), take advantage of cyclonic winds to save energy while gliding over thousands of kilometers.³⁴ Such extreme gliders, which occur in the tropics and Southern Ocean, are rarer in the North Atlantic, an area dominated by alcid species in terms of abundance.³⁵ This seabird family, which is morphologically adapted to using its wings for underwater propulsion, has high wing loading and very limited capacity to ride strong winds. Thus, cyclonic conditions dramatically increase flight costs in alcids³⁰ and therefore tend to prevent them from flying.

High Figure

Figure 2. Mean cyclone exposure index across the North Atlantic Ocean during winter

See STAR Methods for details on index calculation.

Second, because direct measurements of environmental data during cyclones were lacking at the spatiotemporal scales needed for our analyses, we depicted cyclonic conditions in the microclimate module using a limited number of reconstructed data at a broad scale, potentially missing other relevant differences between non-cyclonic and cyclonic conditions. Finally, due to a lack of available information, we only

modeled the effects of temperature, wind speed, and reduced activity on heat loss and energy requirements during cyclones, omitting other variables, such as wind-gust speed, water velocity, wave height, as well as other conditions potentially affecting seabird thermoregulation and thus their energy requirements. For example, localized wind gusts, sea spray, and high-speed precipitation could all reduce plumage insulation, thereby increasing seabird heat loss and energy requirements. Further, tall waves could require the birds to dive more frequently, increasing energy requirements. Thus, current predictions on seabird energy requirements during cyclones should be interpreted as likely conservative estimates. However, given the current lack of information, mechanistic models remain important tools to explore potential impacts and to generate hypotheses that could be further investigated. Additional direct measurements could help to refine these modeling approaches and evaluate initial assumptions. For example, deploving electronic devices to study North Atlantic seabird 3D movements during cyclones is a key objective for future research, which will allow testing our assumption that these animals are less active under passing cyclones. Such deployments will also help infer additional energetic costs in seabirds floating across rough seas, especially if they try to dive under each wave and/or to stay in the same area rather than drifting away with surface currents (up to 2.5 m.s⁻¹ under class 4 cyclones).³⁶ Finally, future efforts will allow researchers to take into account inter- and intra-individual morphological, physiological, and behavioral variability across winter in seabird energetic modeling, improving output accuracy.

Despite these limitations, our study provides essential new knowledge, notably on the vulnerability of wintering seabirds to high-intensity cyclones in the Labrador Sea, off Newfound-land, off southern Greenland, around Iceland, off the Norwe-gian coast, and in the Barents Sea. These results are supported by the numerous seabird winter wrecks observed across years on beaches close to those areas (F. Ravn Merkel, personal communication about Brünnich's guillemot wrecks in south Greenland).^{2,6,23,24,37} Yet as GLS tracking is based on individuals who survived the non-breeding period, we cannot exclude the possibility that wrecked individuals wintered in



Table 1. Results of Kruskal-Wallis tests

	Little auks			Black-legged kittiwakes			Atlantic puffins			Common guillemots			Brünnich's guillemots		
	χ^2	df	р	χ ²	df	р	χ ²	df	р	χ ²	df	р	χ^2	df	р
October	48.6	5	<0.0001 ^a	50.4	5	<0.0001 ^a	12.5	5	0.029 ^a	21.2	5	<0.0001 ^a	6.98	5	0.223
November	48.8	5	<0.0001 ^a	44.5	5	<0.0001 ^a	10.1	5	0.072	8.70	5	0.122	9.98	5	0.076
December	38.6	5	<0.0001 ^a	36.9	5	<0.0001 ^a	26.7	5	<0.0001 ^a	7.28	5	0.201	10.16	5	0.071
January	26.1	5	<0.0001 ^a	25.8	5	<0.0001 ^a	12.9	5	0.024 ^a	12.2	5	0.032 ^a	14.4	5	0.013 ^a
February	14.0	5	0.016 ^a	11.5	5	0.042	17.9	5	0.003 ^a	9.99	5	0.076	10.8	5	0.055

Comparisons were made between mean energy requirements per conditions tested (class 1–4 cyclones, non-cyclonic conditions with usual seabird flight and diving activities, and non-cyclonic conditions with cyclonic seabird flight and diving activities) for each species and each month. Outputs of the corresponding Dunn post hoc test are presented in Data S1G–S1K. χ 2, chi square; df., degrees of freedom; p, p value. ^aStatistically significant results (p < 0.05).

different locations with higher cyclone exposure, even though seabirds are highly gregarious, also at sea. Further, the localization and study of winter wrecks on beaches is impacted by oceanic currents carrying seabird carcasses and by the accessibility of these beaches to observers. This potentially leads to spatiotemporal mismatches between seabird wrecks as perceived by researchers and the general public and areas of major winter high-intensity cyclone impacts on seabirds offshore. Such a mismatch may explain why we found limited overlap between seabird winter distributions and cyclones in the Eastern Atlantic, despite the fact that large winter wrecks have regularly been recorded along the shores of Western Europe.

Seabird storm exposure under climate change

It is well established that seabirds experience direct and indirect negative stressors from global warming.³⁸ Since the 1970s, the frequency and intensity of the strongest tropical cyclones in

the North Atlantic have increased markedly.³⁹ Considering future North Atlantic cyclone regimes, there are still uncertainties, but model outputs predict some common features: there will be fewer cyclones in the future, but the frequency of the strongest ones is predicted to increase with global warming.³⁹ Moreover, cyclone tracks will likely shift northward under climate change,⁴⁰ increasing the storminess of Western Europe⁴¹ and of the high Arctic.^{42,43} Because storm activity is positively correlated to the magnitude of seabird wrecks^{44,45} and the North Atlantic Ocean seabird community is also predicted to shift northward, following its prey base,⁴⁶ we infer that this community will become even more susceptible to mass mortality caused by winter wrecks. This might be particularly marked in areas such as the Bay of Biscay, the North Sea, and the Norwegian and the Barents Seas. In this context, our study provides an essential conceptual and methodological framework to identify the vulnerability of specific populations to cyclone events at the scale of the North Atlantic Ocean.



Figure 3. Average energy requirements for little auks and Atlantic puffins in the studied area off Newfoundland for each wintering month under cyclonic and non-cyclonic conditions between 2000 and 2016

Error bars were halved for clarity reasons and correspond to standard deviations capturing between-year variation. Shown are little auks (A) and Atlantic puffins (B). Results for other species are presented in Data S1G–S1K.



STAR*METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- RESOURCE AVAILABILITY
 - Lead contact
 - Materials availability
 - Data and code availability
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
- METHOD DETAILS
 - Locations and characteristics of North Atlantic cyclones during winter
 - Winter distribution of seabirds
 - Modeling seabird energy requirements under noncyclonic and cyclonic conditions
 - Estimation of fasting endurance
- QUANTIFICATION AND STATISTICAL ANALYSIS

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at https://doi.org/10.1016/j. cub.2021.06.059.

ACKNOWLEDGMENTS

We warmly thank the many fieldworkers involved in this study. Data collection in Norway, including Svalbard and Jan Mayen, was supported by the Norwegian seabird program SEAPOP (https://seapop.no/en). Thanks to the SEATRACK program for making tracking data available for the analyses (http://seatrack. seapop.no/map/). Fieldwork on little auks in Northwest Greenland was funded by the Bureau of Minerals and Petroleum, Greenland Government. The work on the Isle of May was also supported by the Natural Environment Research Council (award NE/R016429/1 as part of the UK-SCaPE programme delivering National Capability). The New Brunswick Wildlife Trust Fund provided funding for GLS tags on Machias Seal Island. We thank the University of New Brunswick, the Federal Government of Canada, the Government of Greenland, and the Danish Ministry of Environment and Food. Thanks to the Polar Continental Shelf Program of Natural Resources Canada (grant 213-08) and core funding from Environment and Climate Change Canada. Thanks also to the French Polar Institute Paul Emile Victor for funding the ADACLIM program (IPEV program 388 to D.G. and J.F.) and the ORNITHO-ENDOCRINO program (IPEV program 330 to O.C.). We thank Simon Benhamou and Benjamin Robira for their help with BRB-MKDE; Doris Gomez, Christine Andraud, and Aurélie Tournié for their valuable help regarding feather reflectivity measurements; and Katherine Brownlie for carcass measurements. We thank Kyle Elliott (McGill University, Canada) and Sarah Wanless (Center for Ecology and Hydrology, UK) for their help when assessing the seabird behavior during cyclones and Antoine Chabrolle for our discussion regarding winter wreck phenomena. This is a contribution to the Excellence Chair ECOMM funded by the Region Nouvelle Aquitaine.

AUTHOR CONTRIBUTIONS

Conceived and designed the project, D.G., J.F., and M.C.; software developers and data providers, W.P., P.M., J.F., H.S., B. Moe, P.F., S.D., H.H.H., V.S.B., B. Merkel, T.A.-N., I.S.B., O.C., S.C.-D., J.D., F.D., N.D., K.E.E., A.E., M.G., Y.K., M.L., S.-H.L., M.N., B.O., T.K.R., G.H.S., T.L.T., M.B., T.D., A.L.F., M.G.F., M.F., H.G.G., T.G., N.P.H., M.J., K.L.J., A.-L.K., J.F.L., L.M.T., M.M., F.R.M., W.M., A.M., A.P., and D.G.; analyzed the data, M.C.; wrote the paper, M.C. and D.G. All authors reviewed the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: February 18, 2021 Revised: April 26, 2021 Accepted: June 21, 2021 Published: July 29, 2021

REFERENCES

- Anker-Nilssen, T., Harris, M.P., Kleven, O., and Langset, M. (2017). Status, origin, and population level impacts of Atlantic Puffins killed in a mass mortality event in southwest Norway early 2016. Seabird 30, 1–14.
- Camphuysen, C.J., Wright, P.J., Leopold, M., Hüppop, O., and Reid, J.B. (1999). A review of the causes, and consequences at the population level, of mass mortality of seabirds. ICES Coop. Res. Rep. 232, 51–56.
- Farque, P.-A. (2014). Échouage massif d'oiseaux marins durant l'hiver 2014 sur la façade Atlantique (LPO France).
- Frederiksen, M., Daunt, F., Harris, M.P., and Wanless, S. (2008). The demographic impact of extreme events: stochastic weather drives survival and population dynamics in a long-lived seabird. J. Anim. Ecol. 77, 1020–1029.
- Guéry, L., Descamps, S., Hodges, K.I., Pradel, R., Moe, B., Hanssen, S.A., Erikstad, K.E., Gabrielsen, G.W., Gilchrist, H.G., Jenouvrier, S., et al. (2019). Winter extratropical cyclone influence on seabird survival: variation between and within common eider *Somateria mollissima* populations. Mar. Ecol. Prog. Ser. 627, 155–170.
- Morley, T.I., Fayet, A.L., Jessop, H., Veron, P., Veron, M., Clark, J., and Wood, M.J. (2016). The seabird wreck in the Bay of Biscay and Southwest Approaches in 2014: a review of reported mortality. Seabird 29, 22–38.
- Porter, W.P., and Mitchell, J.W. (2006). Method and system for calculating the spatial-temporal effects of climate and other environmental conditions on animals. US patent US7155377B2, filed August 23, 2001, and granted December 26, 2006.
- Christensen, J.H., Kanikicharla, K.K., Aldrian, E., An, S.-I., Albuquerque Cavalcanti, I.F., de Castro, M., Dong, W., Goswami, P., Hall, A., Kanyanga, J.K., et al. (2013). Climate phenomena and their relevance for future regional climate change. In Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley, eds. (Cambridge University).
- Grémillet, D., Ponchon, A., Paleczny, M., Palomares, M.D., Karpouzi, V., and Pauly, D. (2018). Persisting worldwide seabird-fishery competition despite seabird community decline. Curr. Biol. 28, 4009–4013.e2.
- Dias, M.P., Martin, R., Pearmain, E.J., Burfield, I.J., Small, C., Phillips, R.A., Yates, O., Lascelles, B., Borboroglu, P.G., and Croxall, J.P. (2019). Threats to seabirds: a global assessment. Biol. Conserv. 237, 525–537.
- Frederiksen, M., Descamps, S., Erikstad, K.E., Gaston, A.J., Gilchrist, H.G., Grémillet, D., Johansen, K.L., Kolbeinsson, Y., Linnebjerg, J.F., Mallory, M.L., et al. (2016). Migration and wintering of a declining seabird, the thick-billed murre *Uria lomvia*, on an ocean basin scale: conservation implications. Biol. Conserv. 200, 26–35.
- 12. Fayet, A.L., Freeman, R., Anker-Nilssen, T., Diamond, A., Erikstad, K.E., Fifield, D., Fitzsimmons, M.G., Hansen, E.S., Harris, M.P., Jessopp, M., et al. (2017). Ocean-wide drivers of migration strategies and their influence on population breeding performance in a declining seabird. Curr. Biol. 27, 3871–3878.e3.
- 13. Dvorak, V. (1984). Tropical cyclone intensity analysis using satellite date. NOAA Tech. Rep. NESDIS 11.
- 14. Benhamou, S. (2011). Dynamic approach to space and habitat use based on biased random bridges. PLoS ONE 6, e14592.
- Jones, P.H., Barrett, C.F., Mudge, G.P., and Harris, M.P. (1984). Physical condition of auks beached in eastern Britain during the wreck of February 1983. Bird Study 31, 95–98.

CellPress

- Fort, J., Lacoue-Labarthe, T., Nguyen, H.L., Boué, A., Spitz, J., and Bustamante, P. (2015). Mercury in wintering seabirds, an aggravating factor to winter wrecks? Sci. Total Environ. 527-528, 448–454.
- Fort, J., Porter, W.P., and Grémillet, D. (2009). Thermodynamic modelling predicts energetic bottleneck for seabirds wintering in the northwest Atlantic. J. Exp. Biol. 212, 2483–2490.
- Amélineau, F., Fort, J., Mathewson, P.D., Speirs, D.C., Courbin, N., Perret, S., Porter, W.P., Wilson, R.J., and Grémillet, D. (2018). Energyscapes and prey fields shape a North Atlantic seabird wintering hotspot under climate change. R. Soc. Open Sci. 5, 171883.
- Fort, J., Beaugrand, G., Grémillet, D., and Phillips, R.A. (2012). Biologging, remotely-sensed oceanography and the continuous plankton recorder reveal the environmental determinants of a seabird wintering hotspot. PLoS ONE 7, e41194.
- 20. Clairbaux, M., Cheung, W.W.L., Mathewson, P., Porter, W., Courbin, N., Fort, J., Strøm, H., Moe, B., Fauchald, P., Descamps, S., et al. (2021). Meeting Paris agreement objectives will temper seabird winter distribution shifts in the North Atlantic Ocean. Glob. Chang. Biol. 27, 1457–1469.
- Grémillet, D., Kuntz, G., Woakes, A.J., Gilbert, C., Robin, J-P., Le Maho, Y., et al. (2005). Year-round recordings of behavioural and physiological parameters reveal the survival strategy of a poorly insulated diving endotherm during the Arctic winter. Journal of Experimental Biology 208, 4231– 4241. https://doi.org/10.1242/jeb.01884.
- 22. Deville, A.-S., Labaude, S., Robin, J.-P., Béchet, A., Gauthier-Clerc, M., Porter, W., Fitzpatrick, M., Mathewson, P., and Grémillet, D. (2014). Impacts of extreme climatic events on the energetics of long-lived vertebrates: the case of the greater flamingo facing cold spells in the Camargue. J. Exp. Biol. 217, 3700–3707.
- Stenhouse, I.J., and Montevecchi, W.A. (1996). Winter distribution and wrecks of little auks (dovekies) *Alle a.alle* in the Northwest Atlantic. Sula 10, 219–228.
- Lucas, Z., Horn, A., and Freedman, B. (2012). Beached bird surveys on Sable Island, Nova Scotia, 1993–2009, show a decline in the incidence of oiling. Proc. N. S. Inst. Sci. 47, 91–129.
- Meyer, X., MacIntosh, A.J.J., Chiaradia, A., Kato, A., Ramírez, F., Sueur, C., and Ropert-Coudert, Y. (2020). Oceanic thermal structure mediates dive sequences in a foraging seabird. Ecol. Evol. 10, 6610–6622.
- Hansen, K.A., Maxwell, A., Siebert, U., Larsen, O.N., and Wahlberg, M. (2017). Great cormorants (*Phalacrocorax carbo*) can detect auditory cues while diving. Naturwissenschaften 104, 45.
- Thiebault, A., Charrier, I., Aubin, T., Green, D.B., and Pistorius, P.A. (2019). First evidence of underwater vocalisations in hunting penguins. PeerJ 7, e8240.
- Udyawer, V., Chin, A., Knip, D.M., Simpfendorfer, C.A., and Heupel, M.R. (2013). Variable response of coastal sharks to severe tropical storms: environmental cues and changes in space use. Mar. Ecol. Prog. Ser. 480, 171–183.
- Patterson, W.F., Watterson, J.C., Shipp, R.L., and Cowan, J.H. (2001). Movement of tagged red snapper in the Northern Gulf of Mexico. Trans. Am. Fish. Soc. 130, 533–545.
- 30. Elliott, K.H., Chivers, L.S., Bessey, L., Gaston, A.J., Hatch, S.A., Kato, A., Osborne, O., Ropert-Coudert, Y., Speakman, J.R., and Hare, J.F. (2014). Windscapes shape seabird instantaneous energy costs but adult behavior buffers impact on offspring. Mov. Ecol. 2, 17.
- Finney, S.K., Wanless, S., and Harris, M.P. (1999). The effect of weather conditions on the feeding behaviour of a diving bird, the common guillemot *Uria aalge. J. Avian Biol. 30, 23–30.*
- Wiley, J.W., and Wunderle, J.M. (1993). The effects of hurricanes on birds, with special reference to Caribbean islands. Bird Conserv. Int. 3, 319–349.
- Wilkinson, B.P., Satgé, Y.G., Lamb, J.S., and Jodice, P.G.R. (2019). Tropical cyclones alter short-term activity patterns of a coastal seabird. Mov. Ecol. 7, 30.

 Weimerskirch, H., Bishop, C., Jeanniard-du-Dot, T., Prudor, A., and Sachs, G. (2016). Frigate birds track atmospheric conditions over months-long transoceanic flights. Science 353, 74–78.

Current Biology

Report

- 35. Barrett, R.T., Chapdelaine, G., Anker-Nilssen, T., Mosbech, A., Montevecchi, W.A., Reid, J.B., and Veit, R.R. (2006). Seabird numbers and prey consumption in the North Atlantic. ICES J. Mar. Sci. 63, 1145– 1158.
- 36. Chang, Y.-C., Chu, P.C., Centurioni, L.R., and Tseng, R.-S. (2014). Observed near-surface currents under four super typhoons. J. Mar. Syst. 139, 311–319.
- McFarlane Tranquilla, L., Hedd, A., Burke, C., Montevecchi, W.A., Regular, P.M., Robertson, G.J., Stapleton, L.A., Wilhelm, S.I., Fifield, D.A., and Buren, A.D. (2010). High Arctic sea ice conditions influence marine birds wintering in Low Arctic regions. Estuar. Coast. Shelf Sci. 89, 97–106.
- Sydeman, W.J., Poloczanska, E., Reed, T.E., and Thompson, S.A. (2015). Climate change and marine vertebrates. Science 350, 772–777.
- 39. IPCC (2014). In Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, R.K. Pachauri, and L.A. Meyer, eds..
- Tamarin-Brodsky, T., and Kaspi, Y. (2017). Enhanced poleward propagation of storms under climate change. Nat. Geosci. 10, 908–913.
- Wolf, J., Woolf, D., and Bricheno, L. (2020). Impacts of climate change on storms and waves relevant to the coastal and marine environment around the UK. MCCIP Sci. Rev. 2020, 132–157.
- 42. Akperov, M., Rinke, A., Mokhov, I.I., Semenov, V.A., Parfenova, M.R., Matthes, H., Adakudlu, M., Boberg, F., Christensen, J.H., Dembitskaya, M.A., et al. (2019). Future projections of cyclone activity in the Arctic for the 21st century from regional climate models (Arctic-CORDEX). Glob. Planet. Change 182, 103005.
- 43. Benestad, R.E., Parding, K.M., Isaksen, K., and Mezghani, A. (2016). Climate change and projections for the Barents region: what is expected to change and what will stay the same? Environ. Res. Lett. 11, 054017.
- 44. Tavares, D.C., Fulgencio de Moura, J., and Siciliano, S. (2016). Environmental predictors of seabird wrecks in a tropical coastal area. PLoS ONE *11*, e0168717.
- Tavares, D.C., Moura, J.F., Merico, A., and Siciliano, S. (2020). Mortality of seabirds migrating across the tropical Atlantic in relation to oceanographic processes. Anim. Conserv. 23, 307–319.
- 46. Beaugrand, G., Conversi, A., Atkinson, A., Cloern, J., Chiba, S., Fonda-Umani, S., Kirby, R.R., Greene, C.H., Goberville, E., Otto, S.A., et al. (2019). Prediction of unprecedented biological shifts in the global ocean. Nat. Clim. Chang. 9, 237–243.
- Serreze, M. (2009). Northern Hemisphere Cyclone Locations and Characteristics from NCEP/NCAR Reanalysis Data, Version 1. https:// nsidc.org/data/NSIDC-0423/versions/1.
- Serreze, M.C., Carse, F., Barry, R.G., and Rogers, J.C. (1997). Icelandic low cyclone activity: Climatological features, linkages with the NAO, and relationships with recent changes in the Northern hemisphere circulation. J. Clim. *10*, 453–464.
- 49. Lisovski, S., Hewson, C.M., Klaassen, R.H.G., Korner-Nievergelt, F., Kristensen, M.W., and Hahn, S. (2012). Geolocation by light: accuracy and precision affected by environmental factors. Methods Ecol. Evol. 3, 603–612.
- Phillips, R.A., Silk, J.R.D., Croxall, J.P., Afanasyev, V., and Briggs, D.R. (2004). Accuracy of geolocation estimates for flying seabirds. Mar. Ecol. Prog. Ser. 266, 265–272.
- Reynolds, R.W., Smith, T.M., Liu, C., Chelton, D.B., Casey, K.S., and Schlax, M.G. (2007). Daily high-resolution-blended analyses for sea surface temperature. J. Clim. 20, 5473–5496.
- Fort, J., Steen, H., Strøm, H., Tremblay, Y., Grønningsæter, E., Pettex, E., Porter, W.P., and Grémillet, D. (2013). Energetic consequences of contrasting winter migratory strategies in a sympatric Arctic seabird duet. J. Avian Biol. 44, 255–262.

Report



- McKnight, A., Irons, D.B., Allyn, A.J., Sullivan, K.M., and Suryan, R.M. (2011). Winter dispersal and activity patterns of post-breeding black-legged kittiwakes *Rissa tridactyla* from Prince William Sound, Alaska. Mar. Ecol. Prog. Ser. 442, 241–253.
- Shoji, A., Elliott, K., Fayet, A., Boyle, D., Perrins, C., and Guilford, T. (2015). Foraging behaviour of sympatric razorbills and puffins. Mar. Ecol. Prog. Ser. 520, 257–267.
- 55. Burke, C.M., and Montevecchi, W.A. (2018). Taking the bite out of winter: common murres (*Uria aalge*) push their dive limits to surmount energy constraints. Front. Mar. Sci. 5, 63.
- Dunn, R.E., Wanless, S., Green, J.A., Harris, M.P., and Daunt, F. (2019). Effects of body size, sex, parental care and moult strategies on auk diving behaviour outside the breeding season. J. Avian Biol. 50, https://doi.org/ 10.1111/jav.02012.
- Medina, I., Newton, E., Kearney, M.R., Mulder, R.A., Porter, W.P., and Stuart-Fox, D. (2018). Reflection of near-infrared light confers thermal protection in birds. Nat. Commun. 9, 3610.

- Amélineau, F., Grémillet, D., Bonnet, D., Le Bot, T., and Fort, J. (2016). Where to forage in the absence of sea ice? Bathymetry as a key factor for an arctic seabird. PLoS ONE *11*, e0157764.
- 59. Burger, A.E., and Simpson, M. (1986). Diving depths of Atlantic puffins and common murres. Auk *103*, 828–830.
- Chimienti, M., Cornulier, T., Owen, E., Bolton, M., Davies, I.M., Travis, J.M.J., and Scott, B.E. (2017). Taking movement data to new depths: inferring prey availability and patch profitability from seabird foraging behavior. Ecol. Evol. 7, 10252–10265.
- Belkhou, R., Cherel, Y., Heitz, A., Robin, J.-P., and Le Maho, Y. (1991). Energy contribution of proteins and lipids during prolonged fasting in the rat. Nutr. Res. *11*, 365–374.
- 62. Schmidt-Nielsen, K. (1990). Animal Physiology, Adaptation and Environment, Fourth Edition (Cambridge University).
- 63. Cherel, Y., and Groscolas, R. (1999). Relationships between nutrient storage and nutrient utilisation in long-term fasting birds and mammals. In 22 Int. Ornithol. Congr., Durban:17-34, N.J. Adams, and R.H. Slotow, eds. (BirdLife South Africa), pp. 1–14.





STAR***METHODS**

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER				
Experimental models: Organism/stains						
Little auk Alle alle	Table S1	N/A				
Black-legged kittiwake Rissa tridactyla	Table S1	N/A				
Atlantic puffin Fratercula arctica	Table S1	N/A				
Common guillemot Uria aalge	Table S1	N/A				
Brunnïch's guillemot Uria Iomvia	Table S1	N/A				
Deposited data						
Custome code to complete all statistical analyses	This study	https://doi.org/10.17632/rzszr9fpt4.1				
Software and algorithms						
R software	https://www.r-project.org/	N/A				
BRB-MKDE	Benhamou ¹⁴	N/A				
NicheMapper TM	Porter and Mitchell ⁷	N/A				

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Manon Clairbaux (clairbauxm@gmail.com).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- Environmental variables, bird and cyclone location are publicly available online. Physiological, behavioral and morphological characteristic of the species studied are available in the Table S2.
- All original code has been deposited at (https://doi.org/10.17632/rzszr9fpt4.1) and is publicly available as of the date of publication. DOIs are listed in the Key resources table.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

We focused on the winter distribution of five seabird species (little auk (*Alle alle*), Atlantic puffin (*Fratercula arctica*), common guillemot (*Uria aalge*), Brünnich's guillemot (*Uria lomvia*) and black-legged kittiwake (*Rissa tridactyla*)). These species, which account for ca. 75% of the total number of seabirds breeding along the North Atlantic Ocean³⁵ are also among the most severely hit by winter high-intensity cyclones^{2,3}. Bird locations were obtained from Global Location Sensors (GLS) deployed and retrieved during the breeding season for 1532 individuals from 39 breeding colonies across the North Atlantic Ocean (see Figure S1 and Table S1). Birds were caught at the nest according to ethical approval from each country's appropriate organization.

METHOD DETAILS

Locations and characteristics of North Atlantic cyclones during winter

According to the National Hurricane Center, cyclones are defined as large-scale air masses that rotate around a center of low atmospheric pressure. Their locations were obtained from the 'Northern Hemisphere Cyclone Locations and Characteristics from NCEP/ NCAR Reanalysis Data, Version 1' online dataset⁴⁷. Locations were calculated using the updated Serreze (1997)⁴⁸ algorithm on daily Sea Level Pressure from the National Center for Environmental Prediction (NCEP) and National Center for Atmospheric Research (NCAR) Reanalysis dataset in a 250 km grid. We focused on winter cyclones (October to February) in the North Atlantic and adjacent seas (100°W–100°E, 30°N–90°N) between 2000 and 2016. Using the Dvorak storm classification¹³ on the cyclone central pressure



we defined four classes of cyclone intensity (> 1009 hPa = Class 1; 1005-1009 hPa = Class 2; 987-1005 hPa = Class 3; < 987 hPa = Class 4) and classified each cyclone in the studied area across its duration. Finally, for each month, we calculated the average number of cyclones of each category that occurred between 2000 and 2016 for each 250 km grid cell.

Winter distribution of seabirds

Two locations per day were obtained through the recorded light levels by Global Location Sensors for each individual with an accuracy of $\pm 200 \text{ km}^{49,50}$. We focused on locations acquired during the winter period (October to February) and removed locations obtained during the two weeks on either side of the equinoxes⁴⁹ as well as spurious locations and those falling on land. For each species, we then calculated monthly utilization distributions (UD) for each individual using the BRB-MKDE software¹⁴ (March 2018 version for location in decimal degrees with hmin = 250 km) before averaging by seabird colony and species. To avoid that some individuals drive the UD of the colony, we performed a kernel stability analysis and for each month, we only considered individuals which had a sufficiently large number of locations to satisfy this stability criteria. To perform this stability analysis we calculated for each individual and for each month (n/4) UD, with n the total number of locations (after quality and equinoxes checks) for the individual and month considered: the first UD was calculated with 4 locations, the second 8, the third one 12, etc. until the total number of locations was included. Then, for each individual and for each month, we calculated the percentage of overlap between each UD and the UD obtained considering all the locations. Finally, for each species, we plot for each month the individual percentage of overlap against the number of locations included in the UD calculation and we visually determined the minimum number of locations needed to reach a high overlap percentage for the maximum of individuals. This threshold was set at 30 locations for little auks and black-legged kittiwakes and was set at 20 locations for Atlantic puffins, common and Brunnïch's guillemots. Therefore individuals having less locations than the number reeded wasn't considered when averaging UD by colony and species.

We then used 25% utilization kernels to approximate areas of core use for each colony and species during the winter months (see Data S1A). For each month, we calculated the average number of cyclones of each category that occurred between 2000 and 2016 within each colony core use areas (Figure 1; Data S1B–S1F). Finally, we interpolated each colony's UD on a 250 km grid before calculating, for each pixel, the following cyclone exposure index (CEI) for each wintering month:

$CEI(i,j) = NC(i,j) \times UDT(i,j)$

where NC(i,j) is the average number of cyclones between 2000 and 2016 within the pixel *i* for the month *j* and UDT(i,j) is the sum, all species considered, of average UD of each colony for the pixel *i* and the month *j*. The monthly cyclone exposure indexes were then averaged to obtain a unique index for the wintering period (see Figure 2).

Modeling seabird energy requirements under non-cyclonic and cyclonic conditions

We used the mechanistic model Niche Mapper^{TM 7} to calculate energy requirements for each species under cyclonic and noncyclonic conditions for each wintering month. Niche MapperTM is based on a microclimate model and an animal module: the microclimate model calculates hourly environmental conditions for the near surroundings of the bird and the animal module uses the resulting outputs together with bird morphological, behavioral and physiological properties to estimate the metabolic rate needed by the bird to maintain its body temperature at a given time with a specific behavioral time budget. We focused on a 1000 km x 1000 km area off North Newfoundland (see Figure S1) where the five seabird species aggregate during winter. This area experienced numerous cyclones during the winter months between 2000 and 2016 (see Figure S3). For each of them, we extracted environmental characteristics (sea surface temperature, air temperature and relative humidity), before averaging the values obtained by intensity class for each winter month (see Figure S4) to parameterize the microclimate module of Niche MapperTM. For each cyclone, we extracted corresponding sea surface temperatures using the NOAA High Resolution SST dataset provided by the NOAA/OAR/ESRL PSL (https://psl.noaa.gov/data/gridded/data.noaa.oisst.v2.highres.html#detail)⁵¹. The corresponding air temperature and relative humidity were extracted from the NCEP/NCAR Reanalysis dataset previously used to define cyclone locations (see above). We set the wind speed of each cyclone according to its intensity (5-13 m.s⁻¹ = Class 1; 13.1-17 m.s⁻¹ = Class 2; 17.1-32.5 m.s⁻¹ = Class 3; > 32.6 m.s⁻¹ = Class 4) using the Dvorak classification¹³. We then averaged the values obtained by intensity class to calculate mean environmental characteristic of each class for each winter month between 2000 and 2016. Finally, for each year, non-cyclonic conditions were defined by calculating the daily mean characteristics of each month using the same environmental dataset after excluding cyclone days. Wind speeds were obtained from the NCEP/NCAR Reanalysis dataset. All environmental data used were interpolated in a 250 km grid. Mean environmental conditions encountered during cyclonic and non-cyclonic events are presented in Figure S4.

Once we had parameterized the microclimate model of Niche MapperTM, we parameterized its animal module to obtain seabird daily energy requirements. As this module had already been parameterized to model energy expenditure in little auks, Brünnich's guillemots and common guillemots^{17,52}, we re-used most Niche Mapper TM input values. Missing values and values required to parameterize Niche Mapper TM for black-legged kittiwakes and Atlantic puffins, were sourced from the literature (see Table S2) and obtained through dedicated measurements. Specifically, we set the proportion of time spent flying per day during winter under non-cyclonic conditions to 9%, 11.8%, 5.7% and 4.5% for little auks, black-legged kittiwakes, Atlantic puffins and guillemots, respectively^{17,53–55}. Further, under those conditions, we considered that those species spent respectively, 24%, 18.8%, 16.9% and 16% of the day diving during winter^{17,52,53,56}. Feather reflectivity was measured with a double beam spectrophotometer (CARY 5000 UV-VIS-NIR, Agilent, USA) with a deuterium-tungsten-mercury light source. We used an integrative sphere to measure



spectral and diffuse reflectance with a 1 nm resolution across all wavelengths between 300 and 2500 nm. This range covers approximately 98% of the solar spectrum that reaches the Earth's surface. Reflectance spectra relative to a Spectralon white standard were then computed with the Cary WinUV software. For each species, measurements were made on one ventral and dorsal patch for six individuals. We then calculated the reflectivity of each sample following the methods of Medina et al.⁵⁷. For each species, the results were averaged across patches. Other morphological properties such as body dimensions were measured on adult bird carcasses of five Atlantic puffins and four black-legged kittiwakes collected on Ré Island (France) during February 2014. All Niche Mapper input data are available in Table S2.

There is limited information about the behavior of seabirds under cyclonic conditions. Yet, we benefited from personal field observations performed by a wide range of North Atlantic seabird experts (e.g., most authors of this publication, as well as Dr. Kyle Elliott and Prof Sarah Wanless, see Table S3). Those led us to assume that the seabird species considered in our study react to winter cyclones by reducing their flight activity to zero when wind speeds exceed 15 m.s⁻¹ (corresponding to cyclones of Class 2, 3 and 4). In surface-feeding black-legged kittiwakes, we also assumed that conditions experienced, when wind speeds exceed this threshold, precluded them from diving and feeding. Further, for little auks diving in the upper 30 m of the water column⁵⁸, we assumed that birds are also severely impacted by those cyclonic conditions, as the water here is too perturbed for them to dive and prey on zooplankton. In contrast, deeper diving species such as puffins and guillemots, which can dive down to 75 and 250 m, respectively^{59,60}, may still manage to feed³¹. Therefore, seabirds were modeled as resting at the surface of water when they did not fly and/or dive. Moreover, flight and diving activities were assumed to remain unchanged under cyclone of Class 1 (for which wind speeds are below 13 m.s⁻¹, see above).

Using climatic, morphological and behavioral information as stated above, we modeled species-specific, monthly energy requirements for each winter between 2000 and 2016, under four intensities of cyclones and under non-cyclonic conditions. To disentangle the effects of changes in behavior (decreased flight/diving activities) and of weather conditions on bird energy requirements during cyclones, we also calculated those requirements for each species under non-cyclonic conditions, setting behavioral parameters as those displayed during a cyclone with wind speed > 15 m.s⁻¹.

Estimation of fasting endurance

Fasting endurance was calculated for each species under Class 2, Class 3 and Class 4 cyclonic conditions in the studied area off Newfoundland for each winter months using the method described in Grémillet et al.²¹. In this particular context, we assumed that none of the species could feed under cyclonic conditions even if deep divers continued diving. The energy requirement to be supplied was calculated previously using Niche MapperTM (see above). The amount of energy available was calculated as the masses of lipids and proteins that could be metabolized during a prolonged fast (95% and 25%, respectively⁶¹), converted into energy (caloric equivalents 39.3 and $18.0 \cdot kJ \cdot g^{-1}$, respectively⁶²). The body fat and protein contents of each species are indicated in Table S2. Integrating daily body component losses through an iterative procedure, we calculated the maximal duration of fasting that lipid and protein reserves could sustain before one of them was used up. We assumed that lipids and proteins accounted for 75 and 25% of the energy expenditure until 3/4 of lipid reserves were used. They then accounted for 63 and 37%, respectively^{61,63}.

QUANTIFICATION AND STATISTICAL ANALYSIS

Monthly mean energy requirements per condition tested (Class 1 to 4 cyclones, non-cyclonic conditions with usual seabird flight/diving activities) were compared for each species using a Kruskal-Wallis test (Table 1). When this test was significant (p < 0.05), a posthoc Dunn's test was conducted (p values adjusted with the Benjamini-Hochberg method) to conclude on the difference between energy requirements experienced under cyclonic and non-cyclonic conditions (Data S1G–S1K).