

Application of satellite altimetry to ocean circulation studies: 1987-1994

Lee-Lueng Fu

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California

Robert E. Cheney

National Ocean Service, National Oceanic and Atmospheric Administration, Silver Spring, Maryland

1. Introduction

The ocean plays a key role in determining the global climate and its time evolution. To understand this role and subsequently develop techniques for predicting future climate, one must understand the dynamics of the global ocean circulation—the movement of water that transports mass, heat, salt, and other biogeochemical properties of the ocean that are closely linked to the processes of climate change. The only viable approach to observing the global ocean circulation with sufficient resolution and consistent sampling is the use of a satellite radar altimeter to measure the height of the sea surface—the sea level (Wunsch and Gaposchkin, 1980; Stewart, 1983; Wunsch, 1992). After removing the effects of the tides and atmospheric pressure from the observation, the deviation of the sea surface from the geoid, called the ocean dynamic topography, is readily related to the velocity of the surface geostrophic flow—a component of the surface flow on which the surface pressure force is balanced by the Coriolis force due to the Earth's rotation. Moreover, the ocean dynamic topography provides a strong constraint for determining the ocean circulation through the entire water column via the dynamic equations governing the fluid motion. Precise measurement of the shape of the global sea surface thus provides a powerful tool for studying the dynamics of the ocean circulation.

Progress in the application of satellite altimetry for ocean circulation studies from 1987-1994 is reviewed in this paper. The reader is referred to Douglas et al. (1987), Brown and Cheney (1983), and Fu (1983) for early development in the subject. The last review of the subject was conducted eight years ago (Douglas et al., 1987). The past eight years have seen the greatest strides in the development of satellite altimetry as an observational tool in oceanography. The U.S. Navy launched Geosat in 1985 with a primary objective of mapping the marine gravity field for military applications. Most of the data collected during Geosat's first 18 month primary mission (the Geodetic Mission) are still classified. However, the declassified portion of the data has led to significant advancement in marine geophysics (e.g., McAdoo and Marks, 1992). When its primary mission was achieved, Geosat was maneuvered into a 17-day repeat orbit, and an extended mission (the Exact Repeat Mission) for oceanographic applications started in October 1986 (Born et al., 1987; Douglas and Cheney, 1990). Although the mission was not designed specifically for oceanographic applications, the 2.5-year data set created the first opportunity for oceanographers to experiment with a multi-year global

data set (see the March and October 1990 issues of the *Journal of Geophysical Research* devoted to Geosat.)

The most serious Geosat error source has been the uncertainty in the radial orbit height, but significant progress continues to be made in this area. The Geosat orbit error was on the order of 2 m in the initial data release, but was later reduced to about 30-50 cm in a later release largely through gravity model improvements (Cheney et al., 1991; Haines et al., 1994). However, the orbit accuracy rapidly degrades to the 1-m level after mid 1988 through the end of the mission due to inadequate modeling of the drag force caused by the increased solar activities. By using the altimeter data as a constraint, Shum et al. (1990 a) achieved an orbit precision of about 20 cm. Finally, in 1993, the U.S. Navy released additional Geosat Doppler tracking data which have led to a further reduction in the orbit error to about 10 cm (S. Nerem and R. Williamson, private communication). Other important error sources include the altimeter signal delay caused by tropospheric water vapor and ionospheric free electrons (Emery et al., 1990; Musman et al., 1990). Based on the best orbit available, the present Geosat error budget is at the level of about 15 cm, an impressive accomplishment in view of the initial low expectations for the mission.

In 1991, two years after the end of the Geosat mission, the European Space Agency launched the ERS-1 (European Remote Sensing Satellite-1) altimeter. An identical satellite, ERS-2, will take its place in 1995. These two missions provide the type of continuous, long-term coverage needed for studies of interannual ocean variability and also provide altimeter data to operational users in near-real time (Cheney and Lillibridge, 1992). Because the ERS series of satellites carry a suite of sensors, a number of different ground track patterns have been followed, including 3-day and 35-day repeats. During most of its final year of operation, the ERS-1 satellite was flown in an orbit that yielded a dense global network of altimeter profiles spaced 15-km apart at the equator for geodetic studies. As was the case for Geosat, orbit error is the primary obstacle to be overcome for ERS-1 altimeter applications. The problem was compounded by early failure of one of the two onboard tracking systems. However, laser tracking together with advanced gravity models have now resulted in orbits with an accuracy of about 15 cm (Lillibridge et al., 1993).

To be useful for studying the ocean circulation, especially at the basin scales, the accuracy of altimetry measurement must be less than 15 cm (TOPEX Science Working Group, 1981). To achieve this goal, a satellite mission with specially designed instrumentation and orbit configuration, and much improved knowledge in the Earth's gravity field for orbit determination was required. The TOPEX/POSEIDON Mission was the result of this conviction by scientists and engineers in both the U.S. and France (Fu et al., 1991 a; The TOPEX/POSEIDON Science Working

Copyright 1995 by the American Geophysical Union.

Paper number 95RG00187.
8755-1209/95/95RG-00187\$15.00

Team, 1991; Fu et al., 1994). TOPEX is an acronym standing for Ocean Topography Experiment, the name originally used by the U.S.; POSEIDON is the original French name for the mission. This joint U.S./France mission was launched in August, 1992 with an expected lifetime of three to five years. The performance of the mission has exceeded its specification. Orbit and altimeter accuracies are estimated to be 3.5 cm and 3.2 cm, respectively, resulting in an absolute accuracy of 4.7 cm for the determination of the geocentric sea level (Fu et al., 1994). For the first time, oceanographers have a global observing system that is providing data of sufficient accuracy and sampling for the study of the global ocean circulation. Preliminary results from TOPEX/POSEIDON can be found in the December 1994 issue of the *Journal of Geophysical Research*. Significant advancement in the knowledge of the large-scale ocean circulation is expected from the mission in the near future.

A brief summary of the accomplishments made in the past eight years is given below in categories based on oceanographic phenomenology. Only those papers addressing direct applications to the ocean circulation are reviewed. A large body of literature on techniques for altimetry correction and data analysis has thus been left out (see Chelton, 1988 for an overview). Of great importance to the utility of altimeter data is the correction for the tidal effects on the sea level observation. The reader is referred to Cartwright and Ray (1990) for estimating tides from altimeter data.

2. The Western Boundary Currents

The western boundary currents can be easily resolved even by the relatively crude Geos-3 radar altimeter because of the large sea level variation across the currents (e.g., Fu et al., 1987). A host of studies of the various western boundary currents of the world's oceans have been conducted using the much improved Geosat data. Among the numerous accomplishments is a powerful technique developed by Kelly and Gille (1990) for estimating the absolute dynamic ocean topography across strong currents from their temporal variabilities. An analytical model is used to describe the dynamic topography of an isolated jet with its kinematic parameters (i.e., location, width, and amplitude) determined from the observed temporal sea level changes as an inverse problem (also see Tai, 1990). Comparison of the altimetry-derived velocities with simultaneous in-situ velocity measurements demonstrated the validity of the technique (Joyce et al., 1990). The technique was applied to the Gulf Stream and the Kuroshio for studying their spatial and temporal characteristics with results consistent with in-situ data (Kelly, 1991; Qiu et al., 1991). Tai and White (1990) extended this approach to addressing the eddy-mean flow interaction of the Kuroshio Extension. Other studies based on the technique have been conducted on the recirculation flanking the main current of the Kuroshio (Qiu, 1992) and the Gulf Stream (Qiu, 1994). By combining the altimetric estimate of the surface dynamic topography with historical hydrographic data, Qiu (1994) estimated the deep circulation of the Gulf Stream region with results comparing well to in-situ data. Kelly and Watts (1994) applied the technique to both altimetry data and inverted echo sounder data and obtained excellent agreement. Qiu and Kelly (1993) used altimetrically derived surface current velocities in conjunction with atmospheric wind and thermal forcing to study the heat budget of the upper ocean in the Gulf Stream and the Kuroshio regions.

Using a straightforward technique to examine the cross stream sea level differences, Zlotnicki (1991) reported that both the Gulf Stream and the Kuroshio, after entering the open ocean, had similar seasonal cycles with peak surface currents occurring in the fall, consistent with the findings of the studies cited in the preceding paragraph. The discrepancy from the finding of Fu et al. (1987) that reported a peak Gulf Stream surface velocity in the spring during the Geos-3 mission could be due to interannual variability or large systematic errors in the Geos-3 data. A downstream decrease in the time scales of the Gulf Stream sea level variability was found by Vazquez et al. (1990). The spatial and temporal characteristics of the meandering of the Gulf Stream was investigated further by Vazquez (1993).

There have been numerous intercomparisons made between Geosat altimetry and in-situ data in the regions of the Gulf Stream and the Kuroshio (Hallock et al., 1989; Hallock and Teague, 1993; Willebrand et al. 1990; Horton et al., 1992; Blaha and Lunde, 1992; Joyce et al., 1990), demonstrating the quality of the Geosat data for the study of the mesoscale structure of the western boundary currents. Carnes et al. (1990) investigated the feasibility of estimating subsurface temperature fields from altimeter data in the Gulf Stream region with mixed results. Using a combination of the Geosat altimeter data with in-situ hydrographic data, a number of investigators constructed for the Gulf Stream region the so-called synthetic geoid (Mitchell et al., 1990; Porter et al., 1989, 1992; Glenn et al., 1991), which was useful for deriving approximate absolute Gulf Stream topography for research as well as operational applications. High-resolution geoids in the Gulf Stream region were also obtained by using geodetic techniques (Rapp and Wang, 1994) and by combining altimetry, hydrography, and direct current velocities (Kelly et al., 1991). Rapp and Smith (1994) used the TOPEX/POSEIDON data and the geoid model of Rapp and Wang (1994) to study the characteristics of the Gulf Stream.

The Geosat data have also been used to study the Loop Current and its eddies in the Gulf of Mexico. D. Johnson et al. (1992) tracked two Loop Current eddies which drifted southwestward across the Gulf. Jacobs and Leben (1990) reported a 10.5-month period for the shedding of eddies by the Loop Current. Leben et al. (1990) constructed maps of the mean sea surface and eddy variability of the Gulf.

Using the Geosat altimeter data with a numerical model, Matano et al. (1993) studied the seasonal variability of the Brazil and the Malvinas Currents and found that the seasonal cycles of the two currents were opposite in phase (with the maximum of the Brazil Current during the austral summer). They interpreted the result in terms of the seasonal migration of the confluence zone of the two currents. Spatial and temporal characteristics of the variability of the Brazil-Malvinas Confluence region were analyzed by Provost and Le Traon (1993). They reported the dominance of semiannual fluctuations characterized by alternating positive and negative sea level anomalies with length scales of 400–500 km. The mesoscale variability was highly inhomogeneous and anisotropic. They also computed the Reynolds stress and discussed its relation to the dynamics of the mean flow.

The variability of the Agulhas Current was investigated using the Geosat data in a number of studies. Fu and Zlotnicki (1989) calculated the wavenumber-frequency spectrum which revealed the dominant wavelengths and periods as 300–800 km and 50–200 days. Quartly and Srokosz (1993) reported that the seasonal variation evidenced in the Geosat data was not reproduced by the Fine Resolution Antarctic Model (FRAM), suggesting possible

directions for the improvement of the model. Wakker et al. (1990) calculated the dynamic topography and the eddy variability statistics in the region.

3. The Tropical Oceans

Altimetric data are particularly useful for observing oceanographic phenomena in the tropical oceans where the zonal scales are large and the meridional scales are relatively small, making the orbit errors less important because their effects are primarily on large meridional scales. However, careful treatment of the orbit errors still makes a difference when the errors are as large as in the early versions of the Geosat data (Cheney and Miller, 1990). There has been significant progress in all three tropical oceans.

The Tropical Pacific Ocean

The 1986–87 El Niño was a major focus of the studies of the tropical Pacific using the Geosat data. Cheney and Miller (1988) and Miller et al. (1988) detected the Kelvin waves that initiated the El Niño as the ocean's response to a burst of anomalous westerly wind events (Kelvin wave is an edge wave involving the Earth's gravity and rotation). Using 4-year long sea level time series constructed from both the Geosat Geodetic Mission and the Exact Repeat Mission, Miller and Cheney (1990) studied the budget of the upper ocean water mass. They identified a seasonal exchange of upper ocean water between the equatorial region (7° S– 7° N) and the region just to the north (8° N– 20° N). This exchange process was enhanced during the El Niño. The generation, propagation, and reflection of Kelvin waves and Rossby waves (a low-frequency wave caused by the vorticity of a rotating fluid) in the equatorial wave guide were studied in detail (White et al., 1990 a; Delcroix et al., 1991; du Penhoat et al., 1992). Extensive comparisons were made between the Geosat observations and in-situ observations from tide gauges and expendable bathythermographs (XBTs) (Tai et al., 1989; Cheney et al., 1989). The Geosat data were also compared to simulations of ocean general circulation models with some success (Cheney et al., 1989; Chao et al., 1993).

Evidence for equatorial waves in the strong shear region of the equatorial current system was reported in the Seasat data (Malarde et al., 1987; Musman, 1989, 1992). Characteristics of these waves were studied in detail based on the Geosat data by Perigaud (1990), who identified wave activities in two shear zones centered at 5° N (between the South Equatorial Current and the North Equatorial Counter Current) and 12° N (between the North Equatorial Counter Current and the North Equatorial Current) with distinctively different periods and wavelengths. She also found correlations of the intensity of these waves to the seasonally varying strength of the shears. Using the Geosat data with various in-situ data, Hansen and Maul (1991) and Maul et al. (1992) studied the formation and evolution of eddies in the eastern end of the North Equatorial Counter Current. These eddies, having strong nonlinear character, were formed during late fall when the current was strongest. Similar results were obtained from the TOPEX/POSEIDON data (Giese et al., 1994). Busalacchi et al. (1994) demonstrated the unique aspect of TOPEX/POSEIDON in observing the 20–30 day instability waves.

The Tropical Indian Ocean

Using the Seasat altimeter crossover differences, Perigaud and Minster (1988) investigated the movement of the "Great Whirl"

formed in the Somali Current and found fair agreement between altimetry data and in-situ data. Variability of the Somali Current was studied by Perigaud and Delecluse (1989) using the Seasat altimeter data and scatterometer data and numerical models. They found that nonlinearity is crucial in explaining the observations. A large anticyclonic eddy formed in the eastern Arabian Sea during the northeast monsoon was studied by Bruce et al. (1994) using the Geosat data with a numerical model. Perigaud and Delecluse (1992) investigated the annual sea level variations using the Geosat altimeter data and a shallow-water model. Significant correlation was found between the first complex empirical orthogonal functions of both the observation and the model simulation. This annual signal was interpreted in terms of Rossby waves with the maximum amplitude located near 12° S and 90° E. Using four years' worth of the Geosat data, Perigaud and Delecluse (1993) investigated the interannual sea level variations and compared the observation to the simulation by a shallow-water model. Averaged over the basin studied (north of 20° S), the observation and the simulation were highly correlated (0.92). The signal was characterized by a sea level rise during the 1986–87 El Niño with the maximum amplitude (5 cm) occurring in the southern domain (10° S– 20° S). They also used the Geosat data to test the quality of the wind driving the model.

The Tropical Atlantic Ocean

Combining the Seasat data with the Geos-3 data into a monthly data base, Menard (1988) studied the seasonal cycle of the North Equatorial Counter Current and obtained encouraging comparisons of the altimetry analysis with in-situ data and model simulations. Carton (1989) studied the variation of the equatorial current systems using the first year of the Geosat data. He also estimated the zonal sea surface slopes associated with the current variabilities. A follow-up study was conducted to examine the seasonal variations of the North Equatorial Counter Current using two years' worth of the Geosat data (Carton and Katz, 1990). The current maximum was found at 38° W with peak transport of 40 Sv (1 Sv = 10^9 kg/sec). Extensive intercomparisons were made between the Geosat data and various in-situ data (Carton and Katz, 1990, Arnault et al., 1990, 1992 a). Arnault et al. (1992 b) made extensive comparisons of the low-frequency variabilities between altimetry analysis and model simulations. Interannual variabilities in the Gulf of Guinea related to the 1986–87 El Niño were discussed. Arnault and Cheney (1994) conducted further investigation using 4.5 years' worth of improved Geosat data. They described 80% of the sea level variance in terms of 3 empirical orthogonal functions and reported an increase of the volume of the equatorial upper ocean during 1987–89, indicative of the effect of the 1986–87 El Niño. Didden and Schott (1992) examined the variabilities of the North Equatorial Counter Current and the North Brazil Current using the Geosat data and the World Ocean Circulation Experiment (WOCE) Community Model and obtained good agreement between the two. Musman (1992) reported evidence for the equatorial instability waves at 4° N and 4° S with a period of 25 days and a wavelength of 970 km.

4. The Mesoscale Eddies

The spatial scales of most of the significant errors in altimeter data (the orbit, the tides, the signal delay in the transmission media, the sea-state bias) are generally longer than 1000 km, making the detection of mesoscale features, whose wavelengths are less than 1000 km, relatively less prone to the measurement errors.

However, certain along-track smoothing is required to reduce the random measurement noise, limiting the wavelengths of the variability examined to greater than about 60 km (e.g., Le Traon et al., 1990). Moreover, a single altimeter mission has only limited capability for synoptic mapping of the details of mesoscale eddies in both space and time (Chelton and Schlax, 1994).

Maps of the statistics of the global mesoscale variability with much higher accuracy and resolution than the Seasat results have been produced from the Geosat data due to its large data volume and dense ground tracks (Zlotnicki et al., 1989; Sandwell and Zhang, 1989; Shum et al., 1990b). Sandwell and Zhang (1989) computed the sea surface slope, which filtered out most of the large-scale errors, and converted it directly to the kinetic energy of ocean currents. Seasonally-varying mesoscale energy was reported in the northeast Atlantic and the northeast Pacific by Zlotnicki et al. (1989). Energy level was found to be correlated with the strength of the wind stress curl. Shum et al. (1990b) obtained similar results. Wavenumber spectra were estimated in a wide range of locations (Fu and Zlotnicki, 1989; Le Traon et al., 1990; Stammer and Boning, 1992). Different spectral slopes were found between the high-energy areas and the low-energy areas, but the interpretations were controversial (Le Traon, 1993; Stammer and Boning, 1993).

De Mey and Menard (1989) combined the Geos-3 data with the Seasat data in an objective analysis of the eddy field in the Polymode region (a 500-km square centered at 29° N and 70° W). The analyzed field was assimilated into an ocean model that yielded realistic subsurface fields. Bisagni (1991) created time series of sea level maps using the Geosat crossover data in a similar region, delineating westward propagating eddy fields that were consistent with previous in-situ data in the region. Tokmakian and Challenor (1993) applied the Geosat data to studying the eddy field in the Azores frontal region and the Canary basin of the North Atlantic. They reported slightly higher eddy energy in winter and discussed evidence for Rossby waves, although the effects of tidal aliasing was a concern for the detection of Rossby waves of the annual period (Schlax and Chelton, 1994). The Geosat data were also applied to the North Atlantic Current (De Mey, 1992) and the frontal zones east of Iceland (Robinson et al., 1989). Significant correlations between Geosat and in-situ observations were reported even in areas of extremely low eddy energy such as the Iberian Basin (Stammer et al., 1991) and the Cape Verde Frontal Zone (Zlotnicki et al., 1993). A systematic study of the relationship between the time scale and the space scale of the eddy field in the North Atlantic was conducted by Le Traon (1991), who identified two regimes: higher energy areas where the space and time scales are proportional to each other, consistent with the quasigeostrophic turbulence theory; low energy areas where the space and time scales are inversely related, consistent with the linear Rossby wave dynamics.

The spatial and temporal characteristics of the eddy field in the South Atlantic were investigated by Le Traon and Minster (1993) and Forbes et al. (1993). Both reported evidence for westward propagating Rossby waves: the former emphasized the waves of the semiannual period at the subtropical latitudes west of the Walvis ridge; the latter identified the Agulhas Retroflexion as a source for the waves with periods of 400–500 days. The variation of the characteristics of the eddy wavenumber spectrum in the region was extensively examined with results consistent with the findings of Fu and Zlotnicki (1989) and Le Traon et al. (1990).

The mesoscale variability of the Antarctic Circumpolar Current was investigated by Chelton et al. (1990), who reported that the

eddy field was highly correlated to the mean flow and both were apparently steered by the bathymetry (also see Sandwell and Zhang, 1989). Eddy Reynolds stress was estimated at the crossover locations of Geosat (Morrow et al., 1992, 1994; T. Johnson et al., 1992). The resulting horizontal eddy momentum flux tends to generally accelerate the mean flow, but the zonally averaged eddy flux is an order of magnitude too small, and in the wrong direction to balance the eastward momentum input from the wind.

Jacobs et al. (1993) fit a Rossby wave model to the Geosat data in the entire Pacific Ocean. They found that less than 5% of the variance could be accounted for by the model. However, most of the estimated wave amplitude was above the estimation error. Van Woert and Price (1993) reported evidence for Rossby waves north of the Hawaii Islands and found close agreement of their propagation characteristics and latitudinal dependence with the theoretical dispersion relation. White et al. (1990b) demonstrated that the annual fluctuations in the California Current region were consistent with Rossby waves propagating into the open ocean with wave fronts parallel to the bathymetry. Kelly et al. (1993) discussed the northern limit (37° N) for Rossby wave propagation in the region. The eddy activities in the California Current were found to be consistent with the simulations of a primitive-equation model (Pares-Sierra et al., 1993), indicating that the eddies were forced by the wind adjacent to the coast. The annual-period Rossby waves were also detected in the Alaska Gyre (Matthews et al., 1992). Again the finding must be viewed with caution due to the aliasing effects of the tides (Schlax and Chelton, 1994). Evidence for topographic planetary waves in the Bering Slope Current was reported by Okkonen (1993).

The formation, propagation, and mass transport of eddies were studied in detail by a number of investigators. Didden and Schott (1993) examined the eddies in the retroflexion region of the North Brazil Current. Individual eddies were traced to the entrance of the Caribbean. Seasonal variation of the eddy energy was correlated to the seasonal cycle of the retroflexion of the North Brazil Current into the North Equatorial Counter Current. The interhemispheric exchange of water mass was estimated to be 3 Sv. Gordon and Haxby (1990) were able to trace the eddies spun off from the retroflexion of the Agulhas Current all the way to the western South Atlantic. They estimated as much as 15 Sv of the Indian Ocean water being transported to the South Atlantic by this process. Okkonen (1992) employed a novel eddy-tracking technique to follow and diagnose the shedding of an eddy from the Alaskan Stream. Ichikawa and Imawaki (1994) documented for the first time the complete life history of a cold-core eddy shed from the Kuroshio.

5. The Large-Scale Variability

The "large-scale" is used here to refer to scales larger than 1000 km, a loosely defined upper limit for the mesoscale discussed in Section 4. In this range of scales, the signal-to-noise ratio is generally small for all the altimeters except for TOPEX/POSEIDON. As noted above, a variety of error sources such as the orbit, the tides, the signal transmission media, and the instrument systematic errors have dominant scales larger than 1000 km. Despite these obstacles, numerous attempts have been made to examine the large-scale ocean variabilities using the pre-TOPEX/POSEIDON altimeter data. The dominant orbit errors were reduced by a variety of methods, ranging from a dynamic approach utilizing gravity model adjustment (Koblinsky et al., 1992) to a purely geometric approach utilizing the spectral

characteristics of the orbit error (Wunsch, 1991 a). The reader is referred to Wagner and Tai (1994) for a discussion of the effects of the orbit error removal procedures on the detection of ocean signals.

Comparison of the low degree and order spherical harmonics of the Geosat sea level variations to a set of tide gauge data yielded a rms difference of 5–10 cm (Wunsch, 1991a; Koblinsky et al., 1992). Similar results were obtained by Harangozo et al. (1993). Wunsch (1991a) demonstrated a method of combining altimeter data with tide gauge data in enhancing the accuracy of the estimate of the global sea level variability. The results were used to investigate the large-scale response of the ocean to atmospheric forcing (Wunsch, 1991b). Up to 50% of the variance was accounted for by the forcing of wind and atmospheric pressure. Spatial pattern of the global annual and semiannual variabilities were examined in detail by Koblinsky et al. (1992) and Jacobs et al. (1992). Contamination of the annual cycle by ocean tides was removed by an empirical method in the latter study. The interannual change due to the 1986–87 El Niño were demonstrated (Wunsch, 1991a; Koblinsky et al., 1992; Koblinsky, 1993). Preliminary results from TOPEX/POSEIDON have demonstrated its capability of more accurate detection of the large-scale variabilities such as a hemispheric asymmetry in the global annual cycle, with the amplitude in the Northern Hemisphere twice as large as in the Southern Hemisphere (Stammer and Wunsch, 1994; Cheney et al., 1994; Knudsen, 1994). Behringer (1994) reported close agreement of the large-scale variabilities of the North Atlantic from the TOPEX/POSEIDON observations with an ocean data assimilation system within 2 cm.

There were also regional studies of the gyre- and basin-scale variabilities. The annual and interannual variabilities of the Alaska Gyre was investigated by Bhaskaran et al. (1993) based on a four-year Geosat database. He found significant correlations of the sea level with atmospheric pressure and the Southern Oscillation Index (an indicator for the climatic condition of the tropical Pacific based on the sea level pressure difference between Darwin and Tahiti). Kelly et al. (1993) examined the interannual variations of the Alaska and the California currents and found that the two currents were out of phase with each other. Chelton et al. (1990) examined the large-scale structure of the temporal variability of the Antarctic Circumpolar Current (ACC) and reported that only a minor portion (33%) of the variance was zonally coherent, corresponding to the annual, semiannual, and interannual variabilities. The current has strong regional characteristics in each basin. Using the Seasat altimeter and scatterometer data, Mestas-Nunez et al. (1992) analyzed the relation between the variability of the ACC and the wind forcing and reported that a quarter of the variance was accounted for by the linear response of the ocean to the forcing of the curl of the wind stress (the time-dependent Sverdrup relation). Gille (1994) used the technique of Kelly and Gille (1990) for estimating the mean dynamic topography of the entire ACC. Her results were also able to account for 40–70% of the sea level variance along the axis of the ACC in terms of the meandering of the current.

By comparing the global sea level derived from Seasat to that from Geosat, Haines et al. (1992) examined the difference between the summer of 1978 to that of 1987 and found good agreement with in-situ observations. This result demonstrated the potential of combining observations from different altimetry missions to monitor long-term changes in the ocean. The success of Haines et al. (1992) was largely due to their recomputing the Seasat orbit using the same gravity model as the one used for

Geosat, thus eliminating the large difference in the geographically-correlated orbit errors that could be aliased to temporal changes (Cheney and Douglas, 1988). Based on the difference between the Geosat and the ERS-1 observations, Jacobs et al. (1994) reported a large-scale sea level change from 1987 to 1992 in the Kuroshio Extension region. Using a model of the Pacific Ocean, they demonstrated that the observed sea level change was caused by a Rossby wave excited by the 1982-83 El Niño. It took ten years for the Rossby wave to travel across the Pacific at about 40° N. This study provides the first evidence for a decadal effect of El Niño.

Change in the global mean sea level has both scientific and practical implications. The dense, global sampling of a satellite altimeter makes it one of the most viable approaches to monitoring the change in the global mean sea level. Born et al. (1986) first demonstrated the idea using the Seasat data. Due to the large orbit error, the result showed more than 10 cm fluctuations in 24 days. Tapley et al. (1992) showed that the rms variability of the global mean sea level calculated from two years' worth of the Geosat data was about 2 cm. The rate of change was estimated ± 0.5 cm/year. Similar results were obtained by Visser et al. (1993). The various errors affecting the global mean sea level estimate were discussed in detail by Wagner and Cheney (1992). Based on the first year of the TOPEX/POSEIDON data, Nerem et al. (1994) reported that the rms variability of the global mean sea level was reduced to a level of 0.5 cm. Although significant progress has been made, there is still a long way before altimeter can detect the 1 mm per year global mean sea level change.

6. The General Circulation

Determining the absolute surface geostrophic circulation of the world's oceans has been a tantalizing prospect of satellite altimetry since the advent of the technique. The difficulties arise from the errors in both the orbit and the geoid. The orbit errors of concern are the systematic (or geographically correlated) component that cannot be reduced by averaging in time. This error must be reduced by improved gravity models or by continuous precision satellite-tracking using the Global Positioning System (Bertiger et al., 1994). As part of the TOPEX/POSEIDON Mission, gravity model improvement efforts were initiated in both the U.S. and France in the 1980s. These efforts have progressively produced improved gravity models (e.g., Marsh et al., 1988, 1990 a; Rapp, 1993), which in turn have also led to improved long-wavelength components of the geoid. For instance, Tai (1988) reported improved estimate of the ocean dynamic topography using the gravity model of Marsh et al. (1988)

The ocean dynamic topography is usually solved as an inverse problem involving simultaneous adjustment of the orbit, the gravity model, and the dynamic topography. Using this approach, a number of groups have estimated the global ocean dynamic topography up to spherical harmonics degree and order 6–10 (Tapley et al., 1988; Marsh et al., 1990 b; Denker and Rapp, 1990; Nerem et al., 1990; Visser et al., 1993). For shorter scales, the geoid errors overwhelm the oceanographic signals. Using an inverse model of the North Atlantic circulation involving a large amount of oceanographic data, Martel and Wunsch (1993) evaluated the Geosat altimetric dynamic topography of Nerem et al. (1990) and concluded that the Geosat result was inconsistent with the conventional oceanographic data and that the error in the Geosat result was at least a factor of two too large to improve the existing knowledge of the circulation. Preliminary examination

of the data from the TOPEX/POSEIDON Mission have shown promise for using the data to improve estimate of the ocean general circulation (Nerem et al., 1994; Tapley et al., 1994; Stammer and Wunsch, 1994).

7. Data Assimilation

It has long been recognized that one needs to combine observations with dynamic models to achieve an optimal description of a turbulent global fluid system such as the ocean. This model/data combination is referred to as "data assimilation" following the terminology in meteorology. The development of data assimilation into numerical ocean models has been hampered by the lack of well-sampled data. For a review of the status of the subject the reader is referred to Ghil and Malanotte-Rizzoli (1991) and Bennett (1992). The availability of satellite altimetry data has spurred significant progress in ocean data assimilation over the past several years.

One approach to data assimilation is direct data insertion, or reinitialization. Berry and Marshall (1989) inserted data only to the surface layer and let the information be dynamically transferred to the lower layers. De Mey and Robinson (1987), De Mey and Menard (1989), and Hurlburt et al. (1990) projected the surface information to the deep layers using statistical methods such as empirical orthogonal functions. Altimetry data along with the subsurface projection were then directly inserted into a model, resulting in a more efficient vertical information transfer. Robinson and Walstad (1987) used a feature model for the Gulf Stream and its eddies to initialize the 3-dimensional field for data insertion. Haines (1991) demonstrated a novel approach in which the data insertion was applied to the surface stream function while the potential vorticity of the lower layers were kept unchanged so as to create an effective adjustment of the lower layers to the data insertion at the top layer.

A relatively simple and effective scheme for assimilating data continuously into a model is the so-called nudging technique (Holland and Malanotte-Rizzoli, 1989; Verron, 1990, 1992). The information of the altimetry data is ingested into the surface layer by placing in the model equations a term that is proportional to the difference between the model sea surface height and the observation. The information of the data is transferred to the deep layers by the model dynamics. This technique works quite well for quasi-geostrophic models and has been applied to the Geosat data in the Agulhas Current (Holland et al., 1991) and the western North Atlantic (Verron et al., 1992). Blayo et al. (1994) applied the same approach to the TOPEX/POSEIDON data in the North Atlantic. Smedstad and Fox (1994) demonstrated the use of the technique with a primitive equation model. However, the effects of observation error and model error are generally not taken into account in the nudging approach.

White et al. (1990 c) employed an optimal interpolation (OI) technique to assimilate the Geosat data into a quasi-geostrophic model of the California Current, specifically taking into account the model and observation errors in the assimilation step. However, the information of the data was assimilated only to the surface layer, while the vertical transfer of information was via the model dynamics. Dombrowski and De Mey (1992) used the empirical orthogonal function technique to project the altimetry information to the lower layers of a quasi-geostrophic model and applied an OI scheme to assimilating the Geosat data into all the vertical layers simultaneously. Using statistical correlation in the transfer of information from the sea surface height to subsurface

fields, Mellor and Ezer (1991) and Ezer and Mellor (1994) demonstrated the use of an OI scheme in assimilating altimetry data into a primitive equation model of the Gulf Stream region. The statistical correlation between the sea surface height and subsurface temperature was demonstrated by Ezer et al. (1993). The OI approach is particularly effective in mapping the horizontal fields, but the vertical transfer of information to subsurface fields is somewhat problematic.

Variational methods have been applied to seeking model solutions that minimize a "cost function" related to the model-data misfit (Moore, 1991; Bourles et al., 1992; Schroter et al., 1993; Greiner and Perigaud, 1994). The solutions obtained are exact solutions to the model equations in the so-called strong constraint approach. Using a "weak-constraint" approach, Egbert et al. (1994) demonstrated the use of the variational method to obtain an optimal estimation of the global ocean tides by combining the TOPEX/POSEIDON data with the Laplace tidal equation. In addition to the optimal model solution, other parameters such as friction coefficient, wind forcing amplitude, and model initial conditions can also be estimated as part of the solutions. Although the approach is quite effective in testing the consistency between models and data, error estimates for the solutions were seldom discussed because of the required large computational burden.

The popular method of the "Kalman filter/smoothing" in the engineering community is a sequential approach to optimal estimation by combining model prediction with observations in a way to minimize the estimation error. Miller (1989) demonstrated the use of the Kalman filter to assimilate the altimeter differences into linear ocean models. Gaspar and Wunsch (1989) applied the technique to extracting Rossby wave signals from the Geosat data in the western North Atlantic. They found that only about 6% of the data variance was consistent with the Rossby wave dynamics. Fu et al. (1991 b, 1993) showed that a significant portion (68%) of the Geosat data in the tropical Pacific was consistent with the dynamics of linear Kelvin waves and Rossby waves. Gourdeau et al. (1992) obtained similar results in the Tropical Atlantic. The computational burden of the Kalman filter/smoothing is notoriously heavy, because the error covariance also needs to be carried forward in time by the model dynamics. Noting the rapid approach to a steady state of the error covariance in many Kalman filter applications, Fukumori et al. (1993) developed a steady-state Kalman filter/smoothing and demonstrated its use in assimilating the Geosat data into a simple, coarse-resolution general circulation model of the North Atlantic. The technique was also evaluated by Fu et al. (1993) by comparing it to an exact, time-varying Kalman filter calculation. To make the technique feasible to a full-scale ocean general circulation model, Fukumori and Malanotte-Rizzoli (1995) demonstrated a further approximation of the Kalman filter by assimilating only the large-scale content of the data. Such approach holds promise for assimilating data into truly global models.

8. Conclusions and Outlook

The last eight years have seen enormous progress in the application of the Geosat altimetry data to the study of oceanographic problems, albeit with a great deal of effort in treating the various errors in the data. Significant advancement has been made in the knowledge of the temporal variabilities of the western boundary currents, the mesoscale eddies, the tropical circulation, and the Antarctic Circumpolar Current. The multiyear, global data set has

generated a great deal of progress in developing methods of data assimilation, setting the stage for using altimetry data as a fundamental constraint for global ocean modeling and description.

The launch of TOPEX/POSEIDON has provided scientists with the first truly global ocean observing system, producing a "snapshot" of the global ocean circulation with an unprecedented accuracy every 10 days. Coupled with two other concurrent developments—the World Ocean Circulation Experiment (WOCE) and the maturing technology of global ocean modeling—TOPEX/POSEIDON is creating enormous opportunities for making significant advancement in the understanding of global ocean dynamics. The high accuracy of TOPEX/POSEIDON would help calibrate and improve other less accurate missions such as ERS-1 and 2 and enhance their value for large-scale circulation studies. To achieve the ultimate goal of understanding the ocean's role in climate, the observation provided by TOPEX/POSEIDON must be sustained with consistent quality. Plans are in place to extend the mission from three to at least six years as long as the flight hardware is functioning. A series of follow-on missions to extend the TOPEX/POSEIDON data stream to the indefinite future is being studied jointly by NASA, NOAA, and the French CNES (Centre National d'Etudes Spatiales). The prospect of having a combination of various altimeters with at least one of them having the TOPEX/POSEIDON accuracy holds great promise for ocean circulation research in the next decade. This series of altimetry missions will become a key element of a global ocean observing system for unravelling the ocean's role in climate change.

Acknowledgments. The bibliography in altimetry research compiled by Dr. C. R. Rapley was a valuable source of information for the review. The authors are indebted to Dudley Chelton for his thorough and constructive review of the paper. The research described in the paper was carried out in part (LLF) by the Jet Propulsion Laboratory, California Institute of Technology, under contract with National Aeronautics and Space Administration. Support from the TOPEX/POSEIDON Project funded under the NASA TOPEX/POSEIDON Announcement of Opportunity is acknowledged.

References

- Arnault, S. and R.E. Cheney, Tropical Atlantic sea level variability from Geosat (1985–1989), *J. Geophys. Res.*, **99**, 18207–18223, 1994.
- Arnault, S., Y. Menard and J. Merle, Observing the Tropical Atlantic Ocean in 1986–1987 from Altimetry, *J. Geophys. Res.*, **95**(C10), 17921–17945, 1990.
- Arnault, S., L. Gourdeau, and Y. Menard, Comparison of the Altimetric Signal with In Situ Measurements in the Tropical Atlantic Ocean, *Deep-Sea Res.*, **39**(3/4), 481–499, 1992 a.
- Arnault, S., A. Morliere, J. Merle and Y. Menard, Low-frequency variability of the Tropical Atlantic surface topography: altimetry and models comparison, *J. Geophys. Res.*, **97**, 14259–14288, 1992 b.
- Behringer, D.W., Sea surface height variations in the Atlantic Ocean: A comparison of TOPEX altimeter data with results from an ocean data assimilation system, *J. Geophys. Res.*, **99**, 24685–24690, 1994.
- Bennett, A.F., *Inverse Methods in Physical Oceanography*, 346 pp., Cambridge University Press, Cambridge, 1992.
- Berry, P.J. and J.C. Marshall, Ocean modeling studies in support of altimetry, *Dyn. Atmos. Oceans*, **13**, 269–300, 1989.
- Bertiger, W. and 19 co-authors, GPS Precise Tracking of Topex/Poseidon: Results and Implications, *J. Geophys. Res.*, **99**, 24449–24464, 1994.
- Bhaskaran, S., G.S. Lagerloef, G.H. Born, W.J. Emery and R.R. Leben, Variability in the Gulf of Alaska from Geosat Altimetry Data, *J. Geophys. Res.*, **98**, 16311–16330, 1993.
- Bisagni, J.J., Ocean surface topography measured by the Geosat radar altimeter during the Frontal Air-Sea Interaction Experiment, *J. Geophys. Res.*, **96**, 22087–22099, 1991.
- Blahe, J. and B. Lunde, Calibrating Altimetry to Geopotential Anomaly and Isotherm Depths in the Western North Atlantic, *J. Geophys. Res.*, **97**(C5), 7465–7478, 1992.
- Blayo, E., J. Verron, and J.M. Molines, Assimilation of TOPEX/POSEIDON altimeter data into a circulation model of the North Atlantic, *J. Geophys. Res.*, **99**, 24691–24705, 1994.
- Born, G.H., B.D. Tapley, J.C. Ries and R.H. Stewart, Accurate Measurement of Mean Sea Level Changes by Altimetric Satellites, *J. Geophys. Res.*, **91**(C10), 11775–11782, 1986.
- Born, G.H., J.L. Mitchell and G.A. Heyler, Geosat ERM—Mission Design, *J. Astron. Sci.*, **35**(2), 119–134, 1987.
- Bourles, B., S. Arnault and C. Provost, Towards Altimetric Data Assimilation in a Tropical Atlantic Model, *J. Geophys. Res.*, **97**(C12), 20271–20284, 1992.
- Brown, O. and R.E. Cheney, Advances in Satellite Oceanography, *Rev. Geophys. Space Phys.*, **21**, 1216–1230, 1983.
- Bruce, J.G., D.R. Johnson and J.C. Kindle, Evidence for eddy formation in the eastern Arabian Sea during the northeast monsoon, *J. Geophys. Res.*, **99**, 7651–7664, 1994.
- Busalacchi, A.J., M.J. McPhaden, and J. Picaut, Variability in equatorial Pacific sea surface topography during the verification phase of the TOPEX/POSEIDON mission, *J. Geophys. Res.*, **99**, 24725–24738, 1994.
- Carnes, M.R., J.L. Mitchell and P.W. de Witt, Synthetic Temperature Profiles Derived from Geosat Altimetry: Comparison with Air-dropped Expendable Bathythermograph Profiles, *J. Geophys. Res.*, **95**(C3), 17979–17992, 1990.
- Carton, J.A., Estimates of Sea Level in the Tropical Atlantic Ocean Using Geosat Altimetry, *J. Geophys. Res.*, **94**(C6), 8029–8039, 1989.
- Carton, J.A. and E.J. Katz, Estimates of the Zonal Slope and the Seasonal Transport of the Atlantic North Equatorial Countercurrent, *J. Geophys. Res.*, **95**(C3), 3091–3100, 1990.
- Carton, J.A., G.A. Chepurin, G.K. Korotaev and Tong Zhu, Comparisons of Dynamic Height Variations in the Tropical Atlantic During 1987–1989 as Viewed in Sections Hydrography and Geosat Altimetry, *J. Geophys. Res.*, **98**(C8), 14369–14378, 1993.
- Cartwright, D.E., and R.D. Ray, Oceanic tides from Geosat altimetry, *J. Geophys. Res.*, **95**, 3069–3090, 1990.
- Chao, Y., D. Halpern and C. Perigaud, Sea Surface Height Variability During 1986–1988 in the Tropical Pacific Ocean, *J. Geophys. Res.*, **98**(C4), 6947–6959, 1993.
- Chelton, D.B., *WOCE/NASA Altimeter Algorithm Workshop*, U.S. WOCE Tech. Rpt. No. 2, 70 pp., US Planning Office for WOCE, Texas A&M University, College Station, TX, 1988.
- Chelton, D.B. and M.G. Schlax, The resolution capability of an irregularly sampled dataset: with application to Geosat altimeter data, *J. Atmos. Oceanic Tech.*, **11**, 534–550, 1994.
- Chelton, D.B., M.G. Schlax, D.L. Witter and J.G. Richman, Geosat Altimeter Observations of the Surface Circulation of the Southern Ocean, *J. Geophys. Res.*, **95**(C10), 17877–17903, 1990.
- Cheney, R.E. and B.C. Douglas, Geographically Correlated Orbit Error in Altimetric Sea Level Time Series, in *Proceedings of the WOCE/NASA Altimeter Algorithm Workshop*, edited by D.B. Chelton, Oregon State University, Corvallis, Oregon, 1988.
- Cheney, R.E. and J.L. Lillibridge, ERS-1 fast delivery altimeter data for tropical ocean monitoring, *EOS Trans. AGU*, **74** (14), 84, 1992.
- Cheney, R.E. and L. Miller, Mapping the 1986–1987 El Niño with Geosat Altimeter data, *EOS Trans. AGU*, **69**(31), 754–755, 1988.
- Cheney, R.E. and L. Miller, Recovery of the Sea Level Signal in the Western Tropical Pacific from Geosat Altimetry, *J. Geophys. Res.*, **95**(C3), 2977–2984, 1990.
- Cheney, R.E., B.C. Douglas and L. Miller, Evaluation of Geosat Altimeter Data with Application to Tropical Pacific Sea Level Variability, *J. Geophys. Res.*, **94**(C4), 4737–4747, 1989.
- Cheney, R.E., W.J. Emery, B.J. Haines and F. Wentz, Recent improvements in Geosat altimeter data, *EOS Trans. AGU*, **72** (51), 577–580, 1991.
- Cheney, R.E., L. Miller, R. Agreen, N. Doyle, and J. Lillibridge, TOPEX/POSEIDON: the 2-cm solution, *J. Geophys. Res.*, **99**, 24555–24563, 1994.
- De Mey, P., Synoptic Estimates of an Eddy Field in the North Atlantic Current, *Oceanologica Acta*, **15**(5), 537–544, 1992.
- De Mey, P. and Y. Menard, Synoptic Analysis and Dynamical Adjustment of Geos 3 and Seasat Altimeter Eddy Fields in the Northwest Atlantic, *J. Geophys. Res.*, **94**, 6221–6230, 1989.
- De Mey, P. and A.R. Robinson, Assimilation of Altimeter Eddy Fields Into a Limited-Area Quasi-Geostrophic Model, *J. Phys. Oceanogr.*, **17**(12), 2280–2293, 1987.
- Delcroix, T., J. Picaut and G. Eldin, Equatorial Kelvin and Rossby Waves

- Evidenced in the Pacific Ocean through Geosat Sea-Level and Surface-Current Anomalies, *J. Geophys. Res.*, *96*, 3249–362, 1991.
- Denker, H. and R.H. Rapp, Geodetic and Oceanographic Results from the Analysis of One Year of Geosat Data, *J. Geophys. Res.*, *95*(C8), 13151–13168, 1990.
- Didden, N. and F. Schott, Seasonal Variations in the Western Tropical Atlantic: Surface Circulation from Geosat Altimetry and WOCE Model Results, *J. Geophys. Res.*, *97*(C3), 3529–3542, 1992.
- Didden, N. and F. Schott, Eddies in the North Brazil Current retroflection region observed by Geosat altimeter data, *J. Geophys. Res.*, *98*(C11), 20121–20131, 1993.
- Dombrowski, E. and P. De Mey, Continuous assimilation in an open domain of the northeast Atlantic 1. Methodology and application to Athena-88, *J. Geophys. Res.*, *97*, 9719–9731, 1992.
- Douglas, B.C. and R.E. Cheney, Geosat: Beginning a New Era in Satellite Oceanography, *J. Geophys. Res.*, *95*(C3), 2833–2836, 1990.
- Douglas, B.C., D.C. McAdoo, and R.E. Cheney, Oceanographic and geophysical applications of satellite altimetry, *Rev. Geophys.*, *25*, 875–880, 1987.
- du Penhoat, Y., T. Delcroix and J. Picaut, Interpretation of Kelvin/Rossby Waves in the Equatorial Pacific from Model-Geosat Data Intercomparison During the 1986–1987 El Niño, *Oceanologica Acta*, *15*(5), 545–554, 1992.
- Egbert, G., A. Bennett and M. Foreman, Topex/Poseidon Tides, *J. Geophys. Res.*, *99*, 24821–24852, 1994.
- Emery, W.J., G.H. Born, D.G. Baldwin and C.L. Norris, Satellite-derived Water Vapor Corrections for Geosat Altimetry, *J. Geophys. Res.*, *95*, 2953–2964, 1990.
- Ezer, T. and G.L. Mellor, Continuous assimilation of Geosat altimeter data into a three-dimensional primitive equation Gulf Stream model, *J. Phys. Oceanogr.*, *24*, 832–847, 1994.
- Ezer, T., G.L. Mellor, D.S. Ko and Z. Sirkes, A comparison of Gulf Stream sea surface height fields derived from Geosat altimeter data and those derived from sea surface temperature data, *J. Atmos. Oceanic Tech.*, *10*, 76–87, 1993.
- Forbes, C., K. Leaman, D. Olson and O. Brown, Eddy and wave dynamics in the South Atlantic as diagnosed from Geosat altimeter data, *J. Geophys. Res.*, *98*, 12297–12314, 1993.
- Fu, L.-L., Recent progress in the application of satellite altimetry to observing the mesoscale variability and general circulation of the oceans, *Rev. Geophys. Space Phys.*, *21*, 1657–1666, 1983.
- Fu, L.-L. and V. Zlotnicki, Observing oceanic mesoscale eddies from Geosat altimetry: preliminary results, *Geophys. Res. Lett.*, *16*, 457–460, 1989.
- Fu, L.-L., J. Vazquez and M. E. Parke, Seasonal variability of the Gulf Stream from satellite altimetry, *J. Geophys. Res.*, *92*, 749–754, 1987.
- Fu, L.-L., M. Lefebvre and E. Christensen, TOPEX/POSEIDON: The Ocean Topography Experiment, *EOS Trans. AGU*, *72*(35), 369–373, 1991 a.
- Fu, L.-L., J. Vazquez and C. Perigaud, Fitting dynamic models to the Geosat sea level observations in the tropical Pacific Ocean. Part 1: A free wave model, *J. Phys. Oceanogr.*, *21*, 798–809, 1991 b.
- Fu, L.-L., I. Fukumori and R.N. Miller, Fitting dynamic models to the Geosat sea level observations in the Tropical Pacific Ocean. Part 2: A linear, wind-driven model, *J. Phys. Oceanogr.*, *23*, 2162–2181, 1993.
- Fu, L.-L., E. J. Christensen, C. Yamarone, M. Lefebvre, Y. Menard, M. Dorrier and P. Escudier, 1994: TOPEX/POSEIDON Mission Overview, *J. Geophys. Res.*, *99*, 24369–24381, 1994.
- Fukumori, I. and P. Malanotte-Rizzoli, An approximate Kalman filter for ocean data assimilation: a reduced-dimension, static, linearized Kalman filter, *J. Geophys. Res.*, *100*, 1995 (in press).
- Fukumori, I., J. Benveniste, C. Wunsch and D. Haidvogel, Assimilation of sea surface topography into an ocean circulation model using a steady-state smoother, *J. Phys. Oceanogr.*, *23*, 1831–1855, 1993.
- Gaspar, P. and C. Wunsch, Estimates from Altimeter Data of Barotropic Rossby Waves in the Northwestern Atlantic Ocean, *J. Phys. Oceanogr.*, *19*(12), 1821–1844, 1989.
- Giese, B.S., J.A. Carton, and L.J. Holl, Sea level variability in the eastern tropical Pacific as observed by TOPEX and Tropical Ocean-Global Atmosphere Tropical Atmosphere-Ocean Experiment, *J. Geophys. Res.*, *99*, 24739–24748, 1994.
- Ghil, M. and P. Malanotte-Rizzoli, Data Assimilation in Meteorology and Oceanography, *Adv. Geophys.*, *33*, 141–266, 1991.
- Gille, S.T., Mean sea surface height of the Antarctic Circumpolar Current from Geosat data: Method and Application, *J. Geophys. Res.*, *99*, 18255–18273, 1994.
- Glenn, S.M., D.L. Porter and A.R. Robinson, A Synthetic Geoid Validation of Geosat Mesoscale Dynamic Topography in the Gulf Stream Region, *J. Geophys. Res.*, *96*, 7145–7166, 1991.
- Gordon, A. and W. Haxby, Agulhas eddies invade the S. Atlantic: evidence from Geosat Altimeter and Shipboard Conductivity–Temperature–Depth Survey, *J. Geophys. Res.*, *95* (C3), 3117–3126, 1990.
- Gourdeau, L., S. Arnault, Y. Menard and J. Merle, Geosat Sea-level Assimilation in a Tropical Atlantic Model Using a Kalman Filter, *Oceanologica Acta*, *15*(5), 567–574, 1992.
- Greiner, E. and C. Perigaud, Assimilation of Geosat Altimetric Data in a Nonlinear Reduced-Gravity Model of the Indian Ocean. Part I: Adjoint Approach and Model-Data Consistency, *J. Phys. Oceanogr.*, *24*, 1783–1804, 1994.
- Haines, B., G.H. Born and C.J. Koblinsky, An estimate of decadal changes in sea surface topography from Seasat and Geosat altimetry, in *Sea Level Changes: Determination and Effects*, Geophysical Monograph Vol. 69, IUGG Vol. 11, pp. 181–187, American Geophysical Union, Washington, D.C., 1992.
- Haines, B.J., G.H. Born, R.G. Williamson and C.J. Koblinsky, Application of the GEM-T2 gravity field to altimetric satellite orbit computation, *J. Geophys. Res.*, *99*, 16237–16254, 1994.
- Haines, K., A Direct Method of Assimilating Sea Surface Height Data into Ocean Models with Adjustments to the Deep Circulation, *J. Phys. Oceanogr.*, *21*(6), 843–868, 1991.
- Hallock, Z.R. and W.J. Teague, Sea Surface Height Fluctuations Observed Simultaneously with Inverted Echo Sounders and Geosat, *J. Geophys. Res.*, *98*(C9), 16341–16350, 1993.
- Hallock, Z.R., J.L. Mitchell and J.D. Thompson, Sea-Surface Topographic Variability Near the New England Seamounts: An Intercomparison Among In-Situ Observations, Numerical Simulations, and Geosat Altimeter from Regional Energetics Experiment, *J. Geophys. Res.*, *94*, 8021–8028, 1989.
- Hansen, D.V. and G.A. Maul, Anticyclonic Current Rings in the Eastern Tropical Pacific, *J. Geophys. Res.*, *96*, 6965–6980, 1991.
- Harangozo, S.A., P.L. Woodworth, R.H. Rapp and Y.M. Yang, A comparison of monthly mean sea level variability determined from Geosat altimetry and a global tide gauge dataset, *Int. J. Remote Sensing*, *14*, 789–796, 1993.
- Holland, W. and P. Malanotte-Rizzoli, Assimilation of Altimeter Data into Ocean Circulation Model: Space Versus Time Resolution Studies, *J. Phys. Oceanogr.*, *19*(10), 1507–1534, 1989.
- Holland, W.R., V. Zlotnicki and L.-L. Fu, Modelled time dependent flow in the Agulhas retroflection region as deduced from altimeter data assimilation, *S. African J. Marine Sci.*, *10*, 407–427, 1991.
- Horton, C.W., D.L. Porter, P.W. DeWitt and W.E. Rankin, Airborne Expendable Bathythermograph Survey of the Kuroshio Extension and Comparison with Simultaneous Altimeter Measurements During Geosat Exact Repeat Mission, *J. Geophys. Res.*, *97*(C5), 7447–7464, 1992.
- Hurlburt, H.E., D.N. Fox and J. Metzger, Statistical inference of weakly correlated subthermocline fields from satellite altimeter data, *J. Geophys. Res.*, *95*, 11375–11409, 1990.
- Ichikawa, K. and S. Imawaki, Life history of a cyclonic ring detached from the Kuroshio Extension as seen by the Geosat altimeter, *J. Geophys. Res.*, *99*, 15953–15966, 1994.
- Jacobs, G.A. and R.R. Leben, Loop Current Eddy Shedding Estimated Using Geosat Altimeter Data, *Geophys. Res. Lett.*, *17*(12), 2385–2388, 1990.
- Jacobs, G.A., G.H. Born, M.E. Parke and P.C. Allen, The Global Structure of the Annual and Semiannual Sea Surface Height Variability from Geosat Altimeter Data, *J. Geophys. Res.*, *97*(C11), 17813–17828, 1992.
- Jacobs, G.A., W.J. Emery and G.H. Born, Rossby waves in the Pacific Ocean extracted from Geosat altimeter data, *J. Phys. Oceanogr.*, *23*, 1155–1175, 1993.
- Jacobs, G.A., H.E. Hurlburt, J.C. Kindle, E.J. Metzger, J.L. Mitchell, W.J. Teague, and A.J. Wallcraft, Decade-scale trans-Pacific propagation and warming effects of an El Niño anomaly, *Nature*, *370* (6488), 360–363, 1994.
- Johnson, D.R., J.D. Thompson and J.D. Hawkins, Circulation in the Gulf of Mexico from Geosat altimetry, *J. Geophys. Res.*, *97*, 2201–2214, 1992.
- Johnson, T.J., R.H. Stewart, C.K. Shum and B.D. Tapley, Distribution of Reynolds stress carried by mesoscale variability in the Antarctic Circumpolar Current, *Geophys. Res. Lett.*, *19*, 1201–1204, 1992.
- Joyce, T.M., K.A. Kelly, D.M. Schubert and M.J. Caruso, Shipboard and

- Altimetric Studies of Rapid Gulf Stream Variability Between Cape Cod and Bermuda, *Deep-Sea Res.*, 37, 897–910, 1990.
- Kelly, K.A., The Meandering of the Gulf Stream as seen by the Geosat Altimeter: Surface Transport, Position and Velocity Variance from 73 deg to 46 deg W, *J. Geophys. Res.*, 96, 16721–16738, 1991.
- Kelly, K.A. and S.T. Gille, Gulf Stream Surface Transport and Statistics at 69° W from the Geosat Altimeter, *J. Geophys. Res.*, 95(C3), 3149–3162, 1990.
- Kelly, K.A. and D.R. Watts, Monitoring Gulf Stream transport by radar altimeter and inverted echo sounders, *J. Phys. Oceanogr.*, 24, 1080–1084, 1994.
- Kelly, K.A., T.M. Joyce, D.M. Schubert and M.J. Caruso, The Mean Sea Surface and Geoid along the Geosat Subtrack from Bermuda to Cape Cod, *J. Geophys. Res.*, 96(C7), 12699–12709, 1991.
- Kelly, K.A., M.J. Caruso and J.A. Austin, Wind-forced variations in sea surface height in the Northeast Pacific Ocean, *J. Phys. Oceanogr.*, 23, 2392–2411, 1993.
- Knudsen, P., Global low harmonic degree models of the seasonal variability and residual ocean tides from TOPEX/POSEIDON altimeter data, *J. Geophys. Res.*, 99, 24643–24655, 1994.
- Koblinsky, C.J., Ocean topography and circulation, in *Atlas of Satellite Observations Related to Global Change*, edited by R.J. Gurney, J.L. Foster, and C.L. Parkinson, pp. 223–236. Cambridge University Press, New York, NY, 1993.
- Koblinsky, C.J., R.S. Nerem, R.G. Williamson and S.M. Klosko, Global scale variations in sea surface topography determined from satellite altimetry, in *Sea Level Changes: Determination and Effects*, Geophysical Monograph Vol. 69, IUGG Vol. 11, pp. 155–165, American Geophysical Union, Washington, D.C., 1992.
- Leben, R.R., G.H. Born, J.D. Thompson and C.A. Fox, Mean sea surface and variability of the Gulf of Mexico using Geosat altimetry data, *J. Geophys. Res.*, 95, 20267–20285, 1990.
- Le Traon, P.Y., Time scales of mesoscale variability and their relationship with space scales in the North Atlantic, *J. Mar. Res.*, 49, 467–492, 1991.
- Le Traon, P.Y., Comments on "Mesoscale Variability in the Atlantic Ocean from Geosat Altimetry and WOCE High-Resolution Numerical Modelling," *J. Phys. Oceanogr.*, 23(12), 2729–2732, 1993.
- Le Traon, P.-Y. and J.-F. Minster, Sea Level Variability and Semiannual Rossby Waves in the South Atlantic Subtropical Gyre, *J. Geophys. Res.*, 98(C7), 12315–12326, 1993.
- Le Traon, P.Y., M.C. Rouquet and C. Boissier, Spatial Scales of Mesoscale Variability in the North Atlantic as Deduced from Geosat Data, *J. Geophys. Res.*, 95, 20267–20286, 1990.
- Lillibridge, J.L., R.E. Cheney and N.S. Doyle, The 1991–93 Los Niños from ERS-1 altimetry, *Proceedings Second ERS-1 Symposium*, pp. 495–499, Hamburg, Germany, 1993.
- Malarde, J.-P., P. De Mey, C. Perigaud and J.-F. Minster, Observations of Long Equatorial Waves in the Pacific Ocean by Seasat Altimetry, *J. Phys. Oceanogr.*, 17, 2273–2279, 1987.
- Marsh, J., et al., A New Gravitational Model for the Earth from Satellite Tracking Data: GEM-T1, *J. Geophys. Res.*, 93(B6), 6169–6215, 1988.
- Marsh, J.G., et al., The GEM-T2 Gravitational Model, *J. Geophys. Res.*, 95(B13), 22043–22070, 1990 a.
- Marsh, J.G., et al., Dynamic Sea Surface Topography, Gravity, and Improved Orbit Accuracies from the Direct Evaluation of Seasat Altimeter Data, *J. Geophys. Res.*, 95(C8), 13129–13150, 1990 b.
- Martel, F. and C. Wunsch, Combined inversion of hydrography, current meter data and altimetric elevations for the North Atlantic circulation, *Manuscripta Geodaetica*, 18, 219–226, 1993.
- Matano, R.P., M.G. Schlax and D.B. Chelton, Seasonal Variability in the Southwestern Atlantic, *J. Geophys. Res.*, 98(C10), 18027–18035, 1993.
- Matthews, P.E., M.A. Johnson and J.J. O'Brien, Observations of Mesoscale Ocean Features in the Northeast Pacific using Geosat Radar Altimetry, *J. Geophys. Res.*, 97(C11), 17829–17840, 1992.
- Maul, G., D.V. Hansen and N.J. Bravo, A note on sea level variability at Clipperton Island from Geosat and in-situ observations, in *Sea Level Changes: Determination and Effects*, Geophysical Monograph 69, IUGG Volume 11, pp. 145–154, American Geophysical Union, Washington, DC, 1992.
- McAdoo, D.C. and K.M. Marks, Gravity fields of the Southern Ocean from Geosat data, *J. Geophys. Res.*, 97, 3247–3260, 1992.
- Mellor, G.L. and T. Ezer, A Gulf Stream Model and an Altimetry Assimilation Scheme, *J. Geophys. Res.*, 96, 8779–8795, 1991.
- Menard, Y., Observing the Seasonal Variability in the Tropical Atlantic from Altimetry, *J. Geophys. Res.*, 93(C11), 13967–13978, 1988.
- Mestas-Nunez, A.M., D.B. Chelton and R.A. DeSzoeko, Evidence of Time-Dependent Sverdrup Circulation in the South Pacific from Seasat Scatterometer and Altimeter, *J. Phys. Oceanogr.*, 22(8), 934–943, 1992.
- Miller, L. and R. Cheney, Large-Scale Meridional Transport in the Tropical Pacific Ocean during the 1986–1987 El Niño from Geosat, *J. Geophys. Res.*, 95(C10), 17905–17920, 1990.
- Miller, L., R.E. Cheney and B.C. Douglas, Geosat Altimeter Observations of Kelvin Waves and the 1986–1987 El Niño, *Science*, 239, 52–54, 1988.
- Miller, R.N., Direct Assimilation of Altimetric Differences Using the Kalman Filter, *Dyn. Atmos. Oceans*, 13, 317–333, 1989.
- Mitchell, J.L., J.M. Dastugue, W.J. Teague and Z.R. Hallock, The Estimation of Geoid Profiles in the Northwest Atlantic from Simultaneous Satellite Altimetry and Airborne Expendable Bathythermograph Sections, *J. Geophys. Res.*, 95(C10), 17965–17978, 1990.
- Moore, A.M., Data assimilation in a quasi-geostrophic open ocean model of the Gulf Stream region using the adjoint method, *J. Phys. Oceanogr.*, 21, 398–427, 1991.
- Morrow, R., J. Church, R. Coleman, D. Chelton and N. White, Eddy momentum flux and its contribution to the Southern Ocean momentum balance, *Nature*, 357, 482–484, 1992.
- Morrow, R., R. Coleman, J. Church and D. Chelton, Surface eddy momentum flux and velocity variances in the Southern Ocean from Geosat altimetry, *J. Phys. Oceanogr.*, 24, 2050–2071, 1994.
- Musman, S., Sea height wave form in equatorial waves and its interpretation, *J. Geophys. Res.*, 94, 3303–3309, 1989.
- Musman, S., Geosat Altimeter Observations of Long Waves in the Equatorial Atlantic, *J. Geophys. Res.*, 97(C3), 3573–3580, 1992.
- Musman, S., A. Drew and B. Douglas, Ionospheric Effects on Geosat Altimeter Observations, *J. Geophys. Res.*, 95, 2965–2968, 1990.
- Nerem, R.S., B.D. Tapley and C.-K. Shum, Determination of the Ocean Circulation using Geosat Altimetry, *J. Geophys. Res.*, 95(C3), 3163–3180, 1990.
- Nerem, R.S., E. Schrama, C. Koblinsky and B. Beckley, A preliminary evaluation of ocean topography from the TOPEX/POSEIDON Mission, *J. Geophys. Res.*, 99, 24565–24583, 1994.
- Okkonen, S.R., The Shedding of an Anticyclonic Eddy from the Alaskan Stream as Observed by the Geosat Altimeter, *Geophys. Res. Lett.*, 19(24), 2397–2400, 1992.
- Okkonen, S.R., Observations of topographic planetary waves in the Bering Slope Current using the Geosat altimeter, *J. Geophys. Res.*, 98, 22603–22613, 1993.
- Pares-Sierra, A., W.B. White and C.-K. Tai, Wind-driven coastal generation of annual mesoscale eddy activity in the California Current, *J. Phys. Oceanogr.*, 23, 1110–1121, 1993.
- Perigaud, C., Sea-Level Oscillations Observed with Geosat along Two Shear Fronts of the Pacific North Equatorial Counter Current, *J. Geophys. Res.*, 95(C5), 7239–7248, 1990.
- Perigaud, C. and P. Delecluse, Simulations of Dynamic Topography in the Northwestern Indian Ocean with Input of Seasat Altimeter and Scatterometer Data, *Ocean-Air Interactions*, 1, 289–309, 1989.
- Perigaud, C. and P. Delecluse, Annual Sea Level Variations in the Southern Tropical Indian Ocean from Geosat and Shallow Water Simulations, *J. Geophys. Res.*, 97(C12), 20169–20178, 1992.
- Perigaud, C. and P. Delecluse, Interannual sea level variations in the Tropical Indian Ocean from Geosat and shallow water simulations, *J. Phys. Oceanogr.*, 23, 1916–1934, 1993.
- Perigaud, C. and J.-F. Minster, Variability of the Somali Current as Observed from Seasat Altimetry, *J. Phys. Oceanogr.*, 18(1), 25–39, 1988.
- Picaud, J., A.J. Busalacchi, M.J. McPhaden and B. Camusat, Validation of the Geostrophic Method for Estimating Zonal Currents at the Equator from Geosat Altimeter Data, *J. Geophys. Res.*, 95(C3), 3015–3024, 1990.
- Porter, D.L., A.R. Robinson, S.M. Glenn and E.B. Dobson, The Synthetic Geoid and the Estimation of Mesoscale Absolute Topography from Altimeter Data, *Technical Digest*, 10(4), 369–379, The Johns Hopkins University Applied Physics Lab., 1989.
- Porter, D.L., E. Dobson and S. Glenn, Measurements of sea surface topography during SYNOP utilizing a Geosat synthetic geoid, *Geophys. Res. Lett.*, 19, 1847–1850, 1992.
- Provost, C. and P.-Y. Le Traon, Spatial and Temporal Scales in Altimetric Variability in the Brazil-Malvinas Current Confluence Region: Dominance of the Semiannual Period and Large Spatial Scales, *J. Geophys. Res.*, 98(C10), 18037–18051, 1993.

- Qiu, B., Recirculation and Seasonal Change of the Kuroshio from Altimetry Observations, *J. Geophys. Res.*, 97(C11), 17801–17813, 1992.
- Qiu, B., Determining the mean Gulf Stream and its recirculations through combining hydrographic and altimetric data, *J. Geophys. Res.*, 99, 951–962, 1994.
- Qiu, B. and K.A. Kelly, Upper-ocean heat balance in the Kuroshio Extension region, *J. Phys. Oceanogr.*, 23, 2027–2041, 1993.
- Qiu, B., K.A. Kelly and T.M. Joyce, Mean Flow and Variability in the Kuroshio Extension from Geosat Altimetry data, *J. Geophys. Res.*, 96, 18491–18507, 1991.
- Quartly, G.D. and M.A. Srokosz, Seasonal variations in the region of the Agulhas retroflection: Studies with Geosat and FRAM, *J. Phys. Oceanogr.*, 23, 2107–2124, 1993.
- Rapp, R.H., Geoid Undulation Accuracy, *IEEE Transactions on Geoscience and Remote Sensing*, 31(2), 365–370, 1993.
- Rapp, R.H., and D.A. Smith, Preliminary estimates of Gulf Stream characteristics from TOPEX data and a precise gravimetric geoid, *J. Geophys. Res.*, 99, 24707–24723, 1994.
- Rapp, R.H. and Y.M. Wang, Dynamic Topography Estimates Using Geosat Data and a Gravimetric Geoid in the Gulf Stream Region, *Geophys. J. Int.*, 117, 511–528, 1994.
- Robinson, A.R. and L.J. Walstad, Altimetric data assimilation for ocean dynamics and forecasting, *Technical Digest*, 8, 267–271, The Johns Hopkins University Applied Physics Lab., 1987.
- Robinson, A.R., L.J. Walstad, J. Calman, E.B. Dobson, D.W. Denbo, S.M. Glenn, D.L. Porter and J. Goldhirsh, Frontal Signals East of Iceland from the Geosat Altimeter, *Geophys. Res. Lett.*, 16, 77–80, 1989.
- Sandwell, D.T. and B. Zhang, Global Mesoscale Variability from the Geosat Exact Repeat Mission: Correlation with Ocean Depth, *J. Geophys. Res.*, 94, 17971–17984, 1989.
- Schlag, M. G. and D. B. Chelton, Detecting aliased tidal errors in altimeter height measurements, *J. Geophys. Res.*, 99, 12603–12612, 1994.
- Schroter, J., U. Seiler and M. Wenzel, Variational Assimilation of Geosat Data into an Eddy-resolving Model of the Gulf Stream Extension Area, *J. Phys. Oceanogr.*, 23, 925–953, 1993.
- Shum, C.-K., D.N. Yuan, J.C. Ries, J.C. Smith, B.E. Schutz and B.D. Tapley, Precision Orbit Determination for the Geosat Exact Repeat Mission, *J. Geophys. Res.*, 95(C3), 2887–2898, 1990 a.
- Shum, C.-K., R.A. Werner, D.T. Sandwell, B.H. Zhang, R.S. Nerem and B.D. Tapley, Variations of Global Mesoscale Eddy Energy Observed from Geosat, *J. Geophys. Res.*, 95(C10), 17865–17876, 1990 b.
- Smedstad, O.M. and D.N. Fox, Assimilation of Altimeter data in a two-layer primitive equation model of the Gulf Stream, *J. Phys. Oceanogr.*, 24, 305–325, 1994.
- Stammer, D. and C.W. Boning, Mesoscale Variability in the Atlantic Ocean from Geosat Altimetry and WOCE High-Resolution Numerical Modelling, *J. Phys. Oceanogr.*, 22(7), 732–752, 1992.
- Stammer, D. and C.W. Boning, Reply, *J. Phys. Oceanogr.*, 23(12), 2733–2735, 1993.
- Stammer, D. and C. Wunsch, Preliminary assessment of the accuracy and precision of TOPEX/POSEIDON altimeter data with respect to the large-scale ocean circulation, *J. Geophys. Res.*, 99, 24584–24604, 1994.
- Stammer, D., H.H. Hinrichsen and R.H. Kase, Can Meddies be Detected by Satellite Altimetry, *J. Geophys. Res.*, 96, 7005–7014, 1991.
- Stewart, R.H., *Methods of Satellite Oceanography*, 360 pp., University of California Press, Berkeley, CA, 1985.
- Tai, C.K., Estimating the Basin-Scale Circulation from Satellite Altimetry. Part 1 : Straightforward Spherical Harmonic Expansion, *J. Phys. Oceanogr.*, 18(10), 1398–1413, 1988.
- Tai, C.-K., Estimating the Surface Transport of Meandering Oceanic Jet Streams from Satellite Altimetry: Surface Transport Estimates for the Gulf Stream and Kuroshio Extension, *J. Phys. Oceanogr.*, 20(6), 860–879, 1990.
- Tai, C.-K. and W. White, Eddy Variability in the Kuroshio Extension as Revealed by Satellite Altimetry: Energy Propagation Away from the Jet, *J. Phys. Oceanogr.*, 20, 1761–1777, 1990.
- Tai, C.-K., W.B. White and S.E. Pazan, Geosat Crossover Analysis in the Tropical Pacific 2 : Verification Analysis of Altimetric Sea Level Maps with Expendable Bathythermograph and Island Sea-Level data, *J. Geophys. Res.*, 94(C1), 897–908, 1989.
- Tapley, B.D., R.S. Nerem, C.K. Shum, J.C. Ries and D.N. Yuan, Determination of the General Ocean Circulation from a Joint Gravity Field Solution, *Geophys. Res. Lett.*, 15, 1109–1112, 1988.
- Tapley, B., C. Shum, J. Ries, R. Suter and B. Schutz, Monitoring of changes in global mean sea level using Geosat altimeter, in *Sea Level Changes: Determination and Effects*, Geophysical Monograph, 69, IUGG Vol. 11, pp. 167–180, American Geophysical Union, Washington, DC, 1992.
- Tapley, B.D., D.P. Chambers, C.K. Shum, R.J. Eanes, J.C. Ries, and R.H. Stewart, Accuracy assessment of the large-scale dynamic topography from TOPEX/POSEIDON altimetry, *J. Geophys. Res.*, 99, 24605–24617, 1994.
- Tokmakian, R.T. and P.G. Challenor, Observations in the Canary Basin and the Azores frontal region using Geosat data, *J. Geophys. Res.*, 98, 4761–4773, 1993.
- TOPEX Science Working Group, *Satellite Altimetric Measurements of the Ocean*, JPL Internal Doc. 400–111, 78 pp., Jet Propulsion Laboratory, Pasadena, CA, 1981.
- TOPEX/POSEIDON Science Working Team, *TOPEX/POSEIDON Science Investigations Plan*, JPL Publication 91–27, 176 pp., Jet Propulsion Laboratory, Pasadena, CA, 1991.
- van Woert, M.L. and J.M. Price, Geosat and Advanced Very High Resolution Radiometer Observations of Oceanic and Planetary Waves Adjacent to the Hawaiian Islands, *J. Geophys. Res.*, 98(C8), 14619–14650, 1993.
- Vazquez, J., Observations on the long-period variability of the Gulf Stream downstream of Cape Hatteras, *J. Geophys. Res.*, 98, 20133–20147, 1993.
- Vazquez, J., V. Zlotnicki and L.-L. Fu, Sea level variabilities in the Gulf Stream between Cape Hatteras and 50° W—A Geosat Study, *J. Geophys. Res.*, 95, 17957–17964, 1990.
- Verron, J., Altimeter Data Assimilation Into Ocean Model: Sensitivity to Orbital Parameters, *J. Geophys. Res.*, 95, 11443–11459, 1990.
- Verron, J., Nudging Satellite Altimeter Data into Quasi-Geostrophic Ocean Models, *J. Geophys. Res.*, 97(C5), 7479–7492, 1992.
- Verron, J., J.-M. Molines, and E. Blayo, Assimilation of Geosat Data into a Quasi-geostrophic Model of the North Atlantic between 20° and 50° N: Preliminary Results, *Oceanologica Acta*, 15(5), 575–583, 1992.
- Visser, P.N.A.M., K.F. Wakker and B.A.C. Ambrosius, Dynamic Sea Surface Topography from GEOSAT Altimetry, *Marine Geodesy*, 16(3), 215–239, 1993.
- Wagner, C.A. and R.E. Cheney, Global Sea Level Change from Satellite Altimetry, *J. Geophys. Res.*, 97(C10), 15607–15616, 1992.
- Wagner, C.A. and C.K. Tai, Degradation of ocean signals in satellite altimetry due to orbit removal processes, *J. Geophys. Res.*, 99, 16255–16267, 1994.
- Wakker, K.F., C.A. Zandbergen, M.C. Naeije and B.A. Ambrosius, Geosat Altimeter Data Analysis for the Oceans around South Africa, *J. Geophys. Res.*, 95(C3), 2991–3006, 1990.
- White, W.B., N. Graham and C.-K. Tai, Reflection of Annual Rossby Waves at the Maritime Western Boundary of the Tropical Pacific, *J. Geophys. Res.*, 95(C3), 3101–3116, 1990 a.
- White, W.B., C.-K. Tai and J. DiMento, Annual Rossby Wave Characteristics in The California Current Region from the Geosat Exact Repeat Mission, *J. Phys. Oceanogr.*, 20(9), 1297–1311, 1990 b.
- White, W.B., C.-K. Tai and W.R. Holland, Continuous Assimilation of Geosat Altimetric Sea Level Observations into a Numerical Synoptic Ocean Model of the California Current, *J. Geophys. Res.*, 95(C3), 3127–3148, 1990 c.
- Willebrand, J., R.H. Kase, D. Stammer, H.-H. Hinrichsen and W. Krauss, Verification of the Geosat Altimeter Sea Surface Topography in the Gulf Stream Extension with Surface Drifting Buoys and Hydrographic Measurements, *J. Geophys. Res.*, 95(C3), 3007–3014, 1990.
- Wunsch, C., Global-Scale Sea Surface Variability from Combined Altimetric and Tide Gauge Measurements, *J. Geophys. Res.*, 96, 15053–15082, 1991 a.
- Wunsch, C., Large-scale response of the ocean to atmospheric forcing at low frequencies, *J. Geophys. Res.*, 96, 15083–15092, 1991 b.
- Wunsch, C., Observing ocean circulation from space, *Oceanus*, 35 (2), 9–17, 1992.
- Wunsch, C. and E.M. Gaposchkin, On Using Satellite Altimetry to Determine the General Circulation of the Oceans with Application to Geoid Improvement, *Rev. Geophys. Space Phys.*, 18(4), 725–745, 1980.
- Zlotnicki, V., Sea Level Differences Across the Gulf Stream and Kuroshio Extension, *J. Phys. Oceanogr.*, 21(4), 599–609, 1991.
- Zlotnicki, V., L.-L. Fu and W. Patzert, Seasonal variability in global sea level observed with Geosat altimetry, *J. Geophys. Res.*, 94, 17959–17969, 1989.

Zlotnicki, V., G. Siedler and B. Klein, Can the Weak Surface Currents of the Cape Verde Frontal Zone Be Measured with Altimetry? *J. Geophys. Res.*, 98(C2), 2485-2493, 1993.

L.-L. Fu, Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, CA 91109-8099. (email: llf@pacific.jpl.nasa.gov)

R. E. Cheney, National Ocean Service, National Oceanic and Atmospheric Administration, 1305 East-West Highway, Silver Spring, MD 20910.

(Received June 10, 1994; revised October 20, 1994; accepted November 21, 1994.)