

The Statistical Analysis of a Random, Moving Surface Author(s): M. S. Longuet-Higgins Source: Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences, Vol. 249, No. 966 (Feb. 21, 1957), pp. 321-387 Published by: The Royal Society Stable URL: <u>http://www.jstor.org/stable/91668</u> Accessed: 21/10/2013 05:45

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



*The Royal Society* is collaborating with JSTOR to digitize, preserve and extend access to *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences.* 

http://www.jstor.org

# THE STATISTICAL ANALYSIS OF A RANDOM, MOVING SURFACE

### By M. S. LONGUET-HIGGINS

National Institute of Oceanography, Wormley

# (Communicated by G. E. R. Deacon, F.R.S.—Received 29 March 1956— Revised 31 July 1956)

The following statistical properties are derived for a random, moving, Gaussian surface: (1) the probability distribution of the surface elevation and of the magnitude and orientation of the gradient; (2) the average number of zero-crossings per unit distance along a line in an arbitrary direction; (3) the average length of the contours per unit area, and the distribution of their direction; (4) the average density of maxima and minima per unit area of the surface, and the average density of specular points (i.e. points where the two components of gradient take given values); (5) the probability distribution of the velocities of zero-crossings along a given line; (6) the probability distribution of the velocities of contours and of specular points; (7) the probability distribution of the heights of maxima and minima and the distribution of the intervals between successive zero-crossings along an arbitrary line. All the results are expressed in terms of the two-dimensional energy spectrum of the surface, and are found to involve the moments of the spectrum up to a finite order only. (1), (3), (4), (5) and (6) are discussed in detail for the special case of a narrow spectrum.

The converse problem is also studied and solved: given certain statistical properties of the surface, to find a convergent sequence of approximations to the energy spectrum.

The problems arise in connexion with the statistical analysis of the sea surface.

(More detailed summaries are given at the beginning of each part of the paper.)

#### CONTENTS

	PAGE		PAGE
INTRODUCTION	321	$2 \cdot 3$ . The length and direction of the	
		contours	<b>345</b>
PART I. DESCRIPTION OF THE SURFACE	324	$2 \cdot 4$ . The density of maxima and minima	<b>348</b>
$1 \cdot 1$ . The representation of simple wave		2.5. The velocities of zeros along a line	356
patterns	325	$2 \cdot 6$ . The motion of the contours	359
1.2. The representation of a surface		2.7. The velocities of specular points	362
having a continuous spectrum	327	$2 \cdot 8$ . Properties of the envelope: the	
1.3. Conditions for degeneracy	328	number of waves in a group	370
1.4. The spectrum of the surface in an		2.9. The heights of maxima	<b>374</b>
arbitrary direction	331	$2 \cdot 10$ . The intervals between successive	
1.5. The wave envelope	333	zeros	376
1.6. A narrow spectrum	335	Part III. A method of determining the	
PART II. STATISTICAL PROPERTIES	336	ENERGY SPECTRUM	<b>379</b>
$2 \cdot 1$ . The distribution of surface elevation		3.1. To obtain $m_n(\theta)$	379
and gradient	338	3.2. To obtain $m_{pq}$	380
$2 \cdot 2$ . The number of zero-crossings along		3.3. To obtain $E$	<b>382</b>
a line	342	References	386

### INTRODUCTION

On observing waves in the open ocean, one is struck by their irregularity: no single wave retains its identity for long, the distance between neighbouring crests varies with time and place, and frequently it is difficult to assign to the waves any predominant direction or

Vol. 249. A. 966. (Price £1. 1s.)

40

[Published 21 February 1957

orientation. Thus although the sea surface may, for some purposes, be treated as a uniform train of waves advancing in one direction only, such a representation is usually far from reality.

The first attempt to treat the sea surface as the sum of more than a finite number of simple sine-waves is due to Barber and his collaborators (1946), who used a harmonic analyzer to resolve a length of record, say of wave height or pressure at a fixed point, into its Fourier components. The physical basis for this procedure is that, if the waves are not too steep, the energy in any particular frequency band may be expected to be propagated independently of the rest of the spectrum, and with a velocity characteristic of its frequency. It was shown by Barber & Ursell (1948) that for ocean swell this is in fact nearly true.

Just as sea waves have no single frequency or wavelength, so they have no single direction. One must therefore consider the Fourier spectrum of the sea surface with regard to both frequency and direction or, what is equivalent, the spectrum with regard to wave-number in two horizontal directions. A two-dimensional Fourier analysis for sea waves was proposed by Longuet-Higgins & Barber (1946), who also suggested apparatus for finding a certain amount of information about the spectrum. Independently, Pierson (1952) has emphasized the importance of the distribution of energy with regard to direction when studying the generation and propagation of waves and swell. Thus waves from a limited storm area will decay more or less rapidly with distance according as the spread in direction of the energy is wide or narrow. Similarly, the angular distribution of the energy in a swell will be more or less concentrated according as the region in which it was generated subtends a wide or narrow angle at the point of observation.\*

A very interesting problem now arises: the relation between the energy spectrum of the surface and its observable statistical properties. To take a simple example, suppose that we measure the surface elevation  $\zeta$  at a fixed point: what is the r.m.s. value of  $\zeta$  with regard to time; what is the average time interval between the maxima of  $\zeta$ ; what proportion of the maxima have heights between two given values?

Questions of this kind have been studied theoretically by several authors, notably by Rice (1944, 1945) in connexion with the analysis of electrical noise currents. Rice considered the function  $\zeta(t) = \sum_{n} c_n \cos(\sigma_n t + \epsilon_n), \qquad (1)$ 

which is the sum of a large number of sine-waves of different frequency  $\sigma_n/2\pi$ . The phases  $\epsilon_n$  are random variables distributed uniformly in the interval  $(0, 2\pi)$ , and the amplitudes  $c_n$  are such that in any small interval of  $\sigma$  of width  $d\sigma$ ,

$$\sum_{n} \frac{1}{2} c_n^2 = E(\sigma) \,\mathrm{d}\sigma,\tag{2}$$

say (our notation is slightly different from Rice's). The function  $E(\sigma)$  may be called the energy spectrum of  $\zeta$ . It is the cosine transform of G. I. Taylor's correlation function

$$\psi(t) = \lim_{T \to \infty} \frac{1}{2T} \int_{-T}^{T} \zeta(t') \, \zeta(t'+t) \, \mathrm{d}t' \tag{3}$$

<sup>\*</sup> Some other applications of the two-dimensional spectrum may be mentioned. It has been used to calculate the seismic energy generated by sea waves, where the directional distribution of energy is essentially involved (Longuet-Higgins 1950). Eckart has used a two-dimensional analysis to calculate the scattering of sound from the sea surface (1953*a*) and the waves caused by a random distribution of pressure pulses (1953*b*). St Denis & Pierson (1953) have applied it to ship motion; see also Cartwright (1956).

(see Khintchine 1934). One can show that the method of harmonic analysis used by Barber & Ursell (1948) is essentially equivalent to making an estimate of  $\sqrt{E}$ , within limits of accuracy imposed by the finite length of the record (see Tukey 1949).

Using the above representation, Rice was able to derive many statistical properties of  $\zeta$ , in particular the probability distribution of  $\zeta$  itself (which is Gaussian), the average number of zero-crossings of  $\zeta$  per unit time, the probability distribution of the maxima, and certain statistical properties of the envelope.

It was found by Rudnick (1950) that records of sea-wave pressure are in fact Gaussian (see also Pierson 1952). Barber (1950) considered the distribution of wave heights, that is, the difference in level between a crest and the preceding trough, and compared some observations with the 'random-walk' (or Rayleigh) distribution,\* which is the theoretical distribution for a narrow-band spectrum. This distribution has been discussed in more detail by the present author (1952), who showed that the theoretical ratios of the mean wave height, the mean of the highest one-third of the waves, and the height of the highest of N waves were in close agreement with observation. Further observations are given by Watters (1953).

Some two-dimensional statistical properties of the sea surface have also been measured. By photographing the pattern and intensity of sunlight reflected from the sea surface, Cox & Munk (1954a, b) have deduced the statistical distribution of the two components of surface slope, in winds of different intensity. They find that the distribution differs only slightly from a normal distribution.<sup>†</sup> One may expect that for swell, which is usually less steep than waves under the action of the wind, the departures from the normal distribution will be still less.

For more than fifty years attempts have been made to construct contour maps of the sea surface. Some results, together with references to earlier work, are given by Schumacher (1952). At the present time some very extensive maps are being made as proposed by Marks (1954). These maps may well be suitable for statistical analysis.

On the theoretical side, Eckart (1946) has considered the intensity of light reflected from a random surface whose gradient and second derivatives are all distributed normally; and he has also calculated the first and second moments of the total curvature. However, no extensive theoretical study of the two-dimensional statistical properties of a random surface appears to have been made.

The purpose of the present paper is to study theoretically the statistical properties of a random, moving Gaussian surface, in relation to its two-dimensional spectrum.

In view of the observations mentioned above, there is reason to believe that some at least of the results are relevant to waves in the open ocean. The analysis may also apply to other geophysical phenomena, for example, to microseisms or perturbations of the ionosphere. In addition, however, the subject is of interest as a branch of geometry, and we shall develop it here on its own account, leaving the application of the results and comparison with observations to a separate study.

\* So called because it was derived by Rayleigh in connexion with the theory of sound. See Rayleigh (1880; 1945, pp. 39-42).

<sup>†</sup> Schooley (1954) has made similar measurements for the river Anacostia. A different technique was used earlier by Duntley (1950) on Lake Winnipeg.

40-2

#### M. S. LONGUET-HIGGINS ON THE

The paper is in three parts. Part I is mainly introductory; we define some convenient parameters for describing the surface: the *long-crestedness*, the *skewness*, the *carrier wave* and the *envelope*, and we find conditions for the surface to split up in various ways into one or more simpler systems of waves.

The chief results are contained in part II. Expressions are derived for the statistical distributions of the surface elevation and the magnitude and direction of the surface slope  $(\S 2 \cdot 1)$ ; for the average number of zero-crossings of  $\zeta$  along a line in an arbitrary direction  $(\S 2 \cdot 2)$ ; for the average length of a given contour and for the distribution of its direction  $(\S 2 \cdot 3)$ ; for the density of maxima and minima (humps and hollows) per unit area of the surface, and the density of specular points (points where the two components of surface gradient take given values)  $(\S 2 \cdot 4)$ ; for the statistical distribution of the velocities of the zero-crossings of  $\zeta$  along a given line  $(\S 2 \cdot 5)$ ; for the statistical distributions of the velocities of the contours  $(\S 2 \cdot 6)$  and of specular points  $(\S 2 \cdot 7)$ . In order to interpret the more complex results, the case when the energy spectrum is narrow, i.e. when the waves are more or less uniform in wavelength and direction, is studied in detail. In  $\S 2 \cdot 8$  some properties of the wave envelope are considered, and from these are deduced the average number of waves in a group, the statistical distribution of the heights of maxima and the distribution of the spacing between successive zeros, all for a narrow spectrum.

In part III the converse problem is considered: given the statistical properties of the surface, to find its energy spectrum. To do this, use is made of a striking feature of the present distributions, that they depend only on the moments of the energy spectrum up to a finite order. Thus the average number of zero-crossings along a line involves only the moments of order 0 and 2. The average number of maxima and minima along a line involves only the moments of order 2 and 4. Properties depending on the motion of the surface involve the odd as well as the even moments. Hence, by considering the statistical properties of the surface along a line in a number of different directions, the moments of the two-dimensional spectrum up to, theoretically, any order can be deduced. From this it is possible to obtain a convergent sequence of approximations to the spectrum  $(\S 3 \cdot 3)$ .

Detailed summaries of the results will be found at the beginning of each part.

## PART I. DESCRIPTION OF THE SURFACE

Section  $1 \cdot 1$  introduces the representation of a simple wave pattern by a point in the wave-number diagram, and defines the concepts of *carrier wave* and *envelope*, which are afterwards to be extended to a surface with a continuous spectrum. The fundamental definition of a random surface in terms of its spectrum is given in §1.2.

In §1·3 conditions are found for the surface to degenerate in various ways. Thus, a simple condition for the surface to be 'long-crested' (i.e. for the energy to travel always in the same direction) is given by (1·3·3). A condition for the surface to consist of no more than two long-crested systems is given by (1·3·7), and a condition for no more than n such systems is given by (1·3·8). All these conditions are expressed in terms of the moments  $m_{pq}$  of the energy spectrum E, which are defined by (1·2·7). A condition for E to degenerate into a 'ring' spectrum, i.e. for the energy to have uniform wavelength though not necessarily constant direction, is given by (1·3·11). For standing waves, both (1·3·3) and (1·3·11) must be satisfied simultaneously. Necessary and sufficient conditions for the spectrum to be *narrow* so that the energy is uniform in both wavelength and direction, are given by (1·3·14). Necessary conditions for the existence of not more than two narrow bands of energy are given by (1·3·17) and (1·3·18).

In  $\S1.4$  the curve of intersection of the surface by a perpendicular plane in an arbitrary direction  $\theta$  is considered. It is shown how the spectrum  $E_{\theta}$  of this curve is related to the spectrum E of the surface. The principal direction is defined as the direction in which the second-order moment of  $E_{a}$ , and so the r.m.s. wave-number, is a maximum. The minimum r.m.s. wave-number is in the perpendicular direction, and the ratios of the r.m.s. wave-numbers in these two directions is a convenient measure of the long-crestedness of the surface.

In §1.5 the carrier wave and the wave envelope are defined for a surface having a continuous spectrum. It is seen that in general the principal direction of the envelope is different from that of the wave surface, so that the waves form a 'staggered' or echelon pattern. The angle between the two principal directions is called the angle of skewness. It is proved that the envelope of the curve in which a vertical plane intersects the surface is the same as the curve in which the plane intersects the envelope.

In  $\S16$  some special properties of a narrow spectrum are deduced; in particular, that the longcrestedness equals the reciprocal of the r.m.s. angular deviation of energy from the principal direction.

### $1 \cdot 1$ . The representation of simple wave patterns

Imagine a single long-crested wave of length  $\lambda$  travelling in a direction which makes an angle  $\theta$  with the x axis (see figure 1a). The wave-number w along a line perpendicular to the crest is defined as

$$w = 2\pi/\lambda. \tag{1.1.1}$$

The wavelength and direction can be specified very conveniently by drawing a vector  $\vec{OP}$ from a fixed point O in a direction  $\theta$ , such that the length of  $\overrightarrow{OP}$  equals w. Then if we consider a section of the surface along any line making an angle  $\theta'$  with the x axis, it is clear that the wavelength along this section is increased in the ratio sec  $(\theta - \theta')$ , so that the wavenumber is multiplied by  $\cos(\theta - \theta')$ . In other words the corresponding wave-number is simply the projection of  $\overrightarrow{OP}$  on a line in that direction. In particular, the wave-numbers parallel to the two fixed directions (x, y) are the co-ordinates of the point P with respect of axes in these directions. The equation of the wave surface is then

$$\zeta = c\cos\left(ux + vy + \sigma t\right), \tag{1.1.2}$$

where

$$u, v = w \cos \theta, \ w \sin \theta, \tag{1.1.3}$$

and  $\sigma$  is a function of u and v. It will be assumed that  $\sigma$  depends only upon the wavelength, that is on  $\sqrt{(u^2+v^2)}=w$ ; π

$$= \sigma(u, v) = \sigma(w). \tag{1.1.4}$$

(1.1.7)

We may take  $\sigma$  to be positive, so that the direction of propagation is opposite to  $\overrightarrow{OP}$ . It follows from  $(1 \cdot 1 \cdot 4)$  that  $\sigma(-u,-v)=\sigma(u,v),$  $(1 \cdot 1 \cdot 5)$ 

that is, waves of the same length but opposite in direction have equal and opposite velocities.

Consider now a pair of long-crested waves of equal amplitude c (figure 1 b). If these are represented in the wave-number diagram by the vectors  $\vec{OP_1}$  and  $\vec{OP_2}$ , where  $P_1 = (u_1, v_1)$ and  $P_2 = (u_2, v_2)$ , we have for the surface elevation

 $(-2c\cos(u'x+v'u+\sigma't)\cos(\overline{u}x+\overline{v}u+\overline{\sigma}t))$ 

$$\zeta = c \cos(u_1 x + v_1 y + \sigma_1 t) + c \cos(u_2 x + v_2 y + \sigma_2 t).$$
 (1.1.6)

This may be written

$$u' = \frac{1}{2}(u_1 - u_2), \quad v' = \frac{1}{2}(v_1 - v_2), \quad \sigma' = \frac{1}{2}(\sigma_1 - \sigma_2),$$

where

$$\bar{u} = \frac{1}{2}(u_1 + u_2), \quad \bar{v} = \frac{1}{2}(v_1 + v_2), \quad \bar{\sigma} = \frac{1}{2}(\sigma_1 + \sigma_2).$$
(1.1.8)

If the wave-numbers  $(u_1, v_1)$  and  $(u_2, v_2)$  of the two original waves are nearly equal, the term  $2c \cos(u'x + v'y + \sigma't)$  (1.1.9)

in  $(1 \cdot 1 \cdot 7)$  represents a slowly varying amplitude which we may call the 'envelope' and

$$\cos\left(\bar{u}x + \bar{v}y + \bar{\sigma}t\right) \tag{1.1.10}$$

represents a 'carrier' wave of approximately the same wavelength and direction as the original waves. The carrier wave is represented in the wave-number diagram by the vector  $\overrightarrow{OM}$ , where M is the mid-point of  $P_1P_2$ . The envelope is represented by  $\overrightarrow{MP}_1$  or  $\overrightarrow{MP}_2$ .



FIGURE 1. Representation in the wave-number plane of (a) a single long-crested wave and (b) the sum of two long-crested waves of different wavelength and direction.

For example, suppose that the two waves are in the same direction but of different wavelength. Then the vectors  $\overrightarrow{OP}_1$  and  $\overrightarrow{OP}_2$  are in the same direction and so also are  $\overrightarrow{OM}$  and  $\overrightarrow{MP}_2$ . Thus the envelope has the same direction as the carrier wave; the crests are infinitely long.

Again, suppose that the two waves are of equal length but in different directions. Then the vectors  $\overrightarrow{OP}_1$  and  $\overrightarrow{OP}_2$  are of equal length but different direction. The carrier wave, represented by  $\overrightarrow{OM}$ , lies in the mean direction, but the direction of the envelope is now at right angles to the carrier wave. The result is a short-crested system of waves.

In the general case (figure 1 b) it will be seen that the wave crests are staggered, or form an echelon pattern one behind the other. The direction of this pattern is perpendicular to  $P_1P_2$ . The wave-number perpendicular to  $P_1P_2$  is given by the length OT of the perpendicular from O to  $P_1P_2$ ; it is the direction in which the wave-numbers of the two component waves are equal.

The angle  $\beta$  between OM and  $P_1P_2$  is a measure of the skewness of the waves.

The envelope  $(1\cdot1\cdot9)$  and the carrier  $(1\cdot1\cdot10)$  are not necessarily 'free' travelling waves, that is, they do not satisfy equations of the form of  $(1\cdot1\cdot4)$ . Their representation in the wavenumber diagram is valid only so far as the spatial periodicity is concerned. However, for a narrow spectrum  $\overline{u}$ ,  $\overline{v}$ ,  $\overline{\sigma}$  are nearly equal to  $u_1$ ,  $v_1$ ,  $\sigma_1$  respectively, and so the carrier wave does move with nearly the free-wave velocity, if the component waves are themselves free waves. But the envelope moves with a velocity whose components are

$$-\left(\frac{\sigma'}{u'},\frac{\sigma'}{v'}\right) = -\left(\frac{\sigma_1 - \sigma_2}{u_1 - u_2},\frac{\sigma_1 - \sigma_2}{v_1 - v_2}\right).$$
(1.1.11)

To a first approximation this is

$$-\left(\frac{\partial\sigma}{\partial u},\frac{\partial\sigma}{\partial v}\right) = -\left(\frac{\partial w}{\partial u},\frac{\partial w}{\partial v}\right)\frac{\mathrm{d}\sigma}{\mathrm{d}w} = -\left(\cos\theta,\sin\theta\right)\frac{\mathrm{d}\sigma}{\mathrm{d}w},\tag{1.1.12}$$

which is the so-called group velocity. Thus in this special case the envelope moves with the group velocity of the carrier wave.

### 1.2. The representation of a surface having a continuous spectrum

We now assume that the surface possesses a continuous noise spectrum in two dimensions. Generalizing the representation used in equation (1) we write

$$\zeta(x,y,t) = \sum_{n} c_n \cos(u_n x + v_n y + \sigma_n t + \epsilon_n), \qquad (1\cdot 2\cdot 1)$$

where it is supposed that the wave-numbers  $(u_n, v_n)$  are densely distributed throughout the u, v plane, i.e. there are an infinite number in any elementary area du dv.  $\sigma_n$  is a function of  $(u_n, v_n)$ :  $\sigma_n = \sigma(u_n, v_n)$ :

$$\sigma_n = \sigma(u_n, v_n); \qquad (1 \cdot 2 \cdot 2)$$

the amplitudes  $c_n$  are random variables such that in any element dudv we may assume

$$\sum_{n} \frac{1}{2} c_n^2 = E(u, v) \operatorname{d} u \operatorname{d} v; \qquad (1 \cdot 2 \cdot 3)$$

the phases  $\epsilon_n$  are distributed randomly and with equal probability in the interval  $(0, 2\pi)$ . The function E(u, v) will be called the energy spectrum of  $\zeta$ ; the mean-square value of  $\zeta$  per unit area of the sea surface per unit time\* is given by

$$\lim_{X, Y, T \to \infty} \frac{1}{8XYT} \int_{-X}^{X} \int_{-Y}^{Y} \int_{-T}^{T} \zeta^2 dx dy dt = \sum_n \frac{1}{2} c_n^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) du dv.$$
(1.2.4)

Thus the contribution to the mean energy from an element du dv is proportional to E du dv. We shall write

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) \, \mathrm{d}u \, \mathrm{d}v = m_{00}, \qquad (1.2.5)$$

\* It is assumed that average values taken with respect to x, y or t are equivalent to average values with respect to the phases  $\epsilon_n$ .

and in general for the (p, q)th moment of E(u, v) about the origin we write

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \, u^p v^q \, \mathrm{d}u \, \mathrm{d}v = m_{pq}. \tag{1.2.6}$$

These quantities will occur repeatedly throughout the following analysis. It is assumed that they exist up to all orders required.

The function E(u, v) is closely related to the correlation function  $\psi(x, y, t)$  defined by

$$\psi(x,y,t) = \lim_{X,Y,T\to\infty} \frac{1}{8XYT} \int_{-X}^{X} \int_{-Y}^{Y} \int_{-T}^{T} \zeta(x',y',t') \,\zeta(x'+x,y'+y,t'+t) \,\mathrm{d}x' \,\mathrm{d}y' \,\mathrm{d}t'. \quad (1\cdot 2\cdot 7)$$

On substituting from  $(1 \cdot 2 \cdot 1)$  in the above we find

$$\psi(x, y, t) = \sum_{n} \frac{1}{2} c_n^2 \cos(u_n x + v_n y + \sigma_n t), \qquad (1.2.8)$$

which can be written

$$\psi(x,y,t) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \cos(ux + vy + \sigma t) \, \mathrm{d}u \, \mathrm{d}v, \qquad (1\cdot 2\cdot 9)$$

so that  $\psi$  is the cosine transform of E. The even moments  $m_{pq}$  are related very simply to the derivatives of  $\psi$  at the origin:

$$m_{pq} = (-1)^r \frac{\partial^{2r} \psi(0,0,0)}{\partial x^p \partial y^q} \quad (p+q=2r).$$
(1.2.10)

### $1 \cdot 3$ . Conditions for degeneracy

Some important features of the surface can be described immediately in terms of the moments. For example, to find a condition that the wave energy shall all travel in one direction, so that the spectrum is effectively one-dimensional, consider the integral

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u_1, v_1) E(u_2, v_2) (u_1 v_2 - u_2 v_1)^2 du_1 dv_1 du_2 dv_2.$$
(1·3·1)

If the spectrum is one-dimensional, the product  $E(u_1, v_1) E(u_2, v_2)$  is zero everywhere except when  $u_1/v_1 = u_2/v_2$ , when the squared factor vanishes. Therefore the integral vanishes. Conversely, if the spectrum is not one-dimensional there will be a contribution to the integral from some pairs of elements  $du_1 dv_1$ ,  $du_2 dv_2$  for which  $u_1/v_1 \pm u_2/v_2$ , and since the integrand is never negative the integral does not vanish. But on expanding the squared factor and separating the integrations with respect to  $u_1, v_1$  and  $u_2, v_2$  we find that (1·3·1) is equal to

$$2(m_{20}m_{02}-m_{11}^2)=2 \begin{vmatrix} m_{20} & m_{11} \\ m_{11} & m_{02} \end{vmatrix} = 2\Delta_2, \qquad (1\cdot 3\cdot 2)$$

say. Thus a necessary and sufficient condition for E to degenerate into a single one-dimensional spectrum is that  $\Delta_{1} = 0$  (1.3.3)

$$\Delta_2 = 0. \tag{1.3.3}$$

By similar reasoning, a condition for E to degenerate into two one-dimensional spectra (see figure 2a) is that

$$\int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} E(u_1, v_1) E(u_2, v_2) E(u_3, v_3) \\ \times (u_2 v_3 - u_3 v_2)^2 (u_3 v_1 - u_1 v_3)^2 (u_1 v_2 - u_2 v_1)^2 du_1 dv_1 du_2 dv_2 du_3 dv_3 \quad (1\cdot3\cdot4)$$

shall vanish. The squared product may be written



FIGURE 2. The form of the energy spectrum for (a) two intersecting long-crested systems of waves, (b) a system of standing waves, (c) a narrow band of waves, uniform in wavelength and direction and (d) two narrow bands of waves.

(where  $\epsilon_{ijk} = \pm 1$  according as (i, j, k) is an even or odd permutation of (1, 2, 3), and so the integral equals

$$6 \begin{vmatrix} m_{40} & m_{31} & m_{22} \\ m_{31} & m_{22} & m_{13} \\ m_{22} & m_{13} & m_{04} \end{vmatrix} = 6\Delta_4, \qquad (1\cdot 3\cdot 6)$$

say. Thus E degenerates into not more than two one-dimensional spectra if and only if

$$\Delta_4 = 0. \tag{1.3.7}$$

There is an obvious generalization to any number of one-dimensional spectra: the condition that E degenerate into not more than n one-dimensional spectra is that

$$\Delta_{2n} \equiv \begin{vmatrix} m_{2n,0} & m_{2n-1,1} & \dots & m_{n,n} \\ m_{2n-1,1} & m_{2n-2,2} & \dots & m_{n-1,n+1} \\ \vdots & \vdots & & \vdots \\ m_{nn} & m_{n-1,n+1} & \dots & m_{0,2n} \end{vmatrix} = 0.$$
(1·3·8)

Vol. 249. A.

(In practice  $\Delta_2$ ,  $\Delta_4$ , etc., must be compared with quantities of the same dimensions. Thus  $\Delta_{2n}$  may be compared with  $(m_{20}+m_{02})^n$ .)

A condition for E to degenerate into a 'ring' spectrum, such that all the energy corresponds to wave components of the same length but possibly different directions, is that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u_1, v_1) E(u_2, v_2) \left[ (u_1^2 + v_1^2) - (u_2^2 + v_2^2) \right]^2 \mathrm{d}u_1 \mathrm{d}v_1 \mathrm{d}u_2 \mathrm{d}v_2 \qquad (1 \cdot 3 \cdot 9)$$

shall vanish. This integral equals

$$2[(m_{40}+2m_{22}+m_{04}) m_{00}-(m_{20}+m_{02})^2], \qquad (1\cdot 3\cdot 10)$$

and so we must have

$$(m_{40}+2m_{22}+m_{04}) m_{00}-(m_{20}+m_{02})^2=0. \tag{1.3.11}$$

The condition for the energy to be situated at two diametrically opposite points of the spectrum (giving a standing-wave pattern) is that  $(1\cdot3\cdot3)$  and  $(1\cdot3\cdot11)$  shall be satisfied simultaneously (see figure 2b).

A condition for the energy to be concentrated about a single point in the spectrum is that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u_1, v_1) E(u_2, v_2) \left[ (u_1 - u_2)^2 + (v_1 - v_2)^2 \right] \mathrm{d}u_1 \mathrm{d}v_1 \mathrm{d}u_2 \mathrm{d}v_2 \qquad (1 \cdot 3 \cdot 12)$$

shall vanish. This is equivalent to the pair of conditions that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u_1, v_1) E(u_2, v_2) (u_1 - u_2)^2 du_1 dv_1 du_2 dv_2$$
(1.3.13)

shall vanish, and a similar integral with factor  $(v_1 - v_2)^2$ . These are the conditions that the energy be concentrated on lines parallel to the v axis and the u axis respectively (see figure 2c) On expanding the integrals we have

$$\begin{vmatrix} m_{20} & m_{10} \\ m_{10} & m_{00} \end{vmatrix} = 0, \qquad \begin{vmatrix} m_{02} & m_{01} \\ m_{01} & m_{00} \end{vmatrix} = 0.$$
(1·3·14)

A condition for the energy to be concentrated about not more than two points in the spectrum (not necessarily opposite) is that

$$\int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} E(u_1, v_1) E(u_2, v_2) E(u_3, v_3) \prod_{i \neq j} \left[ (u_i - u_j)^2 + (v_i - v_j)^2 \right] \mathrm{d}u_1 \mathrm{d}v_1 \mathrm{d}u_2 \mathrm{d}v_2 \mathrm{d}u_3 \mathrm{d}v_3 \quad (1 \cdot 3 \cdot 15)$$

shall vanish. The term under the product sign may be written

$$\begin{aligned} &(u_2 - u_3)^2 \,(u_3 - u_1)^2 \,(u_1 - u_2)^2 + (v_2 - v_3)^2 \,(v_3 - v_1)^2 \,(v_1 - v_2)^2 \\ &+ \Sigma (u_2 - u_3)^2 \,(u_3 - u_1)^2 \,(v_1 - v_2)^2 + \Sigma (u_2 - u_3)^2 \,(v_3 - v_1)^2 \,(v_1 - v_2)^2. \end{aligned} \tag{1.3.16}$$

Since all the terms are non-negative, each separately must vanish. The first two give the conditions

$$egin{array}{cccc} m_{40} & m_{30} & m_{20} \ m_{30} & m_{20} & m_{10} \ m_{20} & m_{10} & m_{00} \end{array} = 0, \qquad egin{array}{cccc} m_{04} & m_{03} & m_{02} \ m_{03} & m_{02} & m_{01} \ m_{02} & m_{01} & m_{00} \end{array} = 0, \qquad (1\cdot3\cdot17)$$

which are the conditions that the energy shall be at the intersections of two pairs of lines parallel to the v and u axes, i.e. at the corners of a rectangle (figure 2d). The remaining

conditions can also be expressed in terms of the moments. Thus the group of terms under the first summation sign in  $(1 \cdot 3 \cdot 16)$  leads to the condition

$$4 \begin{vmatrix} m_{30} & m_{20} & m_{11} \\ m_{21} & m_{11} & m_{02} \\ m_{20} & m_{10} & m_{01} \end{vmatrix} + 2 \begin{vmatrix} m_{30} & m_{20} & m_{22} \\ m_{20} & m_{10} & m_{12} \\ m_{10} & m_{00} & m_{02} \end{vmatrix} + \begin{vmatrix} m_{40} & m_{20} & m_{21} \\ m_{20} & m_{00} & m_{01} \\ m_{21} & m_{01} & m_{02} \end{vmatrix} = 0.$$
(1.3.18)

The last group of terms in  $(1\cdot 3\cdot 16)$  leads to a similar condition, the pair of suffixes in each of the moments  $m_{bq}$  being interchanged.

We have incidentally shown that each of the combinations of moments on the left-hand sides of equations  $(1\cdot 3\cdot 2)$ ,  $(1\cdot 3\cdot 6)$ ,  $(1\cdot 3\cdot 11)$ ,  $(1\cdot 3\cdot 14)$ ,  $(1\cdot 3\cdot 17)$  and  $(1\cdot 3\cdot 18)$  is never negative.

### 1.4. The spectrum of the surface in an arbitrary direction

Let us consider the curve in which the surface  $\zeta$  is intersected by a perpendicular plane in direction  $\theta$ , that is, the plane  $x \sin \theta - y \cos \theta = 0$ . The curve will represent a one-dimensional random function, whose spectrum  $E_{\theta}$  with regard to the wave-number u' in this direction bears a simple relation to the original spectrum E(u, v). We may call  $E_{\theta}(u')$  the spectrum of the surface in the direction  $\theta$ .

First, let x', y' denote co-ordinates in the x, y plane in directions parallel and perpendicular to the direction  $\theta$ :

$$x' = x\cos\theta + y\sin\theta, \quad y' = -x\sin\theta + y\cos\theta.$$
 (1.4.1)

Reciprocally, x, y are given in terms of x', y' by similar relations, but with the sign of  $\theta$ reversed. On substituting in  $(1 \cdot 2 \cdot 1)$  we have

$$\zeta = \sum_{n} c_n \cos\left(u'_n x' + v'_n y' + \sigma'_n t + \epsilon_n\right), \qquad (1 \cdot 4 \cdot 2)$$

where

where 
$$u'_n = u_n \cos \theta + v_n \sin \theta$$
,  $v'_n = -u_n \sin \theta + v_n \cos \theta$ , (1.4.3)  
that is, the new wave-number  $u'_n$  is the co-ordinate, in the direction  $\theta$ , of the point  $(u_n, v_n)$ ,  
and  $v'_n$  is the co-ordinate at right angles. We have also

$$\sigma'_n = \sigma\{\sqrt{(u_n^2 + v_n^2)}\} = \sigma\{\sqrt{(u_n'^2 + v_n'^2)}\}.$$
 (1.4.4)

On the curve of intersection we have y' = 0 and so

$$\zeta = \sum_{n} c_n \cos\left(u'_n x' + \sigma'_n t + \epsilon_n\right). \tag{1.4.5}$$

The spectrum  $E_{\theta}(u')$  of this curve is defined as the function such that the energy corresponding to any small interval (u', u' + du') is  $E_{\theta}(u') du'$ . Thus if  $\sum_{i=1}^{n} denotes$  summation over the strip (u', u' + du'),

$$E_{\theta}(u') \, \mathrm{d}u = \sum_{\mathrm{d}u', \, v'} \frac{1}{2} c_n^2 = \mathrm{d}u' \int_{-\infty}^{\infty} E(u, v) \, \mathrm{d}v', \qquad (1\cdot 4\cdot 6)$$

and therefore

$$E_{\theta}(u') = \int_{-\infty}^{\infty} E(u, v) \, \mathrm{d}v'. \qquad (1\cdot 4\cdot 7)$$

In other words, if we take a section of the surface in any direction  $\theta$ , the spectrum  $E_{\theta}(u')$ of this section is found by integrating E(u, v) along the line through  $P = (u' \cos \theta, u' \sin \theta)$ at right angles to  $\overrightarrow{OP}$ .

41-2

 $(1 \cdot 4 \cdot 3)$ 

# M. S. LONGUET-HIGGINS ON THE

From equation  $(1\cdot 4\cdot 7)$  there follow some simple and fundamental relations between the moments of the spectrum  $E_{\theta}(u')$  and the moments of the original distribution E(u, v). Let the *n*th moment of  $E_{\theta}$  about the origin be denoted by  $m_n(\theta)$ . Then we have

 $u' = u \cos \theta + v \sin \theta, \quad v' = -u \sin \theta + v \cos \theta$ 

$$m_n(\theta) = \int_{-\infty}^{\infty} E_{\theta}(u') \, u'^n \mathrm{d}u' = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) \, u'^n \mathrm{d}u' \, \mathrm{d}v'. \tag{1.4.8}$$

Since

and

$$rac{\partial(u',v')}{\partial(u,v)}=1, \hspace{1.5cm} (1{\cdot}4{\cdot}10)$$

 $(1 \cdot 4 \cdot 9)$ 

 $(1 \cdot 4 \cdot 17)$ 

we have

$$m_n(\theta) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) \, (u \cos \theta + v \sin \theta)^n \, \mathrm{d}u \, \mathrm{d}v. \tag{1.4.11}$$

After expanding the binomial and integrating each term we find

$$m_n(\theta) = m_{n,0} \cos^n \theta + {n \choose 1} m_{n-1,1} \cos^{n-1} \theta \sin \theta + \dots + m_{0,n} \sin^n \theta, \qquad (1 \cdot 4 \cdot 12)$$

where  $m_{pq}$  is the (p, q)th moment of E about the origin (equation  $(1 \cdot 2 \cdot 6)$ ) and  $\binom{n}{r}$  denotes the binomial coefficient.

In particular we have 
$$m_0(\theta) = m_{00},$$
 (1.4.13)

showing that the r.m.s. value of  $\zeta(x')$  is independent of the direction  $\theta$  and equals the r.m.s. value of  $\zeta(x, y)$ . Next,  $m_1(\theta) = m_{10} \cos \theta + m_{01} \sin \theta$ . (1.4.14)

If  $(\bar{u}, \bar{v})$  denotes the centroid of the two-dimensional spectrum:

$$(ar{u},ar{v}) = \left(\!rac{m_{10}}{m_{00}},rac{m_{01}}{m_{00}}\!
ight), \hspace{1.5cm} (1\!\cdot\!4\!\cdot\!15)$$

and if  $\overline{u}'$  denotes the mean wave-number of the spectrum of  $\zeta(x')$  we have

$$\bar{u}' = \frac{m_1(\theta)}{m_0(\theta)} = \bar{u}\cos\theta + \bar{v}\sin\theta, \qquad (1\cdot 4\cdot 16)$$

which can be expressed as

$$(\overline{u},\overline{v}) = (\overline{w}\cos\overline{\theta},\overline{w}\sin\overline{\theta}).$$
 (1.4.18)

where

. —

$$\overline{w}$$
 and  $\theta$  may be called the mean wave-number and mean direction of the two-dimensional spectrum. Thus the mean wave-number of  $E_{\theta}(u')$  is the projection of the mean wave-number of  $E(u, v)$  on to the line of the section. The physical significance of this result will become clearer in §1.5.

 $\overline{u}' = \overline{w}\cos{(\theta - \overline{\theta})},$ 

The second moment  $m_2(\theta)$  is particularly important. From  $(1\cdot 4\cdot 12)$  we have

$$m_2(\theta) = m_{20}\cos^2\theta + 2m_{11}\cos\theta\sin\theta + m_{02}\sin^2\theta.$$
(1.4.19)

The maxima and minima of this expression are given by

$$m_{2\max}, m_{2\min} = \frac{1}{2} [(m_{20} + m_{02}) \pm \sqrt{\{(m_{20} - m_{02})^2 + 4m_{11}^2\}}], \qquad (1 \cdot 4 \cdot 20)$$

and these occur always in two directions at right angles, given by

$$\tan 2\theta_p = \frac{2m_{11}}{m_{20} - m_{02}}.$$
 (1.4.21)

If  $\theta_{b}$  corresponds to the maximum we have

$$m_2(\theta) = m_{2\max} \cos^2{(\theta - \theta_p)} + m_{2\min} \sin^2{(\theta - \theta_p)}.$$
(1.4.22)

The direction  $\theta_p$  corresponding to the maximum will be called the *principal direction* of the waves. Now  $(m_{\ell}(\theta))^{\frac{1}{2}}$ 

$$\left(\frac{m_2(\theta)}{m_{00}}\right)^{\frac{1}{2}} \tag{1.4.23}$$

is the r.m.s. wave-number in the direction  $\theta$ . For a long-crested system of waves the r.m.s. wave-number is a maximum perpendicular to the crests and a minimum parallel to the crests. In general, therefore, a convenient measure of the *long-crestedness* is given by the ratio

$$\left(\frac{m_{2\max.}}{m_{2\min.}}\right)^{\frac{1}{2}},\tag{1.4.24}$$

which we denote by  $1/\gamma$ . Thus we have

$$\gamma^{2} = \frac{m_{2\min.}}{m_{2\max.}} = \frac{(m_{20} + m_{02}) - \sqrt{\{(m_{20} - m_{02})^{2} + 4m_{11}^{2}\}}}{(m_{20} + m_{02}) + \sqrt{\{(m_{20} - m_{02})^{2} + 4m_{11}^{2}\}}}.$$
 (1.4.25)

When the condition  $(1\cdot 3\cdot 3)$  for a one-dimensional spectrum is satisfied we have

$$\gamma = 0, \quad 1/\gamma = \infty.$$
 (1.4.26)

The two quantities  $m_{2\text{max.}}$ ,  $m_{2\text{min.}}$  are clearly invariant under a rotation of the axes. Hence we have also the invariants

$$m_{2\max} + m_{2\min} = m_{20} + m_{02} = m, \qquad (1.4.27)$$

say, and 
$$m_{2\max}, m_{2\min} = m_{20}m_{02} - m_{11}^2 = \Delta_2.$$
 (1.4.28)

### 1.5. The wave envelope

By analogy with §1·1 we define the mean wave-number as the centroid of the energy distribution:

$$m_{00}\overline{u} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) \, u \, \mathrm{d}u \, \mathrm{d}v = m_{10},$$

$$m_{00}\overline{v} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) \, v \, \mathrm{d}u \, \mathrm{d}v = m_{01}$$

$$(1.5.1)$$

and we define also the mean frequency  $\overline{\sigma}/2\pi$  by the analogous equation

$$m_{00}\overline{\sigma} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \,\sigma \,\mathrm{d}u \,\mathrm{d}v = m'_{00}, \qquad (1\cdot 5\cdot 2)$$

say. Now let  $(1 \cdot 2 \cdot 1)$  be written in the form

$$\zeta = \mathscr{R} \sum_{n} c_n \exp\left\{i(u_n x + v_n y + \sigma_n t + \epsilon_n)\right\}$$
(1.5.3)

$$= \mathscr{R}\left[\sum_{n} c_{n} \exp\left\{i\left[\left(u_{n}-\overline{u}\right) x+\left(v_{n}-\overline{v}\right) y+\left(\sigma_{n}-\overline{\sigma}\right) t+\epsilon_{n}\right]\right\} \exp\left\{i\left(\overline{u}x+\overline{v}y+\overline{\sigma}t\right)\right\}\right], \quad (1\cdot5\cdot4)$$

where  $\mathscr{R}$  denotes the real part. This expresses  $\zeta$  as the product of a carrier wave

$$\exp\{i(\bar{u}x+\bar{v}y+\bar{\sigma}t)\},\qquad(1.5.5)$$

and a slowly varying amplitude function

$$\rho e^{i\phi} = \sum_{n} c_n \exp\{i[(u_n - \overline{u}) x + (v_n - \overline{v}) y + (\sigma_n - \overline{\sigma}) t + \epsilon_n]\}, \quad (1.5.6)$$

M. S. LONGUET-HIGGINS ON THE

which may be called the complex envelope. ( $\rho$  and  $\phi$  are real functions of (x, y, t), with  $\rho \ge 0$ .) Any other choice for the frequency of the carrier wave might have been taken; the mean wave-number has the unique property that the secular increase of  $\phi$  with x and y is zero (as will be shown in §2.8).

Comparing  $(1\cdot 5\cdot 3)$  and  $(1\cdot 5\cdot 4)$  we see that *the real part* of the amplitude function, i.e.  $\rho \cos \phi$ , has the same spectrum as  $\zeta$ , only with the origin moved to  $(\bar{u}, \bar{v})$ ; similarly for the imaginary part. ( $\rho$  itself, however, is a different type of function, being essentially positive.) The properties of the envelope, therefore, are defined by the moments of the energy distribution about the mean. Let

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \ (u-\overline{u})^p \ (v-\overline{v})^q \ \mathrm{d}u \ \mathrm{d}v = \mu_{pq}. \tag{1.5.7}$$

It is easily seen that

$$\mu_{00} = m_{00}, \quad \mu_{10} = \mu_{01} = 0,$$
 (1.5.8)

and

$$\begin{array}{l} \mu_{20} = m_{20} - \overline{u}^2 m_{00} = (m_{20} m_{00} - m_{10}^2) / m_{00}, \\ \mu_{11} = m_{11} - \overline{u} \overline{v} m_{00} = (m_{11} m_{00} - m_{10} m_{01}) / m_{00}, \\ \mu_{02} = m_{02} - \overline{v}^2 m_{00} = (m_{02} m_{00} - m_{01}^2) / m_{00}. \end{array} \right\}$$

$$(1 \cdot 5 \cdot 9)$$

The second moment about the mean in a direction  $\theta$  is

$$\mu_2(\theta) = \mu_{20}\cos^2\theta + 2\mu_{11}\cos\theta\sin\theta + \mu_{02}\sin^2\theta, \qquad (1.5.10)$$

and the principal direction of the envelope is given by

$$\tan 2\theta_e = \frac{2\mu_{11}}{\mu_{20} - \mu_{02}}.$$
 (1.5.11)

The angle  $\beta$  between the principal direction of the envelope and the principal direction of the waves is given by

$$\tan 2\beta = \tan 2(\theta_e - \theta_p) = \frac{2\mu_{11}(m_{20} - m_{02}) - 2m_{11}(\mu_{20} - \mu_{02})}{(\mu_{20} - \mu_{02})(m_{20} - m_{02}) + 4\mu_{11}m_{11}}.$$
 (1.5.12)

Thus  $\beta$  is a convenient measure of the *skewness* of the waves (see § 1.1).

Consider again the curve of intersection of the surface with a vertical plane in direction  $\theta$ . We may see that the envelope of this curve is simply the intersection of the two-dimensional envelope with the vertical plane. For on the one hand we have from (1.5.6)

$$\rho e^{i\phi} = \sum_{n} c_n \exp\{i[(u'_n - \overline{u}') x' + (v'_n - \overline{v}) y' + (\sigma'_n - \overline{\sigma}) t + \epsilon_n]\}, \quad (1.5.13)$$

where  $u'_n$  and  $v'_n$  are given by  $(1 \cdot 4 \cdot 3)$  and

$$\bar{u}' = \bar{u}\cos\theta + \bar{v}\sin\theta, \quad \bar{v}' = -\bar{u}\sin\theta + \bar{v}\cos\theta. \tag{1.5.14}$$

The intersection of the envelope by the plane y' = 0 is therefore given by

$$\rho e^{i\phi} = \sum_{n} c_n \exp\{i[(u'_n - \overline{u}') x' + (\sigma'_n - \overline{\sigma}) t + \epsilon_n]\}.$$
(1.5.15)

On the other hand from (1.4.5) we may write

$$\zeta(x',t) = \mathscr{R}\left[\sum_{n} c_n \exp\left\{i\left[\left(u'_n - \overline{u}\right)x' + \left(\sigma'_n - \overline{\sigma}\right)t + \epsilon_n\right]\right\}\exp\left\{i\left(\overline{u}'x' + \overline{\sigma}t\right)\right\}\right], \quad (1\cdot5\cdot16)$$

where  $\overline{u}'$  is given by (1.5.14). But we saw in §1.4 that  $\overline{u}'$  is also the mean wave-number for the function  $\zeta(x',t)$  and therefore exp $\{i(\overline{u}'x'+\overline{\sigma}t)\}\$  is, by definition, the carrier wave for  $\zeta(x',t)$  and  $(1\cdot 5\cdot 15)$  is the envelope; which proves the result.

#### 1.6. A narrow spectrum

A case of special interest is when the energy is concentrated near a single point in the spectrum, so that the component waves are nearly constant in wavelength and direction. As we saw earlier, the conditions satisfied by the first-order and second-order moments are that the left-hand sides of equations  $(1\cdot 3\cdot 14)$  are small. In terms of the moments this implies  $\mu_{20} + \mu_{02} \ll m_{20} + m_{02}$ , or equivalently

$$\mu_{20} + \mu_{02} \ll (\overline{u}^2 + \overline{v}^2) \, m_{00}. \tag{1.6.1}$$

The envelope of the waves, as defined in the previous section, then has some special properties. If in (1.5.2) we expand  $\sigma(u, v)$  in a Taylor series about  $(\bar{u}, \bar{v})$  we have

$$\begin{split} m_{00}\overline{\sigma} &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \left[ \sigma(\overline{u},\overline{v}) + (u-\overline{u}) \frac{\partial}{\partial \overline{u}} \sigma(\overline{u},\overline{v}) + (v-\overline{v}) \frac{\partial}{\partial \overline{v}} \sigma(\overline{u},\overline{v}) \right] \mathrm{d}u \, \mathrm{d}v \\ &= m_{00} \sigma(\overline{u},\overline{v}) + \mu_{10} \frac{\partial}{\partial \overline{u}} \sigma(\overline{u},\overline{v}) + \mu_{01} \frac{\partial}{\partial \overline{v}} \sigma(\overline{u},\overline{v}), \end{split}$$
(1.6.2)

terms of higher order being negligible. Since  $\mu_{10} = \mu_{01} = 0$  we have

$$\overline{\sigma} = \sigma(\overline{u}, \overline{v}). \tag{1.6.3}$$

In other words, the carrier wave is a free wave with the frequency and velocity appropriate to its wave-number. Further, in (1.5.6) we may write

$$\sigma_n - \overline{\sigma} = (u_n - \overline{u}) \frac{\partial \overline{\sigma}}{\partial \overline{u}} + (v_n - \overline{v}) \frac{\partial \overline{\sigma}}{\partial \overline{v}}, \qquad (1 \cdot 6 \cdot 4)$$

so that

that 
$$\rho e^{i\phi} = \sum_{n} \exp\{i[(u_n - \overline{u})(x + t\partial\overline{\sigma}/\partial\overline{u}) + (v_n - \overline{v})(y + t\partial\overline{\sigma}/\partial\overline{v})]\},$$
 (1.6.5)

which is a function of  $(x + t\partial \overline{\sigma}/\partial \overline{u})$  and  $(y + t\partial \overline{\sigma}/\partial \overline{v})$  only. In other words, the envelope moves bodily with velocity

$$\left(-\frac{\partial \overline{\sigma}}{\partial \overline{u}}, -\frac{\partial \overline{\sigma}}{\partial \overline{v}}\right). \tag{1.6.6}$$

 $\sigma$  being a function of  $w = (u^2 + v^2)^{\frac{1}{2}}$  only, this velocity is

$$-\left(\frac{\partial \overline{w}}{\partial \overline{u}}, \frac{\partial \overline{w}}{\partial \overline{v}}\right) \frac{\mathrm{d}\overline{\sigma}}{\mathrm{d}\overline{w}} = -\left(\cos\overline{\theta}, \sin\overline{\theta}\right) \frac{\mathrm{d}\overline{\sigma}}{\mathrm{d}\overline{w}},\tag{1.6.7}$$

which is the group velocity of the carrier wave.

Let axes be chosen so that the u axis passes through the centroid  $(\bar{u}, \bar{v})$ , making  $\bar{v} = 0$ . On expanding  $u^p$ , = { $\overline{u} + (u - \overline{u})^p$ }, by the binomial theorem we have

$$\begin{split} m_{pq} &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \, u^{p} v^{q} \, \mathrm{d}u \, \mathrm{d}v \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \left[ \overline{u}^{p} + p \overline{u}^{p-1} (u - \overline{u}) + \ldots + (u - \overline{u})^{p} \right] (v - \overline{v})^{q} \, \mathrm{d}u \, \mathrm{d}v \\ &= \overline{u}^{p} \mu_{00} + p \overline{u}^{p-1} \mu_{1q} + \ldots + \mu_{pq}. \end{split}$$
(1.6.8)

In particular, since  $\mu_{10} = \mu_{01} = 0$ , we have

$$m_{20} = \overline{u}^2 \mu_{00} + \mu_{20}, \quad m_{11} = \mu_{11}, \quad m_{02} = \mu_{02}.$$
 (1.6.9)

Thus (1.4.9) becomes

$$m_2(\theta) = \bar{u}^2 \mu_{00} \cos^2 \theta + (\mu_{20} \cos^2 \theta + 2\mu_{11} \cos \theta \sin \theta + \mu_{02} \sin^2 \theta).$$
 (1.6.10)

Since  $\mu_{11} \leq (\mu_{20}\mu_{02})^{\frac{1}{2}}$  it follows that all three moments  $\mu_{20}$ ,  $\mu_{11}$ ,  $\mu_{02}$  are small compared with  $\overline{u}^2\mu_{00}$ . Hence  $m_2(\theta)$  has a maximum near  $\theta = 0, \pi$  and a minimum near  $\theta = \pm \frac{1}{2}\pi$ . In other words, the principal direction lies along the axis of u. The long-crestedness  $\gamma^{-1}$  was defined as the ratio of the r.m.s. wave-numbers parallel and perpendicular to the principal directions. Thus]

$$\gamma^2 = \frac{m_2(\frac{1}{2}\pi)}{m_2(0)} = \frac{\mu_{02}}{\overline{u}^2 \mu_{00}}.$$
 (1.6.11)

Now in the neighbourhood of the centroid we have  $v = \bar{u}\theta$  very nearly, so that

$$\mu_{02} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) \,\overline{u}^2 \theta^2 \,\mathrm{d}u \,\mathrm{d}v. \tag{1.6.12}$$

336

$$\gamma^2 \mu_{00} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) \,\theta^2 \,\mathrm{d}u \,\mathrm{d}v. \tag{1.6.13}$$

In other words,  $\gamma$  is the r.m.s. angular deviation of the energy from the mean direction.

Since the principal direction of the waves coincides with the u axis, the angle of skewness  $\beta$  is the angle between the u axis and the principal direction of the envelope, that is,

$$\tan 2\beta = \frac{2\mu_{11}}{\mu_{20} - \mu_{02}}.$$
 (1.6.14)

It will be found convenient to introduce one further parameter for a narrow wave spectrum:  $y = (\mu / \overline{\mu}^2 \mu)^{\frac{1}{2}}$  (1.6.15)

$$\nu = (\mu_{20}/\bar{u}^2\mu_{00})^{\frac{1}{2}}.$$
 (1.6.15)

 $\nu$  is proportional to the r.m.s. width of the spectrum in the principal direction. We shall show in §2.8 that  $\nu^{-1}$  is a measure of the average number of waves per 'group'.

### PART II. STATISTICAL PROPERTIES

The fundamental statistical distributions of  $\zeta$  and its derivatives are given in §2·1. The following three sections are devoted to properties of the surface not involving motion, and the next three sections to the distributions of velocities associated with these properties. Lastly, §§2·8 to 2·10 deal with the envelope of the surface and with properties which can be derived from it.

The distributions of the surface elevation  $\zeta$  and of the two components of gradient  $\partial \zeta / \partial x$ ,  $\partial \zeta / \partial y$ are normal in one and two dimensions respectively (equations (2·1·8) and (2·1·12)). The greatest r.m.s. gradient is in the principal direction of the surface. The distribution of the magnitude  $\alpha$  of the gradient regardless of direction is given by (2·1·31) and figure 3. For very short-crested waves the distribution is a Rayleigh distribution; for very long-crested waves it tends to a normal distribution, with an anomaly near  $\alpha = 0$ , the shape of which is shown in figure 4. The probability distribution of the horizontal direction  $\theta$  of the gradient is given by (2·1·37) and figure 5. It is shown that as the long-crestedness increases, the direction of the gradient becomes more and more certain to be near the principal direction.

In §2·2 is found the mean number of zeros of the surface along a horizontal line in an arbitrary direction  $\theta$ . The number  $N_0$  per unit distance is given by (2·2·5). Thus  $N_0$  is a maximum when  $\theta$  is in the principal direction, and a minimum in the direction at right angles. The ratio  $N_{0\text{max}}/N_{0\text{min}}$ .

is equal to the long-crestedness  $\gamma^{-1}$ . The mean number of times that the surface crosses a line at arbitrary level is also found (2·2·12), and the mean number of crests and troughs of a plane section of the surface in any direction.

The average length of a contour of constant height drawn on the surface is derived in §2·3. The length  $\bar{s}$  per unit area is given by (2·3·16). The distribution of the direction  $\theta$  of the normal to a particular contour, at points distributed uniformly along it, is given by (2·3·23). As in §2·1, when the waves become long-crested, the direction becomes concentrated near the principal direction.

Next ( $\S2\cdot4$ ) the average density of maxima and minima of the surface per unit horizontal area is considered. It is shown that the average density of maxima,  $D_{ma.}$ , is equal to one-half of the average density of saddle-points, and to one-quarter of the total density of stationary points on the surface. The actual density is given by  $(2\cdot4\cdot51)$ , in the general case. For a narrow spectrum, the density is given by  $(2\cdot4\cdot61)$  and table 1.  $D_{ma.}$  depends not only on the long-crestedness but also on a parameter *a* representing the *peakedness* of the energy distribution with regard to direction.

Passing now to properties depending on the motion of the surface, we consider in §2.5 the velocity of the zeros of the surface along an arbitrary line. We find that the velocities have a probability distribution given by (2.5.15). This is symmetrical about a mean value depending on the first-order moments. Similarly, the velocities of maxima and minima of a plane section of the surface have a distribution given by (2.5.19). These distributions are studied in the special case of a narrow spectrum. The width of the distribution depends on both the width of the energy spectrum and on the dispersive properties of the medium.

The motion of a contour on the surface can be defined locally by the velocities of its points of intersection with lines parallel to the axes of x, y (§2.6). The distribution of the reciprocals of the velocities, which is simpler than that of the velocities themselves, is given by (2.6.21). The distribution is discussed in detail for the case of a narrow spectrum; the contours of constant probability are then concentric ellipses.

In §2.7 is considered the motion of the 'specular points' of the surface, that is, points where the gradient of the surface takes a certain value. (Such points on the sea surface are, to a distant observer, points of reflected sunlight.) The probability distribution of the two components of velocity is given by (2.7.31). In the special case of a narrow spectrum the mean velocity of the specular points is equal to the phase velocity of the carrier wave. The departures of the velocities from the mean velocity have a distribution given by (2.7.37). This expression has been computed for three different values of the peakedness a, and is shown in figure 12 a, b and c.

In §2.8 we consider some properties of the wave envelope, from which we derive some other useful distributions. The distribution of the envelope function itself is a Rayleigh distribution  $(2\cdot8\cdot6)$ . The joint distribution of  $\rho$ ,  $\partial\rho/\partial x$  and  $\partial\rho/\partial y$  is given by  $(2\cdot8\cdot15)$ , from which it follows that the envelope possesses a number of properties analogous to the original surface. The envelope also controls the 'grouping' of the waves, and we find, taking a section of the surface in an arbitrary direction  $\theta$ , that the average length of a group is 2/N, where N is given by  $(2\cdot8\cdot26)$ . Hence the average length of a group is least in the principal direction and greatest in the direction at right angles. We find the average number of waves in a group  $(2\cdot8\cdot27)$  and the condition that this shall be independent of the direction  $\theta$   $(2\cdot8\cdot28)$ .

When the spectrum is narrow, the crests of the waves lie practically on the envelope, and so we are able to deduce that the probability distribution of the heights of crests is approximately a Rayleigh distribution  $(2\cdot9\cdot1)$ . The distribution of the heights of maxima is found through the distribution of the heights of the maxima of the envelope  $(2\cdot9\cdot8)$ . This distribution is shown in figure 13 for different values of peakedness *a*. The limiting case of two crossing swells (a=1) is given by  $(2\cdot8\cdot12)$  and is also shown in figure 13.

Finally, in §2·10 is deduced the distribution of the intervals l between successive zero-crossings, or between the successive points of intersection of a straight line with a contour at fixed height. The distribution of l for waves of all heights is given by (2·10·18). However, if the waves are classified according to their height, the distribution of l is given by (2·10·23), and hence it is found that l is less scattered for the high waves than for the low waves. The degree of scattering is inversely proportional to the average number of waves in a group.

VOL. 249. A.

## M. S. LONGUET-HIGGINS ON THE

### 2.1. The distribution of surface elevation and gradient

Let  $\xi_1, ..., \xi_n$  be *n* quantities, each the sum of a large number of independent variables whose expectation is zero. Then under certain general conditions (discussed by Rice 1944, 1945; see also Cramér 1937) the joint-probability distribution of  $\xi_1, ..., \xi_n$  is normal in *n* dimensions:

$$p(\xi_1, \dots, \xi_n) = \frac{1}{(2\pi)^{\frac{1}{2}n} \Delta^{\frac{1}{2}}} \exp\left\{-\frac{1}{2} M_{ij} \xi_i \xi_j\right\},$$
(2.1.1)

where  $(M_{ii})$  is the inverse matrix to

$$(\Xi_{ij}) = \begin{pmatrix} \overline{\xi_1^2} & \overline{\xi_1\xi_2} & \dots & \overline{\xi_1\xi_n} \\ \overline{\xi_2\xi_1} & \overline{\xi_2^2} & \dots & \overline{\xi_2\xi_n} \\ \vdots & \vdots & \vdots & \vdots \\ \overline{\xi_n\xi_1} & \overline{\xi_n\xi_2} & \dots & \overline{\xi_n^2} \end{pmatrix}$$
(2.1.2)

and

The elements of  $(\Xi_{ij})$  are the mean products  $\overline{\xi_i \xi_j}$  of the variables  $\xi_i$  and  $\xi_j$  over the probability space of the independent components.  $(\Xi_{ij})$  is a positive-definite matrix, for if  $\alpha_1, ..., \alpha_n$  are any *n* parameters not all zero

 $\Delta = |\Xi_{ii}|.$ 

$$\alpha_i \alpha_j \overline{\xi_i \xi_j} = \overline{(\alpha_i \xi_i)^2} > 0.$$
(2.1.4)

 $(2 \cdot 1 \cdot 3)$ 

Now according to equation  $(1\cdot 2\cdot 1)$ ,  $\zeta$  and also its derivatives are variables of this type. Further, writing for brevity

$$u_n x + v_n y + \sigma_n t + \epsilon_n = \phi_n, \qquad (2.1.5)$$

we have

$$\overline{\zeta^2} = \overline{(\sum_n c_n \cos \phi_n)^2} = \sum_n \frac{1}{2} c_n^2, \qquad (2 \cdot 1 \cdot 6)$$

since the phases are random. Thus

$$\overline{\zeta^2} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u, v) \, \mathrm{d}u \, \mathrm{d}v = m_{00}, \qquad (2 \cdot 1 \cdot 7)$$

and accordingly the probability distribution of  $\xi_1$ , =  $\zeta$ , is

$$p(\xi_1) = \frac{1}{(2\pi)^{\frac{1}{2}} m_{00}^{\frac{1}{2}}} \exp\left\{-\xi_1^2 / 2m_{00}\right\}.$$
(2.1.8)

Similarly

$$\left\{ \frac{\partial \zeta}{\partial x} \right)^2 = \overline{\left( -\sum_n c_n u_n \sin \phi_n \right)^2} = \sum_n \frac{1}{2} c_n^2 u_n^2 = m_{20}, \\
\overline{\left( \frac{\partial \zeta}{\partial y} \right)^2} = \overline{\left( -\sum_n c_n v_n \sin \phi_n \right)^2} = \sum_n \frac{1}{2} c_n^2 v_n^2 = m_{02}, \\$$
(2.1.9)

$$\frac{\overline{\partial\zeta}}{\partial x}\frac{\partial\zeta}{\partial y} = \overline{\left(\sum_{n} c_{n}u_{n}\sin\phi_{n}\right)\left(\sum_{n} c_{n}v_{n}\sin\phi_{n}\right)} = \sum_{n} \frac{1}{2}c_{n}^{2}u_{n}v_{n} = m_{11}.$$

The matrix of correlations for

$$\xi_2, \xi_3 = \frac{\partial \zeta}{\partial x}, \frac{\partial \zeta}{\partial y}$$
(2.1.10)

is therefore

$$\Xi_{ij}) = egin{pmatrix} m_{20} & m_{11} \ m_{11} & m_{02} \end{pmatrix}, \qquad (2\cdot 1\cdot 11)$$

and the joint-probability distribution is

$$p(\xi_2,\xi_3) = \frac{1}{2\pi\Delta_2^{\frac{1}{2}}} \exp\left\{-\left(m_{02}\xi_2^2 - 2m_{11}\xi_2\xi_3 + m_{20}\xi_3^2\right)/2\Delta_2\right\},$$
(2.1.12)

where

$$\Delta_2 = \left| \begin{array}{cc} m_{20} & m_{11} \\ m_{11} & m_{02} \end{array} \right|. \tag{2.1.13}$$

The cross-correlations between  $\zeta$  and  $\partial \zeta / \partial x$ ,  $\partial \zeta / \partial y$  are given by

$$\zeta \frac{\partial \zeta}{\partial x} = \left( \overline{\sum_{n} c_n \cos \phi_n} \right) \left( - \sum_{n} c_n u_n \sin \phi_n \right) = 0, \\ \overline{\zeta \frac{\partial \zeta}{\partial y}} = \overline{\left( \sum_{n} c_n \cos \phi_n \right) \left( - \sum_{n} c_n v_n \sin \phi_n \right)} = 0, \end{cases}$$
(2.1.14)

so that  $\zeta$  and  $\partial \zeta / \partial x$ ,  $\partial \zeta / \partial y$  are uncorrelated. The joint-probability distribution of

$$(\xi_1, \xi_2, \xi_3) = (\zeta, \partial \zeta / \partial x, \partial \zeta / \partial y)$$
(2.1.15)

is therefore given by

$$p(\xi_1,\xi_2,\xi_3) = p(\xi_1) p(\xi_2,\xi_3), \qquad (2.1.16)$$

where  $p(\xi_1)$  is given by (2.1.8) and  $p(\xi_2, \xi_3)$  by (2.1.12).

In general we find, by repeated differentiation,

$$\left(\frac{\partial^{p+q}\zeta}{\partial x^{p}\partial y^{q}}\right)^{2} = m_{2p, 2q} \\
\frac{\partial^{p+q}\zeta}{\partial x^{p}\partial y^{q}}\frac{\partial^{p'+q'}\zeta}{\partial x^{p'}\partial y^{q'}} = (-1)^{\frac{1}{2}(p+q-p'-q')}m_{p+p',q+q'} \quad \text{or} \quad 0, \qquad (2\cdot1\cdot17)$$

and

according as (p+q-p'-q') is even or odd. For example, the derivatives of order p+q=nare not correlated with those of order p' + q' = n + 1; but they are correlated, negatively in general, with those of order n+2.

Slightly different results apply to derivatives involving the time t. We have from  $(1 \cdot 2 \cdot 1)$ 

$$\frac{\partial \zeta}{\partial t} = -\sum_{n} c_n \sigma_n \sin\left(u_n x + v_n y + \sigma_n t + \epsilon_n\right), \qquad (2.1.18)$$

and so the energy spectrum of  $\partial \zeta / \partial t$  is  $\sigma^2 E(u, v)$ . The correlations between the derivatives of  $\zeta$  and those of  $\partial \zeta / \partial t$  are given in terms of the moments

$$m'_{pq} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \sigma E(u, v) \, u^{p} v^{q} \, \mathrm{d}u \, \mathrm{d}v,$$

$$m''_{pq} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \sigma^{2} E(u, v) \, u^{p} v^{q} \, \mathrm{d}u \, \mathrm{d}v$$

$$(2.1.19)$$

of the functions  $\sigma E(u, v)$  and  $\sigma^2 E(u, v)$ . As in (2.1.9) we have

$$\overline{\left(\frac{\partial\zeta}{\partial t}\right)^2} = m_{00}'', \quad \overline{\frac{\partial\zeta}{\partial t}\frac{\partial\zeta}{\partial x}} = m_{10}', \quad \overline{\frac{\partial\zeta}{\partial t}\frac{\partial\zeta}{\partial y}} = m_{01}', \quad (2.1.20)$$

and in general

$$\left(\overline{\frac{\partial^{p+q+1}\zeta}{\partial x^p \partial y^q \partial t}}\right)^2 = m_{2p,2q}''$$
(2.1.21)

and

 $\overline{\left(\frac{\partial^{p+q+1}\zeta}{\partial x^{p}\partial u^{q}\partial t}\right)}\left(\frac{\partial^{p'+q'\zeta}}{\partial x^{p'}\partial u^{q'}}\right)}=(-1)^{\frac{1}{2}(p+q-p'-q'+1)}m'_{p+p',q+q'}\quad\text{or}\quad 0,$  $(2 \cdot 1 \cdot 22)$ according as (p+q-p'-q') is odd or even. Thus all the correlations of the spatial deri-

vatives of  $\partial \zeta / \partial t$  with the spatial derivatives of  $\zeta$  are expressible in terms of the odd moments 42-2

of  $\sigma E$ . The odd derivatives of  $\partial \zeta / \partial t$  are all independent of the odd derivatives of  $\zeta$  but are correlated with all the even derivatives; and vice versa.

Let us now consider more closely the pattern of surface slopes. If the magnitude of the surface slope is  $\alpha$  and its direction is  $\theta$  we have

$$(\xi_2,\xi_3) = \left(\frac{\partial\zeta}{\partial x},\frac{\partial\zeta}{\partial y}\right) = (\alpha\cos\theta,\alpha\sin\theta), \qquad (2\cdot1\cdot23)$$

and so

$$p(\alpha,\theta) = \left| \frac{\partial(\xi_2,\xi_3)}{\partial(\alpha,\theta)} \right| p(\xi_2,\xi_3) = \alpha p(\xi_2,\xi_3), \qquad (2\cdot 1\cdot 24)$$

or from (2.1.12)

$$p(\alpha,\theta) = \frac{\alpha}{2\pi\Delta_2^{\frac{1}{2}}} \exp\left\{-\alpha^2 (m_{02}\cos^2\theta - 2m_{11}\cos\theta\sin\theta + m_{20}\sin^2\theta)/2\Delta_2\right\}.$$
(2.1.25)

If we take the x axis along the principal direction, so that  $m_{11}$  vanishes and  $m_{20} \ge m_{02}$ , then

$$p(\alpha,\theta) = \frac{\alpha^2}{2\pi\Delta_2^{\frac{1}{2}}} \exp\{-\alpha^2 (m_{02}\cos^2\theta + m_{20}\sin^2\theta)/2\Delta_2\}.$$
 (2.1.26)

For a fixed value of  $\theta$ , the r.m.s. slope is given by

$$\left[\frac{\int_{0}^{\infty} \alpha^{2} p(\alpha, \theta) \, \mathrm{d}\alpha}{\int_{0}^{\infty} p(\alpha, \theta) \, \mathrm{d}\alpha}\right]^{\frac{1}{2}} = \left[\frac{2\Delta_{2}}{m_{02}\cos^{2}\theta + m_{20}\sin^{2}\theta}\right]^{\frac{1}{2}}.$$
(2.1.27)

The maximum r.m.s. slope, therefore, is in a direction  $\theta = 0$ , that is to say, in the principal direction. The minimum slope is in the direction at right angles to this.

The statistical distribution of the slope regardless of direction may be found by integrating  $p(\alpha, \theta)$  with respect to  $\theta$  from 0 to  $2\pi$ . We find

$$p(\alpha) = \frac{\alpha}{\Delta_2^{\frac{1}{2}}} \exp\left\{-\alpha^2 (m_{20} + m_{02})/4\Delta_2\right\} I_0[\alpha^2 (m_{20} - m_{02})/4\Delta_2], \qquad (2 \cdot 1 \cdot 28)$$

$$I_0(z) = \frac{1}{2\pi} \int_0^{2\pi} e^{-z\sin\theta} \,\mathrm{d}\theta = J_0(\mathrm{i}z), \qquad (2.1.29)$$

 $I_0$  being the Bessel function of order zero with imaginary argument (see Whittaker & Watson 1952, chap. 17). Writing

$$\eta = \frac{\alpha}{(m_{20} + m_{02})^{\frac{1}{2}}} = \frac{\alpha}{m}$$
(2.1.30)

for the relative slope, and  $\gamma^{-1} = (m_{20}/m_{02})^{\frac{1}{2}}$  for the long-crestedness, we have

$$p(\eta) = \eta(\gamma + \gamma^{-1}) \exp\{-\eta^2(\gamma + \gamma^{-1})^2/4\} I_0[\eta^2(\gamma^{-2} - \gamma^2)/4].$$
 (2.1.31)

This distribution is shown in figure 3, for  $\gamma = 1$ ,  $\frac{1}{2}$ ,  $\frac{1}{4}$  and 0. When  $\gamma = 1$  we have, since  $I_0(0) = 1$ ,  $p(\eta) = 2\eta e^{-\eta^2}$ . (2.1.32)

Thus for short-crested waves the slopes have a Rayleigh distribution. Now as z tends to infinity,  $I_0(z) \sim (2\pi z)^{-\frac{1}{2}} e^z$  (Whittaker & Watson 1952, p. 373) and so when  $\gamma$  is small we have, for general values of  $\eta$ ,  $(2)^{\frac{1}{2}}$ 

$$p(\eta) \sim \left(\frac{2}{\pi}\right)^{\frac{1}{2}} e^{-\frac{1}{2}\eta^2}.$$
 (2.1.33)



FIGURE 3. The probability distribution of the surface slope  $\eta = \alpha/(m_{20}^2 + m_{02}^2)^{\frac{1}{2}}$ , for different values of the long-crestedness.



FIGURE 4. The limiting form of the slope distribution close to the origin, for a very long-crested surface.

In other words, for long-crested waves the slopes have in general a normal distribution (as we should expect, for since the slopes are nearly all in one plane, the distribution of  $\alpha$  is the same as the distribution of  $\partial \zeta / \partial x$ , which is normal). However, for very small slopes, comparable with  $\gamma m$ , we must use the approximation

$$p(\eta) = (\eta/\gamma) e^{-\eta^2/4\gamma^2} I_0(\eta^2/4\gamma^2), = f(\eta/\gamma),$$
 (2.1.34)

say.  $f(\eta/\gamma)$  is plotted in figure 4. As  $\eta/\gamma \to \infty$ , so  $f \to (2/\pi)^{\frac{1}{2}}$ , which is the value of  $(2 \cdot 1 \cdot 33)$  at the origin. The anomalous distribution near the origin appears to arise from directions  $\theta$  which are nearly perpendicular to the principal direction; since the crests are only of finite length, the chance of a very small slope in this direction is less than if the waves were two-dimensional. Nevertheless, the integral of  $(2 \cdot 1 \cdot 34)$  from 0 to  $\infty$  is equal to 1, so that as the waves become infinitely long-crested the contribution to the integrated probability from the anomalous term is vanishingly small.



FIGURE 5. The probability distribution of the direction  $\theta$  of the surface gradient for different values of the long-crestedness.  $\theta = 0$  is the principal direction.

Even when the waves are not long-crested, still for large values of  $\eta$ 

$$p(\eta) \sim \left(\frac{2}{\pi}\right)^{\frac{1}{2}} \frac{1}{(1-\gamma^2)^{\frac{1}{2}}} e^{-\frac{1}{2}\eta^2}.$$
(2.1.35)

Thus for large slopes the distribution always approaches a normal distribution ultimately, provided  $\gamma < 1$ .

The statistical distribution of the direction  $\theta$  of the gradient is found by integrating (2·1·26) with respect to  $\alpha$  from 0 to  $\infty$ :

$$p(\theta) = \frac{\Delta_2^*}{2\pi (m_{02}\cos^2\theta + m_{20}\sin^2\theta)}.$$
 (2.1.36)

$$p(\theta) = \frac{\gamma}{2\pi(\gamma^2 \cos^2\theta + \sin^2\theta)}.$$
 (2.1.37)

or

When  $\gamma = 1$  (the waves are short-crested),  $p(\theta)$  is independent of  $\theta$  and there is no preferential direction for the slopes. As  $\gamma$  diminishes the slopes become more and more concentrated about the principal direction (see figure 5). When  $\gamma \ll 1$  we have in general

$$p( heta) = rac{\gamma}{2\pi\sin^2 heta},$$

which tends to zero as  $\gamma \rightarrow 0$ . But near the principal direction, i.e. when  $\theta$  is comparable with  $\gamma$ , we have

$$p( heta)=rac{\gamma}{2\pi(\gamma^2+ heta^2)},$$

a distribution whose width is proportional to  $\gamma$ . The integral of the distribution from  $\theta/\gamma = -\infty$  to  $\infty$  is equal to  $\frac{1}{2}$ . Thus the probability that  $\theta$  is near zero is  $\frac{1}{2}$ , and so also is the probability that  $\theta$  is near  $\pi$ . Hence it becomes almost certain that the gradient is nearly in the principal direction.

It should be noted that the probabilities so far discussed are for points distributed randomly and uniformly in the x, y plane. The corresponding probabilities for points selected so as to lie, for example, on a particular contour  $\zeta = \text{constant}$ , are different, as will be seen in §2.3.

### $2 \cdot 2$ . The number of zero-crossings along a line

As in §1.4, let us consider the curve in which the surface is intersected by the vertical plane  $x \sin \theta = y \cos \theta$ . A point where this curve passes through the mean level ( $\zeta = 0$ ) may be called a zero-crossing of  $\zeta$ . We shall now consider the number of zero-crossings of  $\zeta$  per unit distance x' measured along the line of intersection of the vertical plane and the mean level.

The mean number of zeros for a random function of a single variable has been derived by Rice (1944, 1945). We recall his argument briefly.  $\zeta$  and  $\partial \zeta / \partial x'$  are random functions which we shall denote by  $\xi_1$  and  $\xi_2$  respectively. Suppose that  $\zeta$  passes through zero at some point x' in the interval  $(x'_0, x'_0 + dx')$ , and with gradient  $\partial \zeta / \partial x'$  lying in the range  $(\xi_2, \xi_2 + d\xi_2)$ . Then at the point  $x' = x_0$  itself  $\zeta$  lies in the range  $(0, -\xi_2 dx')$ , approximately, i.e. a range of height  $d\xi_1 = |\xi_2| dx'$ . The probability of this occurrence is

$$p(0,\xi_2) | \xi_2 | dx' d\xi_2, \tag{2.2.1}$$

where  $p(\xi_1, \xi_2)$  is the joint-probability distribution of  $(\xi_1, \xi_2)$ . The total probability of a zero in  $(x'_0, x'_0 + dx')$  is found by integrating with respect to  $\xi_2$  from  $-\infty$  to  $\infty$ . Hence the total number  $N_0$  of zeros per unit distance is given by

$$N_0 = \int_{-\infty}^{\infty} p(0,\xi_2) |\xi_2| d\xi_2.$$
 (2.2.2)

Now the matrix of correlations for  $(\xi_1, \xi_2)$  is

$$(\Xi_{ij}) = \begin{pmatrix} m_0 & 0\\ 0 & m_2 \end{pmatrix},$$
 (2.2.3)

and so by  $(2 \cdot 1 \cdot 1)$ 

$$p(\xi_1,\xi_2) = \frac{1}{2\pi (m_0 m_2)^{\frac{1}{2}}} \exp\left\{-\xi_1^2/2m_0 - \xi_2^2/2m_2\right\}. \tag{2.2.4}$$

### M. S. LONGUET-HIGGINS ON THE

On substituting in  $(2 \cdot 2 \cdot 2)$  and carrying out the integration we find

$$N_0 = \frac{1}{\pi} \left( \frac{m_2(\theta)}{m_0(\theta)} \right)^{\frac{1}{2}}.$$
 (2.2.5)

In other words  $\pi N_0$  is equal to the r.m.s. wave-number in the direction  $\theta$ . It follows at once from § 1.4 that

(1) the number of zeros is a maximum and a minimum for two directions at right angles, given by

$$\tan 2\theta_p = \frac{2m_{11}}{m_{20} - m_{02}}; \qquad (2.2.6)$$

(2) the maximum and minimum values of  $N_0$  are given by

$$N_{0\,\text{max.}}, N_{0\,\text{min.}} = \frac{1}{\pi m_{00}^{\frac{1}{2}}} \left[ (m_{20} + m_{02}) \pm \sqrt{\{(m_{20} - m_{02})^2 + 4m_{11}^2\}} \right]^{\frac{1}{2}}; \qquad (2 \cdot 2 \cdot 7)$$

(3) the number of zeros in a general direction  $\theta$  is given by

$$N_0^2 = N_{0\,\text{max.}}^2 \cos^2(\theta - \theta_p) + N_{0\,\text{min.}}^2 \sin^2(\theta - \theta_p); \qquad (2.2.8)$$

(4) the ratio  $N_{0 \min}/N_{0 \max}$  is given by

$$\frac{N_{0\,\mathrm{min.}}}{N_{0\,\mathrm{max.}}} = \gamma, \qquad (2.2.9)$$

where  $\gamma^{-1}$  is the long-crestedness; for a narrow spectrum  $N_{0 \text{ min.}}/N_{0 \text{ max.}}$  equals the r.m.s. angular deviation of the energy from the mean direction.

There are similar relations for the mean number of crests and troughs along the curve, since these are simply zeros of the derivative  $\partial \zeta / \partial x'$ . The energy spectrum of  $\partial \zeta / \partial x'$  is  $u'^2$  times the energy spectrum of  $\zeta$ . So the mean number  $N_1$  of crests and troughs together is

$$N_1 = \frac{1}{\pi} \left( \frac{m_4(\theta)}{m_2(\theta)} \right)^{\frac{1}{2}}$$
(2·2·10)

(the number of crests or troughs separately is half this).  $m_2$  and  $m_4$  can be expressed in terms of the two-dimensional moments of E(u, v) by means of  $(1 \cdot 4 \cdot 12)$ .  $m_4$  is of the fourth degree in  $\cos \theta$  and  $\sin \theta$ , and it is found that in general  $N_1(\theta)$  has four maxima (in two pairs of opposite directions) and similarly four minima; these can be found, if necessary, in terms of the fourth-order moments  $m_{40}, m_{31}, \ldots, m_{04}$  and the second-order moments  $m_{20}, m_{11}, m_{02}$ .

In the same way the mean number  $N_2$  of points of inflexion on the curve is

$$N_2 = \frac{1}{\pi} \left( \frac{m_6(\theta)}{m_4(\theta)} \right), \tag{2.2.11}$$

and the maxima and minima of  $N_2$  are given in terms of the sixth-, fourth- and second-order moments of E(u, v).

We may find similarly the average number of times that the curve of intersection crosses the level  $\zeta = \xi_1$ . For in (2·2·1) and (2·2·2) it is necessary only to replace  $p(0, \xi_2)$  by  $p(\xi_1, \xi_2)$ . Since  $\xi_1$  and  $\xi_2$  are independent, this simply amounts to multiplying by a factor  $\exp\{-\xi_1^2/2m_0\}$ . So in the general case we have

$$N_0 = \frac{1}{\pi} \left( \frac{m_2(\theta)}{m_0(\theta)} \right)^{\frac{1}{2}} \exp\left\{ -\xi_1^2 / 2m_0(\theta) \right\}.$$
(2.2.12)

By similar reasoning, the number of times  $N_1$  per unit distance that the curve has a gradient  $\xi_2$  is given by

$$N_1 = \frac{1}{\pi} \left( \frac{m_4(\theta)}{m_2(\theta)} \right)^{\frac{1}{2}} \exp\left\{ -\xi_2^2 / 2m_2(\theta) \right\}, \tag{2.2.13}$$

and there are similar expressions corresponding to the higher derivatives of  $\zeta(x')$ .

### $2 \cdot 3$ . The length and direction of the contours

Let us consider now a corresponding property in two dimensions. Imagine the surface contours  $\zeta = \text{constant}$  to be drawn in the x, y plane. Contained in any region A of the plane there will be a certain length s of the contour  $\zeta = \zeta_0$ . The average length of contour, being proportional to the area A, may be denoted by  $\bar{s}A$ . The factor  $\bar{s}$  is now to be evaluated.



FIGURE 6. The length s of contour intercepted by a small element of area dA.

Let P be any fixed point in the plane, and dA the area of a small region surrounding P (see figure 6). Let s denote the length of a contour  $\zeta = \zeta_0$  intercepted by the region dA, and let n denote the perpendicular distance of the contour from P. Suppose that the magnitude  $\alpha$  and direction  $\theta$  of the gradient are fixed. Then the height  $\zeta$  of the surface at P is given by

$$|\zeta - \zeta_0| = \alpha n. \tag{2.3.1}$$

For the contour to cut the element of area, the perpendicular n and the height  $\zeta$  must lie in certain small ranges  $(n_1, n_2)$  and  $(\zeta_1, \zeta_2)$ . If now  $\alpha, \theta$  are allowed to vary within small ranges  $(\alpha, \alpha + d\alpha)$ ,  $(\theta + d\theta)$ , the expectation  $\bar{s}_{\alpha, \theta} dA d\alpha d\theta$  of s over the area dA is given by

$$\bar{s}_{\alpha,\theta} \mathrm{d}A \,\mathrm{d}\alpha \,\mathrm{d}\theta = \int_{\zeta_1}^{\zeta_2} sp(\zeta,\alpha,\theta) \,\mathrm{d}\zeta \,\mathrm{d}\alpha \,\mathrm{d}\theta = \int_{n_1}^{n_2} sp(\zeta,\alpha,\theta) \,\alpha \,\mathrm{d}n \,\mathrm{d}\alpha \,\mathrm{d}\theta, \qquad (2\cdot 3\cdot 2)$$

where  $p(\zeta, \alpha, \theta)$  denotes the joint distribution of  $\zeta$ ,  $\alpha$ ,  $\theta$  at P. Since  $\zeta$ ,  $\alpha$ ,  $\theta$  are nearly constant over the small range of integration of n we have

$$\bar{s}_{\alpha,\theta} dA = \alpha p(\zeta, \alpha, \theta) \int_{n_1}^{n_2} s dn = \alpha p(\zeta, \alpha, \theta) dA, \qquad (2.3.3)$$

and so

$$\bar{s}_{\alpha,\theta} = \alpha p(\zeta, \alpha, \theta). \tag{2.3.4}$$

Integrating over all possible values of  $\alpha$ ,  $\theta$  we have

$$\bar{s} = \int_0^\infty \int_0^{2\pi} \bar{s}_{\alpha,\theta} \, \mathrm{d}\alpha \, \mathrm{d}\theta = \int_0^\infty \int_0^{2\pi} \alpha p(\zeta, \alpha, \theta) \, \mathrm{d}\alpha \, \mathrm{d}\theta.$$
(2.3.5)

$$p(\zeta, \alpha, \theta) = p(\zeta) p(\alpha, \theta), \qquad (2.3.6)$$

Vol. 249. A.

Now from  $\S 2 \cdot 1$ 

where  $p(\zeta)$  is given by (2.1.8), with  $\xi_1 = \zeta$ , and  $p(\alpha, \theta)$  is given by (2.1.20). On substituting these values in (2.3.5) we have

$$\bar{s} = \frac{1}{(2\pi)^{\frac{3}{2}} (m_{00}\Delta_2)^{\frac{1}{2}}} \exp\left\{-\zeta^2/2m_{00}\right\} \int_0^\infty \int_0^{2\pi} \alpha^2 \exp\left\{-\alpha^2 (m_{02}\cos^2\theta + m_{20}\sin^2\theta)/2\Delta_2\right\} \mathrm{d}\alpha \,\mathrm{d}\theta.$$
(2·3·7)

(It has been supposed that the x axis is taken in the principal direction, so that  $m_{11}$  vanishes.) Integration with respect to  $\alpha$  gives

$$\bar{s} = \frac{\Delta_2}{4\pi m_{00}^{\frac{1}{2}}} \exp\left\{-\xi^2/2m_{00}\right\} \int_0^{2\pi} \frac{\mathrm{d}\phi}{(m_{02}\sin^2\phi + m_{20}\cos^2\phi)^{\frac{3}{2}}},$$
(2.3.8)

where  $\phi = \theta + \frac{1}{2}\pi$ . That is to say

$$\bar{s} = \frac{m_{02}}{\pi (m_{00}m_{20})^{\frac{1}{2}}} \exp\left\{-\zeta^2/2m_{00}\right\} \int_0^{\frac{1}{2}\pi} \frac{\mathrm{d}\phi}{(1-k^2\sin^2\phi)^{\frac{3}{2}}},$$
(2.3.9)

where

$$k^2 = 1 - \gamma^2. \tag{2.3.10}$$

Now since

$$\frac{\mathrm{d}}{\mathrm{d}\phi} \frac{k^2 \sin\phi \cos\phi}{(1-k^2 \sin^2\phi)^{\frac{1}{2}}} = (1-k^2 \sin^2\phi)^{\frac{1}{2}} + \frac{k^2-1}{(1-k^2 \sin^2\phi)^{\frac{3}{2}}}, \qquad (2\cdot3\cdot11)$$

it follows, on integration between 0 and  $\frac{1}{2}\pi$ , that

$$(1-k^2)\int_0^{\frac{1}{2}\pi} \frac{\mathrm{d}\phi}{(1-k^2\sin^2\phi)^{\frac{3}{2}}} = \int_0^{\frac{1}{2}\pi} (1-k^2\sin^2\phi)^{\frac{1}{2}} \,\mathrm{d}\phi = \mathbf{E}(k), \qquad (2\cdot3\cdot12)$$

where E(k) is Legendre's complete elliptic integral of the first kind (Legendre 1811). Hence we have finally

$$\bar{s} = \frac{1}{\pi} \left( \frac{m_{20} + m_{02}}{m_{00}} \right)^{\frac{1}{2}} \exp\left\{ -\zeta^2 / 2m_{00} \right\} (1 + \gamma^2)^{-\frac{1}{2}} \mathbf{E}\left\{ \sqrt{(1 - \gamma^2)} \right\}.$$
(2.3.13)

In the special case of long-crested waves, when  $\gamma = 0$ , we have E(1) = 1 and further

$$m_{20} = m_2(0), \quad m_{02} = 0,$$
 (2.3.14)

$$\bar{s} = \frac{1}{\pi} \left( \frac{m_2(0)}{m_{00}} \right)^{\frac{1}{2}} \exp\left\{ -\zeta^2 / 2m_{00} \right\}.$$
(2.3.15)

giving

Comparison with  $(2 \cdot 2 \cdot 12)$  shows what we might expect, namely, that the mean length of contour per unit area is equal to the mean number of crossings of the contour level per unit distance by a plane perpendicular to the wave crests.

In general  $(2 \cdot 3 \cdot 13)$  may be written

$$\bar{s} = \frac{1}{\pi} \left( \frac{m_{20} + m_{02}}{m_{00}} \right)^{\frac{1}{2}} \exp\left\{ -\zeta^2 / 2m_{00} \right\} f(\gamma), \qquad (2 \cdot 3 \cdot 16)$$

 $f(\gamma) = (1 + \gamma^2)^{-\frac{1}{2}} \mathbb{E}\{\sqrt{(1 - \gamma^2)}\}.$  (2.3.17)

This function is shown in figure 7. At the two extreme values we have

$$f(0) = 1 (2.3.18)$$

$$f(1) = \frac{\pi}{2\sqrt{2}} = 1.1107\dots$$
 (2.3.19)

and

Throughout its whole range the function departs very little from unity. There is, however, a weak singularity at the origin, where

$$f(\gamma) = 1 + \frac{1}{2}\gamma^2 \left( \ln \frac{4}{\gamma} - \frac{3}{2} \right) + O(\gamma^2 \ln \gamma).$$

$$(2 \cdot 3 \cdot 20)$$

A very closely related distribution is that of the direction  $\theta$  of the normal to a given contour. Let us suppose that  $\theta$  is measured at points randomly and uniformly distributed along the contour.  $\theta$  is also the direction of the surface gradient at the point of measurement. However, the distribution of  $\theta$  for a given contour is quite distinct from the distribution of  $\theta$  found in §2.1, where it was supposed that the angle was measured, not on a particular contour but at points randomly distributed in the x, y plane.



FIGURE 7. Graph of  $f(\gamma) = (1 + \gamma^2)^{-\frac{1}{2}} E[\sqrt{(1 - \gamma^2)}].$ 

To find the distribution  $p(\theta)_{\zeta}$  for the contour  $\zeta = \text{constant}$  we may note that the contribution of a given length of arc to the distribution of  $\theta$  in the interval  $(\theta, \theta + d\theta)$  is simply proportional to the expected length of arc for which  $\theta$  lies between these limits, that is,

$$p(\theta)_{\zeta} d\theta \propto \int_{0}^{\infty} \bar{s}_{\alpha,\theta} d\alpha d\theta.$$
 (2.3.21)

On normalizing the right-hand side by dividing by  $\bar{s}$  we have

$$p(\theta)_{\zeta} = \frac{1}{\overline{s}} \int_{0}^{\infty} \overline{s}_{\alpha,\theta} \, \mathrm{d}\alpha = \frac{1}{\overline{s}} \int_{0}^{\infty} \alpha p(\zeta, \alpha, \theta) \, \mathrm{d}\alpha.$$
 (2.3.22)

Substituting from  $(2 \cdot 3 \cdot 6)$  and  $(2 \cdot 3 \cdot 13)$ , and carrying out the integration we find

$$p(\theta)_{\zeta} = \frac{1}{4 \mathrm{E}\{\sqrt{(1-\gamma^2)}\}} \frac{\gamma^2}{(\gamma^2 \cos^2 \theta + \sin^2 \theta)^{\frac{3}{2}}}$$
(2.3.23)

43-2

(where  $\theta = 0$  is chosen as the principal direction). The form of this expression is somewhat similar to (2.1.37). When  $\gamma = 1$  (for short-crested waves)

$$p(\theta)_{\zeta} = \frac{1}{2\pi}, \qquad (2\cdot 3\cdot 24)$$

i.e. the contours have no preferential direction. As  $\gamma$  diminishes the distribution becomes more and more concentrated about the mean direction  $\theta = 0$ . When  $\gamma$  is small, we have for general directions

$$p(\theta)_{\zeta} = \frac{\gamma^2}{4\sin^3\theta},\tag{2.3.25}$$

which tends to zero as  $\gamma \rightarrow 0$ . But near the mean direction, that is, when  $\theta$  is comparable with  $\gamma$ ,

$$p(\theta)_{\zeta} = \frac{\gamma^2}{4(\gamma^2 + \theta^2)^{\frac{3}{2}}}.$$
 (2.3.26)

The integral of this expression from  $\theta/\gamma = -\infty$  to  $\infty$  is equal to  $\frac{1}{2}$ . Thus the probability that  $\theta$  is near zero is  $\frac{1}{2}$ , and so also is the probability that  $\theta$  is near  $\pi$ . Hence it becomes highly probable that the direction of the contour is near the principal direction.

### 2.4. The density of maxima and minima

Let us consider now the problem of how many maxima and minima (humps and hollows) the surface possesses, on the average, per unit area.

At a maximum or a minimum the two components of gradient  $\partial \zeta / \partial x$ ,  $\partial \zeta / \partial y$  must vanish. But not all such points are maxima or minima; we may also have a col or saddle-point, where the surface tends to rise in one pair of opposite directions and fall in another pair of opposite directions. We shall prove the following theorem:

On a statistically uniform surface the average density of maxima per unit area plus the average density of minima is equal to the average density of saddle-points, or

$$D_{\mathrm{ma.}} + D_{\mathrm{mi.}} = D_{\mathrm{sa.}}.$$
 (2.4.1)

Let a contour map of the surface be drawn, and let a direction  $\phi$  be assigned to each contour, say to the right when facing up-hill. Thus at each point of the plane, except the stationary points, there is a unique direction  $\phi$ . Consider now the variation of  $\phi$  round a small closed curve C on the map (see figure 8). C may at first be so small as to contain no stationary point, in which case  $\phi$  will return to its initial value after the circuit is completed (figure 8 a). If now C is expanded so as to enclose a single stationary point,  $\phi$  will increase by  $2\pi$  on completion of the circuit C if the stationary point is a maximum or a minimum (figure 8 b and c), and will decrease by  $2\pi$  if the stationary point is a saddle-point (figure 8 d). As C is further increased in size, so as to enclose  $d_{ma}$ . maxima,  $d_{mi}$ . minima and  $d_{sa}$ . saddle-points, say, the variation of  $\phi$  round C will be  $2\pi(d_{ma}+d_{mi}-d_{sa})$ , or  $2\pi A(D_{ma}+D_{mi}-D_{sa})$  approximately, where A is the area enclosed by C. But since the surface is statistically uniform, the variation of  $\phi$  round C will increase like  $L^2$ , supposing C is of constant shape. Thus  $2\pi(D_{ma}+D_{mi}-D_{sa})$  is proportional to  $L^{-1}$  at most, and letting L tend to infinity we see that  $(D_{ma}+D_{mi}-D_{sa})$  must vanish. This proves the result.

\* In fact it may be shown that the increase is proportional only to  $L^{\frac{1}{2}}$ .



FIGURE 8. Illustrating the way in which the contour direction varies round a curve enclosing (a) no stationary point, (b) a maximum, (c) a minimum and (d) a saddle-point.



FIGURE 9. (a) Stationary points on a surface which consists of two intersecting wave systems.  $\bullet = a$  maximum,  $\circ = a$  minimum, + = a saddle-point. (b) Stationary points in a hexagonal pattern.

A simple example is shown in figure 9a. The surface consists of two long-crested systems of waves of slowly varying amplitude. Where a crest from one system intersects a crest from the other system there is a maximum, and where two troughs intersect there is a minimum. But where a crest from one system intersects a trough from theother there is a saddle-point.

The pattern of saddle-points is similar and congruent to the pattern of maxima and minima together, so that  $(2\cdot 4\cdot 1)$  is satisfied.

On this surface the density of maxima is equal to the density of minima. But a case in which this is not so is illustrated in figure 9 *b*. Here the maxima are at the centres of the cells of a hexagonal honeycomb, the minima are at the vertices and the saddle-points are half-way along the edges. There are twice as many maxima as minima, and three times as many saddle-points, so that  $D = \frac{2}{2}D = \frac{1}{2}D = \frac{1}$ 

$$D_{\rm ma.} = \frac{2}{3} D_{\rm sa.}, \quad D_{\rm mi.} = \frac{1}{3} D_{\rm sa.}.$$
 (2.4.2)

In general it can be shown that the stationary points must form a cellular pattern, and the theorem  $(2\cdot 4\cdot 1)$  then follows from Euler's relation V+F=E+2 connecting the number of vertices V, faces F and edges E of a convex polyhedron (Euler 1752-3; Sommerville 1929, chap. IX).

The class of random surface represented by  $(1 \cdot 2 \cdot 1)$  satisfies the further relation

$$D_{\mathrm{ma.}} = D_{\mathrm{mi.}}$$
 (2.4.3)

For the phases  $\epsilon_n$  of the component waves are randomly and uniformly distributed between 0 and  $2\pi$ . The statistical properties of the surface are unaffected if a constant,  $\pi$ , is added to each phase. But this reverses the sign of  $\zeta$  and converts maxima into minima, and vice versa.

From  $(2\cdot 4\cdot 1)$  and  $(2\cdot 4\cdot 3)$  it follows that

$$D_{\rm sa.} = 2D_{\rm ma.} = 2D_{\rm mi.} \tag{2.4.4}$$

and if  $D_{\text{sta.}}$  denotes the total density of stationary points per unit area of the surface

$$D_{\rm sta.} = 2D_{\rm sa.} = 4D_{\rm ma.} = 4D_{\rm mi.}$$
 (2.4.5)

In other words, of all the stationary points on the surface, one-quarter are maxima, onequarter are minima, and the remaining half are saddle-points.

We proceed now to evaluate  $D_{\text{sta.}}$  in terms of the energy spectrum of  $\zeta$ . The variables entering the problem are  $\frac{\partial r}{\partial t} = \frac{\partial r}{\partial t}$ 

$$\frac{\partial \zeta}{\partial x}, \frac{\partial \zeta}{\partial y} = \xi_2, \xi_3 \tag{2.4.6}$$

and

where

$$\frac{\partial^2 \zeta}{\partial x^2}, \frac{\partial^2 \zeta}{\partial x \partial y}, \frac{\partial^2 \zeta}{\partial y^2} = \xi_4, \xi_5, \xi_6, \qquad (2.4.7)$$

say.  $(\xi_2, \xi_3)$  is a pair of functions of (x, y), and if (x, y) varies within a certain small region  $dA_1 = (x, x + dx; y, y + dy), (\xi_2, \xi_3)$  will vary within a region  $d\Sigma$  of area

$$\mathrm{d}\Sigma| = |J| |\mathrm{d}A|, \qquad (2\cdot4\cdot8)$$

$$J=rac{\partial(\xi_2,\xi_3)}{\partial(x,y)}=\xi_4\xi_6-\xi_5^2.$$

The probability that a given point, say a stationary point, lies in dA is equal to the probability that  $(\xi_2, \xi_3)$  lies in the corresponding region  $d\Sigma$ , which is

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} d\xi_4 d\xi_5 d\xi_6 \iint_{d\xi_2} d\xi_2 d\xi_3 p(\xi_2, \xi_3, \xi_4, \xi_5, \xi_6).$$
(2.4.10)

Since  $(\xi_2, \xi_3) = (0, 0)$  somewhere in  $d\Sigma$ ,  $p(\xi_2, \xi_3, \xi_4, \xi_5, \xi_6)$  may be replaced by  $p(0, 0, \xi_4, \xi_5, \xi_6)$ 

 $\mathbf{350}$ 

when  $d\Sigma$  is sufficiently small, and since

$$\iint_{\mathbf{d}\Sigma} \mathbf{d}\xi_2 \mathbf{d}\xi_3 = |\mathbf{d}\Sigma|, \qquad (2\cdot 4\cdot 11)$$

the above probability becomes

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} d\xi_4 d\xi_5 d\xi_6 | d\Sigma| p(0, 0, \xi_4, \xi_5, \xi_6).$$
(2.4.12)

On substituting from (2.4.8) we have for the probability of a stationary value of  $\zeta$  in dA,

$$D_{\text{sta.}} dA = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p(0, 0, \xi_4, \xi_5, \xi_6) |\xi_4 \xi_6 - \xi_5^2| d\xi_4 d\xi_5 d\xi_6 dA.$$
(2·4·13)

For a true maximum of the surface we must have  $\xi_4 \leq 0, \xi \leq_6 0$  and  $J \geq 0$ ; for a true minimum,  $\xi_4 \ge 0, \xi_6 \ge 0$  and  $J \ge 0$ . Thus the true maxima and minima correspond to the region of the  $(\xi_4, \xi_5, \xi_6)$  space given by

$$J = \xi_4 \xi_6 - \xi_5^2 \ge 0. \tag{2.4.14}$$

The boundary of this region is the surface J = 0, which is a cone with vertex at the origin. The remaining part of the  $(\xi_4, \xi_5, \xi_6)$  space corresponds to the saddle-points.

Now since the second derivatives  $\xi_4$ ,  $\xi_5$ ,  $\xi_6$  are uncorrelated with the first derivatives  $\xi_2, \xi_3 \text{ (see § 2.1) it follows that}$ 

$$p(\xi_2,\xi_3,\xi_4,\xi_5,\xi_6) = p(\xi_2,\xi_3) \, p(\xi_4,\xi_5,\xi_6), \qquad (2\cdot 4\cdot 15)$$

where  $p(\xi_2,\xi_3)$  is given by (2.1.12) and  $p(\xi_4,\xi_5,\xi_6)$  is the distribution for  $(\xi_4,\xi_5,\xi_6)$  independently of the other variables. The matrix of correlations is

$$(\Xi_{ij}) = \begin{pmatrix} m_{40} & m_{31} & m_{22} \\ m_{31} & m_{22} & m_{13} \\ m_{22} & m_{13} & m_{04} \end{pmatrix}, \qquad (2\cdot4\cdot16)$$

and hence

$$p(\xi_4,\xi_5,\xi_6) = \frac{1}{(2\pi)^{\frac{3}{2}} \Delta_4^{\frac{1}{2}}} \exp\left\{-\frac{1}{2} M_{ij} \xi_{i+3} \xi_{j+3}\right\}, \qquad (2.4.17)$$

where  $(M_{ij})$  is the inverse matrix to  $(\Xi_{ij})$  and

$$\Delta_4 = \mid \Xi_{ij} \mid. \tag{2.4.18}$$

Therefore, altogether we have for the density of stationary points

$$D_{\rm sta.} = \frac{1}{(2\pi)^{\frac{5}{2}} \Delta_2^{\frac{1}{2}} \Delta_4^{\frac{1}{2}}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp\left\{-\frac{1}{2} M_{ij} \xi_{i+3} \xi_{j+3}\right\} \left|\xi_4 \xi_6 - \xi_5^2\right| d\xi_4 d\xi_5 d\xi_6. \quad (2.4.19)$$

The density of maxima is given by a similar integral taken over the region  $\xi_4 \leq 0, \xi_6 \leq 0$ ,  $J \ge 0$ . The density of saddle-points is given by the same integral taken over the region J < 0.

Since  $(\Xi_{ij})$  is a positive-definite matrix, so also is its inverse  $(M_{ij})$ , and there exists a real linear transformation  $(\xi_4,\xi_5,\xi_6) = T(\eta_1,\eta_2,\eta_3)$  $(2 \cdot 4 \cdot 20)$ 

which simultaneously reduces the exponent in  $(2 \cdot 4 \cdot 19)$  to the unit form

$$M_{ij}\xi_{i+3}\xi_{j+3} = \eta_1^2 + \eta_2^2 + \eta_3^2 \tag{2.4.21}$$

and J to a diagonal form 
$$\xi_4 \xi_6 - \xi_5^2 = l_1 \eta_1^2 + l_2 \eta_2^2 + l_3 \eta_3^2$$
. (2.4.22)

### M. S. LONGUET-HIGGINS ON THE

The quantities  $l_1$ ,  $l_2$ ,  $l_3$  are easily found, for they are the roots of

$$|\sigma_{ij} - lM_{ij}| = 0, \qquad (2\cdot 4\cdot 23)$$

$$(\sigma_{ij}) = \begin{pmatrix} 0 & 0 & \frac{1}{2} \\ 0 & -1 & 0 \\ \frac{1}{2} & 0 & 0 \end{pmatrix}.$$
 (2.4.24)

On multiplying (2·4·23) by  $|\Xi_{ij}|$ ,  $= |M_{ij}^{-1}|$ , we have

$$|\Xi_{ij}\sigma_{jk}-l\delta_{ik}|=0, \qquad (2\cdot4\cdot25)$$

where  $\delta_{ij}$  is the unit matrix of order 3. In other words  $l_1, l_2, l_3$  are the latent roots of  $(\Xi_{ij}\sigma_{jk})$ :

$$\begin{vmatrix} \frac{1}{2}m_{22}-l & -m_{31} & \frac{1}{2}m_{40} \\ \frac{1}{2}m_{13} & -m_{22}-l & \frac{1}{2}m_{31} \\ \frac{1}{2}m_{04} & -m_{13} & \frac{1}{2}m_{22}-l \end{vmatrix} = 0.$$
 (2.4.26)

On expanding the determinant we find

where  $(\sigma_{ij})$  is the matrix of J:

$$4l^3 - 3Hl - \Delta_4 = 0,$$
 (2.4.27)

 $(2 \cdot 4 \cdot 28)$ 

 $(2 \cdot 4 \cdot 30)$ 

where Hence

 $l_1 + l_2 + l_3 = 0 \tag{2.4.29}$ 

and

It follows that one of the roots, say  $l_1$ , is positive and the other two, say  $l_2$ ,  $l_3$ , are negative. We write  $l_1 > 0 > l_2 \ge l_3$ . (2.4.31)

 $l_1 l_2 l_3 = \frac{1}{4} \Delta_4 > 0.$ 

 $3H = m_{40}m_{04} - 4m_{31}m_{13} + 3m_{22}^2$ .

The solution of the cubic equation  $(2 \cdot 4 \cdot 27)$  is

$$(l_1, l_2, l_3) = H^{\frac{1}{2}}(\cos\psi_1, \cos\psi_2, \cos\psi_3), \qquad (2.4.32)$$

where  $\psi_1, \psi_2, \psi_3$  are the roots of

$$\cos 3\psi = \Delta_4 / H^{\frac{3}{2}}.$$
 (2.4.33)

The modulus of the transformation T is

$$\frac{\partial(\xi_4,\xi_5,\xi_6)}{\partial(\eta_1,\eta_2,\eta_3)} = |M_{ij}|^{-\frac{1}{2}} = \Delta_4^{\frac{1}{2}}.$$
(2.4.34)

$$D_{\rm sta.} = \frac{1}{(2\pi)^{\frac{5}{2}} \Delta_2^{\frac{1}{2}}} I(l_1, l_2, l_3), \qquad (2 \cdot 4 \cdot 35)$$

where

$$I(l_1, l_2, l_3) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp\left\{-\frac{1}{2}(\eta_1^2 + \eta_2^2 + \eta_3^2)\right\} \left| l_1 \eta_1^2 + l_2 \eta_2^2 + l_3 \eta_3^2 \right| \mathrm{d}\eta_1 \mathrm{d}\eta_2 \mathrm{d}\eta_3. \quad (2 \cdot 4 \cdot 36)$$

The density of maxima is given by

$$D_{
m ma.} = rac{1}{(2\pi)^{rac{k}{2}}\Delta_2^{rac{1}{2}}} I'(l_1, l_2, l_3), \hspace{1.5cm} (2{\cdot}4{\cdot}37)$$

where 
$$I'(l_1, l_2, l_3) = \iiint_V \{-\frac{1}{2}(\eta_1^2 + \eta_2^2 + \eta_3^2)\} | l_1 \eta_1^2 + l_2 \eta_2^2 + l_3 \eta_3^2 | d\eta_1 d\eta_2 d\eta_3, \quad (2 \cdot 4 \cdot 38)$$

and V is the conical region

$$\eta_1 > 0, \quad l_1 \eta_1^2 + l_2 \eta_2^2 + l_3 \eta_3^2 > 0.$$
 (2·4·39)

Clearly 
$$4I' - I = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp\left\{-\frac{1}{2}(\eta_1^2 + \eta_2^2 + \eta_3^2)\right\} (l_1\eta_1^2 + l_2\eta_2^2 + l_3\eta_3^2) \,\mathrm{d}\eta_1 \,\mathrm{d}\eta_2 \,\mathrm{d}\eta_3$$
  
=  $(2\pi)^{\frac{3}{2}} (l_1 + l_2 + l_3),$  (2.4.40)

which vanishes, by (2.4.29). Thus

$$I = 4I', \quad D_{\text{sta.}} = 4D_{\text{ma.}},$$
 (2.4.41)

in agreement with  $(2\cdot 4\cdot 5)$ . The integral I' may be evaluated by means of the substitution

$$\eta_1 = l_1^{-\frac{1}{2}}r,$$

$$\eta_{2} = (-l_{2})^{-\frac{1}{2}} r \sin \theta \cos \chi,$$

$$\eta_{3} = (-l_{3})^{-\frac{1}{2}} r \sin \theta \sin \chi,$$

$$(2.4.42)$$

$$0 < r < \infty, \quad 0 < \theta < \frac{1}{2}\pi, \quad 0 < \chi < 2\pi.$$
 (2.4.43)

We have then

$$I' = \frac{1}{(l_1 l_2 l_3)^{\frac{1}{2}}} \int_0^\infty \mathrm{d}r \int_0^{\frac{1}{2}\pi} \mathrm{d}\theta \int_0^{2\pi} \mathrm{d}\chi \exp\left\{-\left(1 + f \sin^2 \theta\right) r^2 / 2l_1\right\} r^4 \cos^3 \theta \sin \theta, \quad (2 \cdot 4 \cdot 44)$$

where

where

$$f = f(\chi) = -\frac{l_1}{l_2} \cos^2 \chi - \frac{l_1}{l_3} \sin^2 \chi.$$
 (2.4.45)

Integration with respect to r gives

$$I' = 3(\frac{1}{2}\pi)^{\frac{1}{2}} \frac{l_1^2}{(l_2 l_3)^{\frac{1}{2}}} \int_0^{\frac{1}{2}\pi} \mathrm{d}\theta \int_0^{2\pi} \mathrm{d}\chi \, \frac{\cos^3\theta \, \sin\theta}{(1 + f \sin^2\theta)^{\frac{3}{2}}}.$$
 (2.4.46)

Further integration with respect to  $\theta$  gives

$$I' = 4(\frac{1}{2}\pi)^{\frac{1}{2}} \frac{l_1^2}{(l_2 l_3)^{\frac{1}{2}}} \int_0^{\frac{1}{2}\pi} d\chi \left[ \frac{1}{f} - \frac{2}{f^2} + \frac{2}{f^2(1+f)^{\frac{1}{2}}} \right].$$
(2.4.47)

This is an elliptic integral and may be evaluated by known methods (Legendre 1811). We find finally

$$\begin{split} I' &= (8\pi)^{\frac{1}{2}} \bigg[ (l_2 l_3)^{\frac{1}{2}} \Big\{ \Big( \frac{l_2 - l_1}{l_2} \Big)^{\frac{1}{2}} \operatorname{E}(k, \frac{1}{2}\pi) - \Big( \frac{l_2}{l_2 - l_1} \Big)^{\frac{1}{2}} \operatorname{F}(k, \frac{1}{2}\pi) \Big\} \\ &- (l_1 + l_2 + l_3) \left\{ \operatorname{F}(k', \phi) \operatorname{E}(k, \frac{1}{2}\pi) + \operatorname{E}(k', \phi) \operatorname{F}(k, \frac{1}{2}\pi) - \operatorname{F}(k', \phi) \operatorname{F}(k, \frac{1}{2}\pi) - \frac{1}{2}\pi \right\} \bigg], \quad (2 \cdot 4 \cdot 48) \end{split}$$

where E and F are the Legendre elliptic integrals of the first and second kind:

$$\begin{split} \mathbf{E}(k,\phi) &= \int_{0}^{\phi} (1-k^{2}\sin^{2}\phi)^{\frac{1}{2}} \,\mathrm{d}\phi, \\ \mathbf{F}(k,\phi) &= \int_{0}^{\phi} (1-k^{2}\sin^{2}\phi)^{-\frac{1}{2}} \,\mathrm{d}\phi, \end{split}$$
 (2.4.49)

and

$$k^{2} = \frac{l_{1}(l_{3} - l_{2})}{l_{3}(l_{1} - l_{2})}, \qquad k'^{2} = 1 - k^{2} = \frac{l_{2}(l_{1} - l_{3})}{l_{3}(l_{1} - l_{2})}, \qquad \phi = \tan^{-1} \left(-\frac{l^{3}}{l_{1}}\right)^{\frac{1}{2}}.$$
 (2.4.50)

If we now make use of the condition  $(2 \cdot 4 \cdot 29)$  we find for the density of maxima

$$D_{\rm ma.} = \frac{1}{2\pi^2} \frac{(l_2 l_3)^{\frac{1}{2}}}{\Delta_2^{\frac{1}{2}}} \left[ \left( \frac{l_2 - l_1}{l_2} \right)^{\frac{1}{2}} \mathbf{E}(k, \frac{1}{2}\pi) - \left( \frac{l_2}{l_2 - l_1} \right)^{\frac{1}{2}} \mathbf{F}(k, \frac{1}{2}\pi) \right].$$
(2.4.51)

Vol. 249. A.

The density of stationary points  $D_{sta}$  is four times this value.

Equation  $(2 \cdot 4 \cdot 51)$  may also be written

$$D_{\rm ma.} = \frac{1}{2\pi^2} \frac{l_1}{\Delta_2^{\frac{1}{2}}} \Phi(-l_2/l_1), \qquad (2\cdot 4\cdot 52)$$

where

$$\Phi(\alpha) = \{\alpha(1-\alpha)\}^{\frac{1}{2}} \left[ \left( \frac{1+\alpha}{\alpha} \right)^{\frac{1}{2}} \mathbf{E}(k, \frac{1}{2}\pi) - \left( \frac{\alpha}{1+\alpha} \right)^{\frac{1}{2}} \mathbf{F}(k, \frac{1}{2}\pi) \right], \\
k^{2} = \frac{1-2\alpha}{1-\alpha^{2}} \quad (0 < \alpha \leq \frac{1}{2}).$$
(2.4.53)



FIGURE 10. Graph of  $\Phi(\alpha)$  (defined by  $(2 \cdot 4 \cdot 53)$ ).

The form of  $\Phi(\alpha)$  is shown in figure 10. When  $\alpha \to 0, k^2 \to 1$  and  $F(k, \frac{1}{2}\pi) \to \infty$  logarithmically. Hence  $\alpha F(k, \frac{1}{2}\pi) \to 0$  and  $\lim \Phi(\alpha) = 1.$  (2.4.54)

Also when 
$$\alpha = \frac{1}{2}$$
,  $k^2 = 0$  and so

$$\Phi(\frac{1}{2}) = (3^{\frac{1}{2}} - 3^{-\frac{1}{2}}) \pi/4 = 0.917 \dots$$
(2.4.55)

Throughout the whole of its range,  $\Phi$  departs very little from unity.

A particularly interesting case is when the energy spectrum E(u, v) is narrow. Let us take axes of (u, v) so that the *u* axis passes through the centroid  $(\bar{u}, \bar{v})$ , making  $\bar{v} = 0$ . Then on substituting from (1.6.8 and 1.6.9) and retaining only the terms of highest order we find

$$\begin{aligned} \Delta_{2} &= \overline{u}^{2} \mu_{00} \mu_{02}, \\ 3H &= \overline{u}^{4} (\mu_{00} \mu_{04} + 3\mu_{02}^{2}), \\ \Delta_{4} &= \overline{u}^{6} (\mu_{00} \mu_{02} \mu_{04} - \mu_{00} \mu_{03}^{2} - \mu_{02}^{3}). \end{aligned}$$

$$(2.4.56)$$

Now  $\mu_{00}$ ,  $\mu_{02}$ ,  $\mu_{03}$ ,  $\mu_{04}$  are moments of the energy spectrum  $E_{\frac{1}{2}\pi}$  of a section of the surface at right angles to the mean direction. We have

$$\mu_{02} = (\gamma \overline{u})^2 m_{00}, \quad \mu_{03} = b(\gamma \overline{u})^3 m_{00}, \quad \mu_{04} = a^2 (\gamma \overline{u})^4 m_{00}, \quad (2.4.57)$$

where  $\gamma^{-1}$  is the long-crestedness (defined in §1.4),  $a^2$  is a non-dimensional parameter (always greater than 1) which represents the *peakedness* of the spectrum  $E_{\frac{1}{2}\pi}$ , and b is a

measure of the asymmetry of  $E_{\frac{1}{2}\pi}$  about its mean (but is independent of the angle of skewness  $\beta$ ). If we assume  $\mu_{03} = 0$  and so b = 0, we have

$$\begin{split} &\Delta_2 = (\gamma \overline{u}^2)^2 \, m_{00}^2, \\ &H = (\gamma \overline{u}^2)^4 \, m_{00}^2 (1 + a^2/3), \\ &\Delta_4 = (\gamma \overline{u}^2)^6 \, m_{00}^3 (a^2 - 1) \end{split}$$

and so from  $(2 \cdot 4 \cdot 32)$ 

$$(l_1, l_2, l_3) = (\gamma \overline{u}^2)^2 m_{00} (1 + a^2/3)^{\frac{1}{2}} (\cos \psi_1, \cos \psi_2, \cos \psi_3), \qquad (2 \cdot 4 \cdot 59)$$

where

$$\cos 3\psi = \frac{a^2 - 1}{(1 + a^2/3)^{\frac{3}{2}}}.$$
 (2.4.60)

Thus from  $(2 \cdot 4 \cdot 52)$  we have

$$D_{\rm ma.} = \gamma \overline{u}^2 C(a), \qquad (2.4.61)$$

where

and hence

$$C(a) = \frac{1}{2\pi^2} (1 + a^2/3)^{\frac{1}{2}} \cos \psi_1 \Phi\left(-\frac{\cos \psi_2}{\cos \psi_1}\right), \qquad (2.4.62)$$

which is a quantity depending only on the peakedness a. For a given peakedness,  $D_{ma.}$  is proportional to the square of the wave-number of the carrier wave and inversely proportional to the long-crestedness  $\gamma^{-1}$ . To illustrate the effect of varying peakedness, C(a)has been computed for a number of different values of a, including some interesting special cases. The results are given in table 1.

### TABLE 1

$a^2$	C(a)	$a^2$	C(a)	$a^2$	C(a)
1	0.0507	4	0.0695	8	0.0880
<u>9</u> 5	0.0562	5	0.0747	9	0.0919
$\check{2}$	0.0578	6	0.0794	10	0.0956
3	0.0639	7	0.0838	<b>20</b>	0.1265

 $a^2 = 1$  corresponds to a pair of intersecting wave trains, for then  $\Delta_4$  vanishes (by  $(2\cdot 4\cdot 54)$ ), which is the condition for the spectrum to degenerate into two one-dimensional spectra (equation  $(1\cdot 3\cdot 7)$ ). In the limit as  $a^2 \rightarrow 1$  we find from  $(2\cdot 4\cdot 60)$  that

$$(\psi_1, \psi_2, \psi_3) = (\pm \frac{1}{6}\pi, \pm \frac{1}{2}\pi, \pm \frac{5}{6}\pi),$$
  
 $C = \frac{1}{2\pi^2} = 0.05066....$  (2.4.63)

This is what we should expect, for the wavelengths of the pattern in the *u* direction and the v direction are  $2\pi/\bar{u}$  and  $2\pi/\gamma\bar{u}$  respectively. Reference to figure 9 *a* will show that each maximum is at the centre of a parallelogram (bounded by troughs) whose diagonals are of length  $2\pi/\bar{u}$  and  $2\pi/\gamma\bar{u}$ , and whose area is therefore  $2\pi^2/\gamma\bar{u}^2$ . The density of maxima is the reciprocal of this area, i.e.  $\gamma\bar{u}^2/2\pi^2$ .

The case  $a^2 = \frac{9}{5}$  has been included, since this is the peakedness of a low-pass spectrum, when the wave energy is uniformly distributed with regard to direction over a narrow sector.  $a^2 = 3$  corresponds to a normal distribution of energy with regard to direction. Another special case is  $a^2 = 9$ , when  $(\psi_1, \psi_2, \psi_3) = (0, \pm \frac{2}{3}\pi, \pm \frac{2}{3}\pi)$ , and so

$$C = \frac{1}{4\pi} \left( 3^{\frac{1}{2}} - 3^{-\frac{1}{2}} \right) = 0.09189 \dots$$
(2.4.64)
  
44-2

M. S. LONGUET-HIGGINS ON THE

Finally as  $a \rightarrow \infty$  we find

356

$$C \sim \frac{a}{4\pi^2} \rightarrow \infty.$$
 (2.4.65)

On the whole, however, the variation of C with a is slight. As  $a^2$  increases from 0 to 10, C is less than doubled.

The density of *specular points*, i.e. points on the surface where the gradient takes a given value, not necessarily zero, may be found similarly. For it is only necessary to replace  $p(0, 0, \xi_4, \xi_5, \xi_6)$  in equation  $(2 \cdot 4 \cdot 13)$  by  $p(\xi_2, \xi_3, \xi_4, \xi_5, \xi_6)$ , and, from equation  $(2 \cdot 4 \cdot 16)$ , this amounts to multiplying by the exponential factor in  $(2 \cdot 1 \cdot 12)$ . So the density  $D_{sp}$  of specular points with gradient  $(\xi_2, \xi_3)$  is given by

$$D_{\rm sp.} = 4D_{\rm ma.} \exp\left\{-\left(m_{02}\xi_2^2 - 2m_{11}\xi_2\xi_3 + m_{20}\xi_3^2\right)/2\Delta_2\right\}, \qquad (2\cdot 4\cdot 66)$$

where  $D_{\text{ma.}}$  is the density of maxima.

#### $2 \cdot 5$ . The velocities of zeros along a line

In this and the following two sections will be considered some statistical properties of the surface which depend on its motion, that is to say properties involving the time t.

Let  $\zeta(x', t)$  denote the curve in which the surface is intersected by a fixed vertical plane in direction  $\theta$ . Consider the movement of a point where the curve crosses a fixed level  $\zeta$ . If x' and x' + dx' are its co-ordinates at two successive instants of time t and t + dt, then we have

$$0 = d\zeta = \frac{\partial \zeta}{\partial x'} dx' + \frac{\partial \zeta}{\partial t} dt. \qquad (2.5.1)$$

Therefore the velocity of the point is given by

$$c = \frac{\mathrm{d}x'}{\mathrm{d}t} = -\frac{\partial\zeta/\partial t}{\partial\zeta/\partial x'}.$$
(2.5.2)

Consider now the statistical distribution of *c*.

Let the variables  $\zeta$ ,  $\partial \zeta / \partial x'$ ,  $\partial \zeta / \partial t$  be denoted by  $\xi_1$ ,  $\xi_2$ ,  $\xi_3$  respectively, so that  $c = -\xi_3/\xi_2$ , and let  $p(\xi_1, \xi_2, \xi_3)$  denote the joint distribution of  $\xi_1, \xi_2, \xi_3$  at an arbitrary point x' on the plane section. The probability distribution of  $\xi_2$ ,  $\xi_3$  at points x' where  $\xi_1$  takes a given value will be denoted by  $p(\xi_2, \xi_3)_{\xi_1}$ . This may be found as follows. If (x', x' + dx') is any fixed interval of distance, the probability of  $\xi_1$  taking the given value in (x', x' + dx') is

$$N_0(\xi_1) \,\mathrm{d}x'$$
 (2.5.3)

(evaluated in § 2.2). But if, at the point x', the variables  $\xi_1, \xi_2, \xi_3$  lie in certain ranges of width  $d\xi_1, d\xi_2, d\xi_3$ , then we have  $d\xi_1 = |\xi_2| dx'$ , (2.5.4)

so that the probability of  $\xi_1$  taking the given value in (x', x' + dx') and of  $\xi_2$ ,  $\xi_3$  lying in the given range is  $p(\xi_1, \xi_2, \xi_3) d\xi_1 d\xi_2 d\xi_3 = p(\xi_1, \xi_2, \xi_3) |\xi_1| dx' d\xi_2 d\xi_3.$  (2.5.5)

The probability of  $\xi_2$ ,  $\xi_3$  lying in the ranges  $d\xi_2$ ,  $d\xi_3$  given that  $\xi_1$  crosses the given level in (x', x' + dx') is the quotient of  $(2 \cdot 5 \cdot 5)$  and  $(2 \cdot 5 \cdot 3)$ . Hence

$$p(\xi_2,\xi_3)_{\xi_1} = \frac{p(\xi_1,\xi_2,\xi_3) |\xi_2|}{N_0(\xi_1)}.$$
(2.5.6)

STATISTICAL ANALYSIS OF A RANDOM, MOVING SURFACE 357

Now the matrix of correlations for  $\xi_1, \xi_2, \xi_3$  is

$$(\Xi_{ij}) = egin{pmatrix} m_0 & 0 & 0 \ 0 & m_2 & m_1' \ 0 & m_1' & m_0'' \end{pmatrix}, \qquad (2.5.7)$$

and therefore

in  $(2\cdot 5\cdot 10)$ , so that

$$p(\xi_1,\xi_2,\xi_3) = \frac{1}{(2\pi)^{\frac{3}{2}} (m_0 \Delta_{2'})^{\frac{1}{2}}} \exp\left\{-\xi_1^2/2m_0\right\} \exp\left\{-(m_0''\xi_2^2 - 2m_1'\xi_2\xi_3 + m_2\xi_3^2)/2\Delta_{2'}\right\}, \quad (2\cdot5\cdot8)$$
  
ere 
$$\Delta_{2'} = m_2 m_0'' - m_1'^2. \quad (2\cdot5\cdot9)$$

where

 $\Delta_{2'} = m_2 m_0'' - m_1'^2.$ 

From this and  $(2 \cdot 2 \cdot 12)$  we have

$$p(\xi_2,\xi_3)_{\xi_1} = \frac{1}{2(2\pi)^{\frac{1}{2}}(m_2\Delta_{2'})^{\frac{1}{2}}} |\xi_2| \exp\{-(m_0''\xi_2^2 - 2m_1'\xi_2\xi_3 + m_2\xi_3^2)/2\Delta_{2'}\}.$$
 (2.5.10)

We require now the statistical distribution of  $-\xi_3/\xi_2$ . Writing

$$-\xi_3/\xi_2 = c, \quad \xi_3 = c'$$
 (2.5.11)

$$\frac{\partial(c,c')}{\partial(\xi_2,\xi_3)} = \frac{\xi_3}{\xi_2^2} = \frac{c^2}{c'},$$
(2.5.12)

we have 
$$p(c,c')_{\xi_1} = \frac{1}{2(2\pi)^{\frac{1}{2}} (m_2 \Delta_{2'})^{\frac{1}{2}}} \left| \frac{c'^2}{c^3} \right| \exp\left\{ - (m_0''/c^2 + 2m_1'/c + m_2) c'^2/2\Delta_{2'} \right\}.$$
 (2.5.13)

The distribution of c is found by integrating with respect to c' from  $-\infty$  to  $\infty$ . Thus

$$p(c)_{\xi_1} = \frac{1}{2} \frac{\Delta_{2'}/m_2^{\frac{1}{2}}}{(m_0'' + 2m_1'c + m_2c^2)^{\frac{3}{2}}}, \qquad (2.5.14)$$

or

$$p(c)_{\xi_1} = \frac{1}{2} \frac{\Delta_{2'}/m_2^2}{\left[(c - \bar{c})^2 + \Delta_{2'}/m_2^2\right]^{\frac{3}{2}}},$$
(2.5.15)

where

$$\bar{c} = -m_1'/m_2.$$
 (2.5.16)

This distribution has a maximum or mode when  $c = \overline{c}$  and is symmetrical about this mean value. The second moment of the distribution is divergent, but the interquartile range is 0 11 given by

$$\frac{2}{\sqrt{3}}\frac{\Delta_{2'}^2}{m_2}.$$
 (2.5.17)

It will be seen that the distribution  $(2\cdot 5\cdot 15)$  is independent of the height  $\xi_1$  at which the velocity is measured (provided this height is constant).

We may consider similarly the distribution of velocities of points on the curve having a given gradient (say, zero). The velocity of such a point is given by

$$c_1 = -\frac{\partial^2 \zeta / \partial x' \, \partial t}{\partial^2 \zeta / \partial x'^2}. \tag{2.5.18}$$

Hence the probability distribution is the same as for the velocities of zeros, except that the index of each of the moments is increased by two. Thus

$$p(c_1) = \frac{1}{2} \frac{\Delta_{4'}/m_4^2}{\left[(c - \bar{c}_1)^2 + \Delta_{4'}/m_4^2\right]^{\frac{3}{2}}},$$
(2.5.19)

$$\Delta_{4'} = m_4 m_2'' - m_3'^2 \tag{2.5.20}$$

$$\bar{c}_1 = -m'_3/m_4.$$
 (2.5.21)

where and

This is a distribution with mean  $\bar{c}_1$  and interquartile range equal to

$$\frac{2}{\sqrt{3}}\frac{\Delta_{4'}^{\frac{1}{2}}}{m_4}.$$
 (2.5.22)

Similar distributions can be written down for the velocities of points having given higher derivatives of  $\zeta$ .

Let us interpret the above results for a narrow spectrum. Without loss of generality we may take the u axis to pass through the centroid. On expanding in a Taylor series about this point we have

$$m_{n} = u'^{n} \mu_{00} + \frac{1}{2} \left( \mu_{20} \frac{\partial^{2}}{\partial u^{2}} + 2\mu_{11} \frac{\partial^{2}}{\partial u \partial v} + \mu_{02} \frac{\partial^{2}}{\partial v^{2}} \right) u'^{n} + \dots,$$

$$m_{n}' = u'^{n} \sigma \mu_{00} + \frac{1}{2} \left( \mu_{20} \frac{\partial^{2}}{\partial u^{2}} + 2\mu_{11} \frac{\partial^{2}}{\partial u \partial v} + \mu_{02} \frac{\partial^{2}}{\partial v^{2}} \right) (u'^{n} \sigma) + \dots,$$

$$m_{n}'' = u'^{n} \sigma^{2} \mu_{00} + \frac{1}{2} \left( \mu_{20} \frac{\partial^{2}}{\partial u^{2}} + 2\mu_{11} \frac{\partial^{2}}{\partial u \partial v} + \mu_{02} \frac{\partial^{2}}{\partial v^{2}} \right) (u'^{n} \sigma^{2}) + \dots,$$
(2.5.23)

where u', v' and  $\sigma$  are to be evaluated at  $(\bar{u}, 0)$ . Suppose that  $\theta = 0$ , that is, let the plane of intersection be taken parallel to the principal direction. Then  $u' \equiv u$  and,  $\sigma$  being a function of  $(u^2+v^2)$  only,  $\partial \sigma = \partial^2 \sigma$ 

$$rac{\partial\sigma}{\partial v} = 0, \quad rac{\partial^2\sigma}{\partial u \,\partial v} = 0.$$
 (2.5.24)

From  $(2 \cdot 5 \cdot 16)$  and  $(2 \cdot 5 \cdot 21)$  we find first

$$\bar{c} = \bar{c}_1 = -\sigma/\mu, \qquad (2 \cdot 5 \cdot 25)$$

showing that the mean velocities of zeros and of specular points are equal to the phase velocity of the carrier wave. Also from (2.5.9) and (2.5.20)

$$\begin{aligned} \Delta_{2'} &= \mu_{00} \mu_{20} u^2 (\sigma/u - \partial \sigma/\partial u)^2, \\ \Delta_{4'} &= \mu_{00} \mu_{20} u^6 (\sigma/u - \partial \sigma/\partial u)^2, \end{aligned}$$
 (2.5.26)

$$\frac{\Delta_{2'}^{\frac{1}{2}}}{m_2} = \frac{\Delta_{4'}^{\frac{1}{2}}}{m_4} = \left(\frac{\mu_{20}}{\mu_{00}}\right)^{\frac{1}{2}} \frac{\sigma/u - \partial\sigma/\partial u}{u}.$$
(2.5.27)

so that

Thus the interquartile ranges of both the velocities of zeros and the velocities of specular points are equal to

$$\frac{2}{\sqrt{3}}\nu(\bar{c}-\Gamma), \qquad (2.5.28)$$

where  $\nu$  is defined by (1.6.15) and  $\Gamma = \partial \sigma / \partial u$  is the group velocity of the carrier wave. Thus we see that the width of the velocity distribution depends both on the r.m.s. width of the spectrum (given by  $\nu u$ ) and also on the dispersive properties of the medium. If the medium is non-dispersive,  $\Gamma = \bar{c}$  and so (2.5.28) vanishes. This is what we should expect, since in a non-dispersive medium a long-crested disturbance advances without change of form and the zeros and specular points move with uniform velocity in the direction of wave propagation.

For gravity waves in deep water the group velocity is half the wave velocity and so the interquartile range equals  $\nu \bar{c}/\sqrt{3}$ .

### $2 \cdot 6$ . The motion of the contours

Consider first how to define this motion. Let P be a point in the plane lying on the contour  $\zeta = \text{constant}$ . A moment later the contour at the same level will have moved to a new position, say QSR (see figure 11 a). If PQ and PR are axes parallel to Ox and Oy, the rates at which PQ and PR are increasing, which we denote by  $c_x$  and  $c_y$ , define the local displacement of the contour uniquely, and we have



FIGURE 11. Definition of the motion of a contour (a) by its intercept on an arbitrary line, (b) by its normal displacement.

However, if we take a line through P in an arbitrary direction  $\theta$ , and if this line intersects the displaced contour in S (figure 11), then it may be shown that

$$\frac{1}{PS} = \frac{1}{PQ}\cos\theta + \frac{1}{PR}\sin\theta, \qquad (2.6.2)$$

and so if c is the rate at which the intercept PS is increasing

$$\frac{1}{c} = \frac{1}{c_x} \cos \theta + \frac{1}{c_y} \sin \theta.$$
(2.6.3)

This shows that the reciprocal quantity  $(1/c_x, 1/c_y)$  is transformed like a vector, but not  $(c_x, c_y)$  itself. It is therefore more appropriate, and in fact more convenient, to consider the distribution of  $(1/c_x, 1/c_y) = (1/c_x, 1/c_y) = (1/c_x, 1/c_y)$ 

$$(\kappa_x, \kappa_y) = (1/c_x, 1/c_y) = \left(-\frac{\partial \zeta/\partial x}{\partial \zeta/\partial t}, -\frac{\partial \zeta/\partial y}{\partial \zeta/\partial t}\right), \qquad (2.6.4)$$

rather than the distribution of  $(c_x, c_y)$ . However, each may be derived from the other by a simple substitution. For since

$$\frac{\partial(\kappa_x,\kappa_y)}{\partial(c_x,c_y)} = \frac{1}{c_x^2 c_y^2} = \kappa_x^2 \kappa_y^2, \qquad (2.6.5)$$

$$p(c_x, c_y) = \frac{1}{c_x^2 c_y^2} p(\kappa_x, \kappa_y). \qquad (2 \cdot 6 \cdot 6)$$

The distribution of the velocities of the contours normal to themselves can also be found. In figure 11 b, T is the foot of the perpendicular from P to QR, and TU, TV are drawn perpendicular to PQ, PR. It can be shown that

$$PU = \frac{PQPR^2}{QR^2}, \quad PV = \frac{PQ^2PR}{QR^2}, \quad (2.6.7)$$

and hence the components of the normal velocity are

$$(q_x, q_y) = \left(\frac{c_x c_y^2}{c_x^2 + c_y^2}, \frac{c_x^2 c_y}{c_x^2 + c_y^2}\right) = \left(\frac{\kappa_x}{\kappa_x^2 + \kappa_y^2}, \frac{\kappa_y}{\kappa_x^2 + \kappa_y^2}\right).$$
(2.6.8)

Solving, we have

$$(\kappa_x,\kappa_y) = \left(\frac{q_x}{q_x^2 + q_y^2}, \frac{q_y}{q_x^2 + q_y^2}\right), \quad \frac{\partial(\kappa_x,\kappa_y)}{\partial(q_x,q_y)} = -\frac{1}{(q_x^2 + q_y^2)^2}, \quad (2.6.9)$$

and so

$$p(q_x, q_y) = \frac{1}{(q_x^2 + q_y^2)^2} p(\kappa_x, \kappa_y).$$
(2.6.10)

Let us write

$$\zeta, \frac{\partial \zeta}{\partial x}, \frac{\partial \zeta}{\partial y}, \frac{\partial \zeta}{\partial t} = \xi_1, \xi_2, \xi_3, \xi_4, \qquad (2.6.11)$$

and let  $p(\xi_1, \xi_2, \xi_3, \xi_4)$  denote the joint distribution of  $\xi_1, \xi_2, \xi_3, \xi_4$  at an arbitrary point Pin the (x, y) plane. We require the joint distribution of  $\xi_2, \xi_3, \xi_4$  at points distributed uniformly along the contour  $\xi_1 = \text{constant}$ . Let this be denoted by  $p(\xi_2, \xi_3, \xi_4)_{\xi_1}$ . To find this distribution let dA be a small area surrounding P. If  $\xi_2, \xi_3, \xi_4$  at P are restricted to lie in certain ranges of width  $d\xi_2, d\xi_3, d\xi_4$ , the contribution of the area dA to the distribution over these ranges is, by the argument of § 2.3,

$$p(\xi_1,\xi_2,\xi_3,\xi_4) \,\alpha \mathrm{d}A \,\mathrm{d}\xi_2 \,\mathrm{d}\xi_3 \,\mathrm{d}\xi_4, \qquad (2 \cdot 6 \cdot 12)$$

where  $\alpha = (\xi_2^2 + \xi_3^2)^{\frac{1}{2}}$ . But the total expectation of contour length over the area dA is  $\overline{s} dA$  (see §2·3). Hence we have

$$p(\xi_2,\xi_3,\xi_4)_{\xi_1} = \frac{(\xi_2^2 + \xi_3^2)^{\frac{1}{2}} p(\xi_1,\xi_2,\xi_3,\xi_4)}{\overline{s}}.$$
(2.6.13)

Now by § 2·1, the elevation  $\xi_1$  is uncorrelated with the first derivatives  $\xi_2$ ,  $\xi_3$ ,  $\xi_4$ . Therefore

$$p(\xi_1,\xi_2,\xi_3,\xi_4) = p(\xi_1) p(\xi_2,\xi_3,\xi_4), \qquad (2.6.14)$$

where  $p(\xi_1)$  is given by (2.1.8). The matrix of correlations for  $(\xi_2, \xi_3, \xi_4)$  is

$$(\Xi_{ij}) = \begin{pmatrix} m_{20} & m_{11} & m'_{10} \\ m_{11} & m_{02} & m'_{01} \\ m'_{10} & m'_{01} & m''_{00} \end{pmatrix}.$$
 (2.6.15)

Hence

$$p(\xi_2,\xi_3,\xi_4) = \frac{1}{(2\pi)^{\frac{3}{2}} \Delta_3^{\frac{1}{2}}} \exp\left\{-\frac{1}{2} M_{ij} \xi_{i+1} \xi_{j+1}\right\}, \qquad (2.6.16)$$

where  $(M_{ij})$  is the inverse matrix to  $(\Xi_{ij})$  and where  $\Delta_3 = |\Xi_{ij}|$ . On substituting these values in  $(2 \cdot 6 \cdot 13)$  we have

$$p(\xi_2,\xi_3,\xi_4)_{\xi_1} = \frac{1}{4\pi (m_{20}+m_{02})^{\frac{1}{2}} \Delta_3^{\frac{1}{2}}} \frac{(1+\gamma^2)^{\frac{1}{2}}}{\mathbf{E}\{\sqrt{(1-\gamma)^2}\}} (\xi_2^2+\xi_3^2)^{\frac{1}{2}} \exp\{-\frac{1}{2}M_{ij}\xi_{i+1}\xi_{j+1}\}. \quad (2.6.17)$$

$$\kappa_x = -\xi_2/\xi_4, \quad \kappa_y = -\xi_3/\xi_4, \quad \kappa = \xi_4$$
(2.6.18)

Writing now

$$\frac{\partial(\kappa_x,\kappa_y,\kappa)}{\partial(\xi_2,\xi_3,\xi_4)} = \frac{1}{\xi_4^2} = \frac{1}{\kappa^2},$$
(2.6.19)

so that we have

### STATISTICAL ANALYSIS OF A RANDOM, MOVING SURFACE 361

To obtain the distribution of  $\kappa_x$ ,  $\kappa_y$ , we eliminate  $\kappa$  by integrating from  $-\infty$  to  $\infty$ ; thus

$$p(\kappa_x,\kappa_y)_{\xi_1} = \frac{1}{\pi \Delta_3^{\frac{1}{2}} (m_{20} + m_{02})^{\frac{1}{2}}} \frac{(1+\gamma^2)^{\frac{1}{2}}}{\mathrm{E}\{\sqrt{(1-\gamma^2)}\}} \frac{(\kappa_x^2 + \kappa_y^2)^{\frac{1}{2}}}{R^2}, \qquad (2.6.21)$$

where

$$R = M_{11}\kappa_x^2 + 2M_{12}\kappa_x\kappa_y + M_{22}\kappa_y^2 - 2M_{13}\kappa_x - 2M_{23}\kappa_y + M_{33}.$$
 (2.6.22)

We may also write

$$R = M_{11}(\kappa_x - \bar{\kappa}_x)^2 + 2M_{12}(\kappa_x - \bar{\kappa}_x) (\kappa_y - \bar{\kappa}_y) + M_{22}(\kappa_y - \bar{\kappa}_y)^2 + M, \qquad (2.6.23)$$

$$= M_{13}M_{22} - M_{12}M_{22} - m'_{10} \qquad )$$

where

$$\kappa_{x} = \frac{M_{11}M_{22} - M_{12}^{2}}{M_{11}M_{22} - M_{12}^{2}} = \frac{m_{01}'}{m_{00}''},$$

$$\bar{\kappa}_{y} = \frac{M_{11}M_{23} - M_{12}M_{13}}{M_{11}M_{22} - M_{12}^{2}} = -\frac{m_{01}'}{m_{00}''},$$

$$M = M_{33} - (M_{11}\bar{\kappa}_{x}^{2} + 2M_{12}\bar{\kappa}_{x}\bar{\kappa}_{y} + M_{22}\bar{\kappa}_{y}^{2}) = \frac{1}{m_{00}''}.$$

$$(2.6.24)$$

The denominator R is thus a symmetrical expression with a maximum at

$$(\bar{\kappa}_x, \bar{\kappa}_y) = \left(-\frac{m'_{10}}{m''_{00}}, -\frac{m'_{01}}{m''_{00}}\right).$$
(2.6.25)

The distribution (2.6.22) itself is not in general symmetrical. However, when the spectrum is narrow, R is appreciable only in the neighbourhood of  $(\bar{\kappa}_x, \bar{\kappa}_y)$ , giving

$$p(\kappa_x,\kappa_y)_{\xi_1} = \frac{1}{\pi m \Delta_3^{\frac{1}{2}}} \frac{(\overline{\kappa}_x^2 + \overline{\kappa}_y^2)^{\frac{1}{2}}}{R^2}, \qquad (2.6.26)$$

approximately. The curves of constant probability are then the ellipses R = constant. The major axis of each ellipse makes an angle  $\varpi$  with the x axis given by

$$\tan 2\varpi = \frac{2M_{12}}{M_{11} - M_{22}}.$$
 (2.6.27)

Since  $p(\kappa_x, \kappa_y)_{\xi_1}$  is proportional to  $R^{-2}$ , it may be shown that the fraction of the distribution lying *outside* the ellipse is proportional to  $R^{-1}$ . At the centre, R = M. Therefore the ellipse enclosing just half the distribution is

$$R = 2M. \tag{2.6.28}$$

The semi-axes of this ellipse are of length

$$r_1, r_2 = \left[\frac{2M}{(M_{11} + M_{22}) \pm \sqrt{\{(M_{11} - M_{22})^2 + 4M_{12}^2\}}}\right]^{\frac{1}{2}}.$$
 (2.6.29)

To interpret these results, let the *u* axis be taken so as to pass through the centroid  $(\bar{u}, \bar{v})$  of the energy spectrum. The spectrum being narrow, we may expand in a Taylor series about  $(\bar{u}, 0)$  thus:

$$m_{00}'' = \sigma^{2} \mu_{00} + \frac{1}{2} \left( \mu_{20} \frac{\partial^{2}}{\partial u^{2}} + \mu_{02} \frac{\partial^{2}}{\partial v^{2}} \right) \sigma^{2} + \dots,$$
  

$$m_{10}' = u \sigma \mu_{00} + \frac{1}{2} \left( \mu_{20} \frac{\partial^{2}}{\partial u^{2}} + \mu_{02} \frac{\partial^{2}}{\partial v^{2}} \right) (u\sigma) + \dots,$$
  

$$m_{01}' = \mu_{11} \frac{\partial^{2}}{\partial u \partial v} (v\sigma) + \dots,$$
  
(2.6.30)

Vol. 249. A.

where  $u = \overline{u}$ . Making use of (1.6.9) we obtain

$$M_{11} = \frac{\mu_{02}}{\mu_{20}\mu_{02} - \mu_{11}^2} \frac{(\sigma/u)^2}{(\sigma/u - \partial\sigma/\partial u)^2},$$

$$M_{12} = \frac{\mu_{11}}{\mu_{20}\mu_{02} - \mu_{11}^2} \frac{\sigma/u}{\sigma/u - \partial\sigma/\partial u},$$

$$M_{22} = \frac{\mu_{20}}{\mu_{20}\mu_{02} - \mu_{11}^2},$$

$$\Delta_3 = \mu_{00}(\mu_{20}\mu_{02} - \mu_{11}^2) u^2(\sigma/u - \partial\sigma/\partial u)^2.$$

$$(\overline{\kappa}_x, \overline{\kappa}_y) = \left(-\frac{u}{\sigma}, 0\right),$$
(2.6.32)

Hence

$$\tan 2\varpi = \frac{2\mu_{11}\sigma/u(\sigma/u - \partial\sigma/\partial u)}{\mu_{20}(\sigma/u - \partial\sigma/\partial u)^2 - \mu_{02}(\sigma/u)^2},$$
(2.6.33)

and

showing that the centre of the distribution is the inverse of the phase velocity, and that  $\tan 2\omega$ , like  $\tan 2\beta$ , is proportional to  $\mu_{11}$  (cf. equation (1.6.14)). When the spectrum is symmetrical the semi-axes of the distribution are given by

$$r_{1} = \left(\frac{\mu_{20}}{\mu_{00}}\right)^{\frac{1}{2}} \left| \frac{\sigma/u - \partial\sigma/\partial u}{\sigma^{2}/u} \right|, \quad r_{2} = \left(\frac{\mu_{02}}{\mu_{00}}\right)^{\frac{1}{2}} \frac{1}{\sigma}, \quad (2.6.34)$$

$$r_{1} = \left| \nu \overline{\kappa} (1 - \Gamma \overline{\kappa}) \right|, \quad r_{2} = \left| \gamma \overline{\kappa} \right|, \quad (2 \cdot 6 \cdot 35)$$

where  $\bar{\kappa}_{,} = -u/\sigma_{,}$  is the mean reciprocal velocity;  $\Gamma_{,} = -\partial\sigma/\partial u_{,}$  is the corresponding group velocity;  $\nu = (\mu_{20}/\mu_{00}u^2)^{\frac{1}{2}}$  and  $\gamma = (\mu_{02}/\mu_{00}u^2)^{\frac{1}{2}}$ . This shows that  $r_2/\bar{\kappa}$ , which represents the width of the distribution perpendicular to the principal direction of the waves, depends only on the long-crestedness  $\gamma^{-1}$  and is proportional to  $\gamma$ . On the other hand,  $r_2/\bar{\kappa}$ , which represents the width of the distribution parallel to the principal direction, depends not only on the r.m.s. width of the spectrum, represented by  $\nu$ , but also on the dispersive properties of the medium. For gravity waves in deep water  $\Gamma\bar{\kappa} = \frac{1}{2}$  and so

$$r_1 = \frac{1}{2} \left| \nu \overline{\kappa} \right|. \tag{2.6.36}$$

### 2.7. The velocities of specular points

A specular point on the surface is defined as a point where the two components of the gradient take given values. Such points would be indicated to a distant observer as the points where light was reflected from a distant source. In  $\S 2 \cdot 4$  we deduced the mean density of such points per unit area; let us now consider the statistical distribution of their velocities.

If (x, y) are the co-ordinates at time t of a point whose components of gradient

$$\frac{\partial \zeta}{\partial x}, \frac{\partial \zeta}{\partial y} = \xi_2, \xi_3 \tag{2.7.1}$$

are fixed. At a subsequent time t + dt the point will have moved to a position (x + dx, y + dy), where  $(\partial t) = \partial^2 t = \partial^2 t = \partial^2 t$ 

$$0 = d\left(\frac{\partial\zeta}{\partial x}\right) = \frac{\partial^{2}\zeta}{\partial x^{2}} dx + \frac{\partial^{2}\zeta}{\partial x \partial y} dy + \frac{\partial^{2}\zeta}{\partial x \partial t} dt,$$

$$0 = d\left(\frac{\partial\zeta}{\partial y}\right) = \frac{\partial^{2}\zeta}{\partial x \partial y} dx + \frac{\partial^{2}\zeta}{\partial y^{2}} dy + \frac{\partial^{2}\zeta}{\partial y \partial t} dt.$$
(2.7.2)

# that is,

This content downloaded from 134.246.158.75 on Mon, 21 Oct 2013 05:45:01 AM All use subject to JSTOR Terms and Conditions

The ratios

$$\frac{\mathrm{d}x}{\mathrm{d}t}, \frac{\mathrm{d}y}{\mathrm{d}t} = c_x, c_y \tag{2.7.3}$$

are the required velocities. Writing

$$\frac{\partial^{2}\zeta}{\partial x^{2}}, \frac{\partial^{2}\zeta}{\partial x \partial y}, \frac{\partial^{2}\zeta}{\partial y^{2}} = \xi_{4}, \xi_{5}, \xi_{6}, 
\frac{\partial^{2}\zeta}{\partial x \partial t}, \frac{\partial^{2}\zeta}{\partial y \partial t} = \xi_{7}, \xi_{8}$$

$$\frac{\xi_{4}c_{x} + \xi_{4}c_{y} = -\xi_{7}, \\
\xi_{5}c_{x} + \xi_{6}c_{y} = -\xi_{8}.$$
(2.7.5)

in  $(2 \cdot 7 \cdot 2)$  we have

In other words

Since the velocities  $c_x$ ,  $c_y$  are given in terms of  $\xi_4$ , ...,  $\xi_8$ , we require first the joint distribution  $p(\xi_4, ..., \xi_8)_{\xi_2, \xi_3}$  of these quantities at points where  $\xi_2$ ,  $\xi_3$  take the given values.

Let dA denote any small area of the x, y plane, and P a neighbouring point. As usual,  $p(\xi_2, ..., \xi_8)$  will denote the ordinary distribution of  $\xi_2, ..., \xi_8$  at P. Now if  $\xi_2, \xi_3$  take the given values at some point in dA, and  $\xi_4, ..., \xi_8$  are fixed, then  $(\xi_2, \xi_3)$  at P lies within a certain region d $\Sigma$  of area  $d\Sigma = |\xi_4\xi_6 - \xi_5^2| dA$  (2.7.6)

(cf. §2·4). Hence the probability that  $\xi_2$ ,  $\xi_3$  take the given values in dA and that  $\xi_4, \ldots, \xi_8$  lie in ranges of width  $d\xi_4, \ldots, d\xi_8$  respectively is

$$p(\xi_2,\xi_3,...,\xi_8) | \xi_4\xi_6 - \xi_5^2 | dA d\xi_4 d\xi_5 \dots d\xi_8.$$
(2.7.7)

But the total probability of  $\xi_2$ ,  $\xi_3$  taking the given values in dA is

$$D_{\rm sp.} dA,$$
 (2.7.8)

where  $D_{sp.}$  is the density of specular points with gradient  $(\xi_2, \xi_3)$ . Therefore the probability that  $\xi_4 \dots \xi_8$  lie in their respective ranges given that there is a specular point in dA is the quotient of  $(2\cdot7\cdot7)$  and  $(2\cdot7\cdot8)$ , that is

$$\frac{p(\xi_2,...,\xi_8) |\xi_4\xi_6 - \xi_5^2| d\xi_4 \dots d\xi_8}{D_{\rm sp.}}.$$
(2.7.9)

 $p(\xi_4, \dots, \xi_8)_{\xi_2, \xi_3} = \frac{p(\xi_2, \dots, \xi_8) | \xi_4 \xi_6 - \xi_5^2 |}{D_{\rm sp.}}.$ (2.7.10)

Now the first derivatives  $\xi_2$ ,  $\xi_3$  are statistically independent of the second derivatives  $\xi_4 \dots \xi_8$ . Therefore  $h(\xi - \xi) - h(\xi - \xi) h(\xi - \xi)$  (9.7.11)

$$p(\xi_2, \dots, \xi_8) = p(\xi_2, \xi_3) \, p(\xi_4, \dots, \xi_8), \tag{2.7.11}$$

where  $p(\xi_2, \xi_3)$  is the ordinary distribution of  $\xi_2$ ,  $\xi_3$  given by  $(2 \cdot 1 \cdot 12)$  and  $p(\xi_4, ..., \xi_8)$  is the ordinary distribution of  $\xi_4, ..., \xi_8$ . The matrix of correlations for  $\xi_4, ..., \xi_8$  is

$$(\Xi_{ij}) = \begin{pmatrix} m_{40} & m_{31} & m_{22} & m'_{30} & m'_{21} \\ m_{31} & m_{22} & m_{13} & m'_{21} & m'_{12} \\ m_{22} & m_{13} & m_{04} & m'_{12} & m'_{03} \\ \hline m'_{30} & m'_{21} & m'_{12} & m'_{20} & m''_{11} \\ m'_{21} & m'_{12} & m'_{03} & m''_{11} & m''_{02} \end{pmatrix}, \qquad (2\cdot7\cdot12)$$

45-2

and so

$$p(\xi_4,...,\xi_8) = rac{1}{(2\pi)^{rac{5}{2}} \Delta_5^{rac{1}{2}}} \exp{\{-rac{1}{2} M_{ij} \xi_{i+3} \xi_{j+3}\}},$$
 (2.7.13)

where  $(M_{ij})$  is the inverse matrix to  $(\Xi_{ij})$  and where

$$\Delta_5 = |\Xi_{ij}|. \tag{2.7.14}$$

Substituting in  $(2 \cdot 7 \cdot 10)$  we have then

$$p(\xi_4, \dots, \xi_8)_{\xi_2, \xi_3} = \frac{1}{4(2\pi)^{\frac{7}{2}} (\Delta_2 \Delta_5)^{\frac{1}{2}} D_{\text{ma.}}} |\xi_4 \xi_6 - \xi_5^2| \exp\{-\frac{1}{2} M_{ij} \xi_{i+3} \xi_{j+3}\}, \quad (2.7.15)$$

where  $D_{\text{ma.}}$  is given by  $(2 \cdot 4 \cdot 51)$ .

The distribution of the velocities  $c_x$ ,  $c_y$  is now obtained from the relations (2.7.5). If we write also  $c_1, c_2, c_3 = \xi_4, \xi_5, \xi_6$ (2.7.16)

and transform to the variables  $c_x$ ,  $c_y$ ,  $c_1$ ,  $c_2$ ,  $c_3$  we have

$$\frac{\partial(\xi_4, \dots, \xi_8)}{\partial(c_x, c_y, c_1, c_2, c_3)} = \xi_4 \xi_6 - \xi_5^2 = c_1 c_3 - c_2^2.$$
(2.7.17)

Hence 
$$p(c_x, c_y, c_1, c_2, c_3)_{\xi_2, \xi_3} = \frac{1}{4(2\pi)^{\frac{7}{2}} (\Delta_2 \Delta_5)^{\frac{1}{2}} D_{\text{ma.}}} (c_1 c_3 - c_2^2)^2 \exp\{-\frac{1}{2} N_{ij} c_i c_j\},$$
 (2.7.18)

where  $(N_{ij})$  is the  $(3 \times 3)$  matrix whose elements are

On eliminating  $c_1, c_2, c_3$  by integration between  $\pm \infty$  we have

$$p(c_x, c_y)_{\xi_2, \xi_3} = \frac{1}{4(2\pi)^{\frac{7}{2}} (\Delta_2 \Delta_5)^{\frac{1}{2}} D_{\text{ma.}}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (c_1 c_3 - c_2^2)^2 \exp\left\{-\frac{1}{2} N_{ij} c_i c_j\right\} dc_1 dc_2 dc_3. \quad (2.7.20)$$

The matrix  $(N_{ij})$  is positive-definite. For, if any real values of  $c_1$ ,  $c_2$ ,  $c_3$ , not all zero existed which made the quadratic form  $N_{ij}c_ic_j$  zero or negative, a corresponding set of values of  $\xi_4 \dots \xi_8$  could be found from (2.7.5) which made  $M_{ij}\xi_{i+3}\xi_{j+3}$  zero or negative. But this is impossible, since  $(M_{ij})$  is positive-definite. Therefore  $(N_{ij})$  is positive-definite. Therefore by a real linear transformation of variables  $c_1, c_2, c_3 \rightarrow \eta_1, \eta_2, \eta_3$  we have

$$\begin{array}{c} N_{ij}c_ic_j = \eta_1^2 + \eta_2^2 + \eta_3^2, \\ c_1c_3 - c_2^2 = \mathbf{l}_1\eta_1^2 + \mathbf{l}_2\eta_2^2 + \mathbf{l}_3\eta_3^2, \end{array}$$

$$(2.7.21)$$

where  $\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3$  are roots of the equation

$$|\sigma_{ij} - \mathbf{1}N_{ij}| = 0. \tag{2.7.22}$$

As in  $\S 2 \cdot 4$  we have then

$$p(c_x, c_y)_{\xi_2, \xi_3} = \frac{1}{4(2\pi)^{\frac{7}{2}} (\Delta_2 \Delta_5 |N_{ij}|)^{\frac{1}{2}} D_{\text{ma.}}} \mathbf{I}(\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3), \qquad (2.7.23)$$

where

$$\mathbf{I}(\mathbf{l}_1, \mathbf{l}_2, \mathbf{l}_3) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (\mathbf{l}_1 \eta_1^2 + \mathbf{l}_2 \eta_2^2 + \mathbf{l}_3 \eta_3^2)^2 \exp\left\{-\frac{1}{2}(\eta_1^2 + \eta_2^2 + \eta_3^2)\right\} \mathrm{d}\eta_1 \mathrm{d}\eta_2 \mathrm{d}\eta_3. \quad (2 \cdot 7 \cdot 24)$$

This integral is much easier to evaluate than the similar integral  $I(l_1, l_2, l_3)$  of equation  $(2\cdot 4\cdot 36)$  on account of the factor in the integrand being squared. In fact we have immediately

$$\begin{split} \mathbf{I}(\mathbf{l}_1,\mathbf{l}_2,\mathbf{l}_3) &= (2\pi)^{\frac{3}{2}} \big[ \mathbf{3}(\mathbf{l}_1^2 + \mathbf{l}_2^2 + \mathbf{l}_3^2) + 2(\mathbf{l}_2\mathbf{l}_3 + \mathbf{l}_3\mathbf{l} + \mathbf{l}_1\mathbf{l}_2) \big]. \\ &= (2\pi)^{\frac{3}{2}} \big[ \mathbf{3}(\mathbf{l}_1 + \mathbf{l}_2 + \mathbf{l}_3)^2 - \mathbf{4}(\mathbf{l}_2\mathbf{l}_3 + \mathbf{l}_3\mathbf{l}_1 + \mathbf{l}_1\mathbf{l}_2) \big]. \end{split} \tag{2.7.25}$$

Therefore

$$p(c_x, c_y)_{\xi_2, \xi_3} = \frac{1}{16\pi^2 (\Delta_2 \Delta_5 |N_{ij}|)^{\frac{1}{2}} D_{\text{ma.}}} [3(\sum_i \mathbf{l}_i)^2 - 4\sum_{i\neq j} \mathbf{l}_i \mathbf{l}_j].$$
(2.7.26)

Equation (2.7.22) on expansion becomes

$$N\mathbf{l}^{3} - \left(\frac{1}{2}n_{13} - n_{22} + \frac{1}{2}n_{31}\right)\mathbf{l}^{2} + \left(\frac{1}{2}N_{13} - \frac{1}{4}N_{22} + \frac{1}{2}N_{31}\right)\mathbf{l} - \frac{1}{4} = 0, \qquad (2.7.27)$$

where

$$N = |N_{ij}| \tag{2.7.28}$$

and

$$n_{13} = N_{21}N_{32} - N_{22}N_{31} = n_{31}, \qquad n_{22} = N_{11}N_{33} - N_{22}^2. \tag{2.7.29}$$

Therefore

$$\sum_{i} \mathbf{1}_{i} = \frac{n_{13} - n_{22}}{N}, \qquad \sum_{i+j} \mathbf{1}_{i} \mathbf{1}_{j} = \frac{N_{13} - \frac{1}{4}N_{22}}{N}, \qquad (2.7.30)$$

and so finally

$$p(c_x, c_y)_{\xi_1, \xi_2} = \frac{1}{16\pi^2 (\Delta_2 \Delta_5)^{\frac{1}{2}} D_{\text{ma.}}} \frac{3(n_{13} - n_{22})^2 + (N_{22} - 4N_{13}) N}{N^{\frac{5}{2}}}.$$
 (2.7.31)

It will be seen that in general the quantities  $N_{ij}$ ,  $n_{ij}$  and N which occur in this expression are polynomials in  $c_x$ ,  $c_y$  of degree 2, 4 and 6 respectively.

As before, we may study this distribution in the special case when the energy spectrum is narrow and has symmetry about the principal direction. Taking the u axis along the line of symmetry we have  $m_{pq} = 0$  whenever q is odd, and so

$$(\Xi_{ij}) = \begin{pmatrix} m_{40} & 0 & m_{22} & m'_{30} & 0 \\ 0 & m_{22} & 0 & 0 & m'_{12} \\ m_{22} & 0 & m_{04} & m'_{12} & 0 \\ m'_{30} & 0 & m'_{12} & m''_{20} & 0 \\ 0 & m'_{12} & 0 & 0 & m''_{02} \end{pmatrix}.$$

$$(2.7.32)$$

The reciprocal matrix  $(M_{ii})$  is given by

$$(M_{ij}) = \begin{pmatrix} A_{11} & 0 & A_{12} & A_{13} & 0 \\ 0 & B_{11} & 0 & 0 & B_{12} \\ A_{21} & 0 & A_{22} & A_{23} & 0 \\ A_{31} & 0 & A_{32} & A_{33} & 0 \\ 0 & B_{21} & 0 & 0 & B_{22} \end{pmatrix},$$
(2.7.33)

(2.7.34)

 $(A_{ij}) = \begin{pmatrix} m_{40} & m_{22} & m'_{30} \\ m_{22} & m_{04} & m'_{12} \\ m'_{30} & m'_{12} & m''_{20} \end{pmatrix}^{-1}, \qquad (B_{ij}) = \begin{pmatrix} m_{22} & m'_{12} \\ m'_{12} & m''_{02} \end{pmatrix}^{-1}.$ 

Since the spectrum is narrow, each coefficient may be expanded in a Taylor series about the centroid  $(\bar{u}, 0)$ . Thus

$$\begin{split} m_{40} &= \mu_{00} u^{4} \qquad \qquad + \frac{1}{2} \mu_{20} \frac{\partial^{2}}{\partial u^{2}} (u^{4}) + \dots, \\ m_{22} &= \mu_{02} u^{2} + \mu_{12} \frac{\partial}{\partial u} (u^{2}) \qquad + \frac{1}{2} \mu_{22} \frac{\partial^{2}}{\partial u^{2}} (u^{2}), \\ m_{04} &= \mu_{04}, \\ m'_{30} &= \mu_{00} u^{3} \sigma \qquad \qquad + \frac{1}{2} \left( \mu_{20} \frac{\partial^{2}}{\partial u^{2}} + \mu_{02} \frac{\partial^{2}}{\partial v^{2}} \right) (u^{3} \sigma) + \dots, \\ m'_{12} &= \mu_{02} u \sigma + \mu_{12} \frac{\partial}{\partial u} (u \sigma) + \frac{1}{2} \left( \mu_{22} \frac{\partial^{2}}{\partial u^{2}} + \mu_{04} \frac{\partial^{2}}{\partial v^{2}} \right) (u \sigma) + \dots, \\ m''_{20} &= \mu_{00} u^{2} \sigma^{2} \qquad \qquad + \frac{1}{2} \left( \mu_{20} \frac{\partial^{2}}{\partial u^{2}} + \mu_{02} \frac{\partial^{2}}{\partial v^{2}} \right) (u^{2} \sigma^{2}) + \dots, \\ m''_{02} &= \mu_{02} \sigma^{2} + \mu_{12} \frac{\partial}{\partial u} (\sigma^{2}) + \frac{1}{2} \left( \mu_{22} \frac{\partial^{2}}{\partial u^{2}} + \mu_{04} \frac{\partial^{2}}{\partial v^{2}} \right) (\sigma^{2}) + \dots, \end{split}$$

where  $u = \overline{u}$  and  $\sigma = \sigma(\overline{u}, 0)$ . Thus we find for the determinants of  $(A_{ij})^{-1}$  and  $(B_{ij})^{-1}$ 

$$\begin{array}{l} |A_{ij}|^{-1} = \delta_3 u^6 (\sigma/u - \partial \sigma/\partial u)^2, \\ |B_{ij}|^{-1} = \delta_2 u^2 (\sigma/u - \partial \sigma/\partial u)^2, \end{array}$$

$$(2.7.36)$$

$$\delta_{3} = \begin{vmatrix} \mu_{00} & 0 & \mu_{02} \\ 0 & \mu_{20} & \mu_{12} \\ \mu_{02} & \mu_{12} & \mu_{04} \end{vmatrix}, \qquad \delta_{2} = \begin{vmatrix} \mu_{02} & \mu_{12} \\ \mu_{12} & \mu_{22} \end{vmatrix}.$$
(2.7.37)

(Each zero term in  $\delta_3$  could be replaced by  $\mu_{10}$ .) Further, on evaluating  $(A_{ij})$  and  $(B_{ij})$  and substituting in  $(2\cdot7\cdot19)$  we find eventually

where  $\alpha_{ij}$  and  $\beta_{ij}$  are the (i, j)th elements of the reciprocal matrices of  $\delta_3$  and  $\delta_2$  respectively, and  $\alpha_1 + \frac{\sigma}{2}$ 

$$q_1, q_2 = \frac{c_x + \sigma/u}{\sigma/u - \partial\sigma/\partial u}, \frac{c_y}{\sigma/u - \partial\sigma/\partial u}.$$
 (2.7.39)

Thus  $q_1$ ,  $q_2$  are non-dimensional quantities proportional to the departures of  $c_x$ ,  $c_y$  from their mean values  $(-\sigma/u, 0)$ . In deriving  $(2\cdot7\cdot38)$  it has been assumed that  $q_1$  is of the same order of magnitude as  $\gamma$ ,  $= (\mu_{02}/u^2\mu_{00})^{\frac{1}{2}}$ , but that  $q_2$  is of order 1; this makes the matrix  $(N_{ij})$  more homogeneous. Before proceeding further we may make the additional restrictions

$$\mu_{12} = 0, \quad \mu_{22} = \mu_{20} \mu_{02} / \mu_{00}, \quad (2.7.40)$$

and we may write  $\mu_{20} = \nu^2 u^2 \mu_{00}, \quad \mu_{02} = \gamma^2 u^2 \mu_{00}, \quad \mu_{04} = a^2 \gamma^4 u^4 \mu_{00},$  (2.7.41)

#### This content downloaded from 134.246.158.75 on Mon, 21 Oct 2013 05:45:01 AM All use subject to JSTOR Terms and Conditions

where v, y and a have the same meanings as before, namely,  $v^{-1}$  is a measure of the average length of a group of waves,  $\gamma^{-1}$  is the long-crestedness and a is the peakedness of the spectrum in the v direction. Then  $(2 \cdot 7 \cdot 38)$  reduces to

$$(N_{ij}) = \frac{1}{u^4 \mu_{00}} \begin{pmatrix} (\xi^2 + d^2 + 1) & \xi \eta / \gamma & -d^2 / \gamma^2 \\ \xi \eta / \gamma & (\xi^2 + \eta^2 + 1) / \gamma^2 & \xi \eta / \gamma^3 \\ -d^2 / \gamma^2 & \xi \eta / \gamma^3 & (\xi^2 + \eta^2) / \gamma^4 \end{pmatrix}, \qquad (2.7.42)$$

where

 $\xi = q_1/\nu, \quad \eta = \gamma q_2/\nu, \quad d^2 = 1/(a^2 - 1).$ (2.7.43)

Clearly the (i, j)th term of  $(N_{ij})$  is of order  $1/\gamma^{i+j}$ . Therefore

$$\mathbf{l}_{1}, \mathbf{l}_{2}, \mathbf{l}_{3} = \gamma^{2} u^{4} \mu_{00} \left( \mathbf{l}_{1}', \mathbf{l}_{2}', \mathbf{l}_{3}' \right)$$
(2.7.44)

where  $l'_1$ ,  $l'_2$ ,  $l'_3$  are the corresponding roots of the equation

$$|\sigma_{ij} - \mathbf{l}' N_{ij}'| = 0 \tag{2.7.45}$$

and  $(N'_{ii})$  is the non-dimensional matrix

$$(N'_{ij}) = \begin{pmatrix} \xi^2 + d^2 + 1 & \xi\eta & -d^2 \\ \xi\eta & \xi^2 + \eta^2 + 1 & \xi\eta \\ -d^2 & \xi\eta & \eta^2 + d^2 \end{pmatrix}.$$
 (2.7.46)

Also

 $\mid N_{ij} \mid = \mid N_{ij}' \mid / (\gamma^{6} u^{12} \mu_{00}^{3}).$ (2.7.47)

So from equation  $(2 \cdot 7 \cdot 26)$ 

$$p(c_x, c_y)_{\xi_2, \xi_3} = \frac{\gamma^7 u^{14} \mu_{00}^2}{16\pi^2 (\Delta_2 \Delta_5)^{\frac{1}{2}} D_{\text{ma.}}} \frac{3(\sum_i \mathbf{l}'_i)^2 - 4\sum_{i \neq j} \mathbf{l}'_i \mathbf{l}'_j}{|N'_{ij}|^{\frac{1}{2}}}.$$
(2.7.48)

It is convenient to state the solution in terms of the non-dimensional variables

$$\xi, \eta = \frac{c_x + \sigma/u}{\nu(\sigma/u - \partial\sigma/\partial u)}, \frac{\gamma c_y}{\nu(\sigma/u - \partial\sigma/\partial u)}, \qquad (2.7.49)$$

$$p(\xi,\eta)_{\xi_2,\xi_3} = \frac{\nu^2 (\sigma/u - \partial \sigma/\partial u)^2}{\gamma} p(c_x,c_y)_{\xi_2,\xi_3}.$$
(2.7.50)

First, on expanding the left-hand side of  $(2 \cdot 7 \cdot 45)$  we have

$$N'\mathbf{l}'^{3} + \eta^{2}\mathbf{l}'^{2} - \left[\frac{1}{4}(\xi^{2} + \eta^{2} + 1) + d^{2}\right]\mathbf{l}' - \frac{1}{4} = 0, \qquad (2.7.51)$$

(2.7.55)

where

giving

so that

$$N' = |N'_{ij}| = (\xi^2 + \eta^2 + 1) \eta^2 + \{(\xi + \eta)^2 + 1\} \{(\xi - \eta)^2 + 1\} d^2, \qquad (2.7.52)$$
  
$$\sum_{i} \mathbf{l}'_i = -\eta^2 / N', \qquad \sum_{i+j} \mathbf{l}'_i \mathbf{l}'_j = -\left[\frac{1}{4}(\xi^2 + \eta^2 + 1) + d^2\right] / N'. \qquad (2.7.53)$$

$$\begin{split} \Delta_2 &= \gamma^2 u^4 \mu_{00}^2, \qquad \Delta_5 = |A_{ij}|^{-1} |B_{ij}|^{-1} = \delta_2 \delta_3 u^8 (\sigma/u - \partial \sigma/\partial u)^4, \qquad (2 \cdot 7 \cdot 54) \\ \delta_2 \delta_3 &= \mu_{20} \mu_{22} \mu_{02} (\mu_{00} \mu_{04} - \mu_{02}^2) = \nu^4 \gamma^8 u^{12} \mu_{00}^5 (a^2 - 1). \qquad (2 \cdot 7 \cdot 55) \end{split}$$

 $D_{\rm ma.} = \gamma u^2 C(a)$ . Thirdly from  $(2 \cdot 4 \cdot 61)$ (2.7.56)

#### This content downloaded from 134.246.158.75 on Mon, 21 Oct 2013 05:45:01 AM All use subject to JSTOR Terms and Conditions





FIGURE 12. The probability distribution of the velocities of specular points, for a narrow spectrum

Therefore, altogether we have

$$p(\xi,\eta)_{\xi_2,\xi_3} = \frac{d}{16\pi^2 C(a)} \frac{3\eta^4 + \left[(\xi^2 + \eta^2 + 1) + 4d^2\right]N'}{N'^{\frac{5}{2}}}, \qquad (2.7.57)$$

where C(a) is given by  $(2 \cdot 4 \cdot 62)$  and N' by  $(2 \cdot 7 \cdot 52)$ .

Two special cases are of interest. Suppose first that the surface consists of two systems of long-crested waves, intersecting at a small angle  $2\gamma$ . As we saw in §2.4, this corresponds to the limiting case when  $a \rightarrow 1$  and  $d \rightarrow \infty$ . Equation (2.7.57) then becomes

$$p(\xi,\eta)_{\xi_2,\xi_3} = \frac{1}{2} \frac{1}{\{(\xi+\eta)^2+1\}^{\frac{3}{2}} \{(\xi-\eta)^2+1\}^{\frac{3}{2}}}.$$
(2.7.58)

The distribution is shown in figure 12 a. There are two ridges of high probability, in the directions  $\xi = \pm \eta$ , that is  $c_x + \sigma/u = \pm \gamma c_u$ , (2.7.59)

or when the vector difference between the specular velocity  $(c_x, c_y)$  and the mean velocity  $(-\sigma/u, 0)$  is in the direction of the crests of one of the two wave systems.

This particular case may also be derived quite simply as follows. We have seen that the velocities of specular points have the same probability distribution as the velocities of the maxima only. Now with two intersecting systems of long-crested waves, the maxima occur at the points of intersection of the crests of the two systems, and at no other points (see

VOL. 249. A.

### M. S. LONGUET-HIGGINS ON THE

figure 9*a*). If the point of intersection of two crests has components of velocity  $c_x$ ,  $c_y$  parallel and perpendicular to the mean direction, then the rates of advance of the crests in the two systems of waves are

$$c_1^{(1)} = c_x \cos \gamma + c_y \sin \gamma, \quad c_1^{(2)} = c_x \cos \gamma - c_y \sin \gamma, \quad (2 \cdot 7 \cdot 60)$$

where  $2\gamma$  is the angle between the two wave systems. But the distribution of  $c_1$  for a longcrested system of waves was found in § 2.5 to be

$$p(c_1) = \frac{1}{2} \frac{\nu^2 (\sigma/u - \partial \sigma/\partial u)^2}{\left[ (c_1 + \sigma/u)^2 + \nu^2 (\sigma/u - \partial \sigma/\partial u)^2 \right]^{\frac{3}{2}}}, \qquad (2.7.61)$$

where u is measured in the direction of propagation. Since the two systems are independent,

$$p(c_1^{(1)}, c_1^{(2)}) = p(c_1^{(1)}) p(c_1^{(2)})$$
(2.7.62)

When the angle of separation  $2\gamma$  is small,  $\nu$  and u are effectively the same for the two systems and for the combined system. Thus

$$c_1^{(1)} + \sigma/u = (c_x + \sigma/u) + \gamma c_y = (\xi + \eta) \nu(\sigma/u - \partial\sigma/\partial u),$$

$$c_1^{(2)} + \sigma/u = (c_x + \sigma/u) - \gamma c_y = (\xi - \eta) \nu(\sigma/u - \partial\sigma/\partial u).$$

$$(2.7.63)$$

$$p(\xi,\eta)_{\xi_2,\xi_3} = \left| \frac{\partial(c_1^{(1)}, c_1^{(2)})}{\partial(\xi,\eta)} \right| p(c_1^{(1)}, c_1^{(2)})_{\xi_2,\xi_3}, \tag{2.7.64}$$

370

$$p(\xi,\eta)_{\xi_2,\xi_3} = 2\nu^2 (\sigma/u - \partial\sigma/\partial u)^2 p(c_1^{(1)}) p(c_1^{(2)}), \qquad (2.7.65)$$

from which (2.7.58) follows.

A second case of interest is that of infinite peakedness:  $a \to \infty$  and  $d \to 0$ . For large values of a,  $(2 \cdot 7 \cdot 57)$  becomes  $1 \quad 3n^2 + (\xi^2 + n^2 + 1)^2$ 

$$p(\xi,\eta)_{\xi_2,\xi_3} = \frac{1}{4a^2} \frac{3\eta^2 + (\xi^2 + \eta^2 + 1)^2}{\eta^3 (\xi^2 + \eta^2 + 1)^{\frac{5}{2}}}.$$
(2.7.66)

This distribution is shown in figure 12 c. There is only one ridge of high probability, namely, that in the principal direction of the waves. The expression is valid only asymptotically, as is shown by the presence of the factor  $1/4a^2$  and the fact that  $\iint p(\xi, \eta)_{\xi_2, \xi_3} d\xi d\eta$  diverges.

An intermediate case,  $a^2 = 3$ ,  $d^2 = \frac{1}{2}$ , is shown in figure 12 *b*. This corresponds to a distribution of energy distributed normally with regard to direction, over a narrow range.

### 2.8. Properties of the envelope: the number of waves in a group

We shall now consider briefly some statistical properties of the envelope of the wave surface, as defined in §1.5. The envelope function  $\rho$  is essentially different from the surface elevation  $\zeta$ , in that  $\rho$  is always positive whereas  $\zeta$  has a mean value zero. Nevertheless, many of the properties of  $\rho$  will be seen to be analogous to corresponding properties of  $\zeta$ .

It is convenient to introduce the auxiliary variables

$$\begin{aligned} \xi_1 &= \rho \cos \phi = \sum_n c_n \cos \left\{ (u_n - \overline{u}) \, x + (v_n - \overline{v}) \, y + (\sigma_n - \overline{\sigma}) \, t + \epsilon_n \right\}, \\ \xi_2 &= \rho \sin \phi = \sum_n c_n \sin \left\{ (u_n - \overline{u}) \, x + (v_n - \overline{v}) \, y + (\sigma_n - \overline{\sigma}) \, t + \epsilon_n \right\}, \end{aligned}$$

$$(2.8.1)$$

which are the real and imaginary parts of the complex envelope function  $\rho e^{i\phi}$  (see equation (1.5.6)).  $\xi_1, \xi_2$  have the same form as  $\zeta$ , each being the sum of an infinite number of sinusoidal

components with random phase. In fact, the energy spectrum of  $\xi_1$ ,  $\xi_2$  is the same as that of  $\zeta$ , but with the origin moved to the mean wave-number  $(\overline{u}, \overline{v})$ . Thus  $\xi_1, \xi_2$  are normally distributed with mean value zero. Since

$$\overline{\xi_1^2} = \overline{\xi_2^2} = m_{00}, \quad \overline{\xi_1 \xi_2} = 0,$$
 (2.8.2)

we have

$$p(\xi_1,\xi_2) = \frac{1}{2\pi m_{00}} \exp\left\{-\frac{(\xi_1^2 + \xi_2^2)}{2m_{00}}\right\}.$$
(2.8.3)

We now transform back to the variables  $\rho$ ,  $\phi$ . From (2.8.1)

$$\xi_1^2 + \xi_2^2 = \rho^2, \quad \frac{\partial(\xi_1, \xi_2)}{\partial(\rho, \phi)} = \rho,$$
 (2.8.4)

$$p(\rho,\phi) = \frac{1}{2\pi m_{00}} \rho \exp\{-\rho^2/2m_{00}\}.$$
(2.8.5)

This is independent of the phase angle  $\phi$ . The distribution of  $\rho$  alone is found by integrating with respect to  $\phi$  from 0 to  $2\pi$ :

$$p(\rho) = \frac{1}{m_{00}} \rho \exp\{-\rho^2/2m_{00}\}, \qquad (2.8.6)$$

which is the well-known Rayleigh distribution.

etc.,

The joint distribution of  $\rho$ ,  $\phi$  and their first-order derivatives with respect to x, y may be found as follows. Let 17 17 25 25

$$\xi_3, \xi_4 = \frac{\partial \xi_1}{\partial x}, \frac{\partial \xi_1}{\partial y}; \quad \xi_5, \xi_6 = \frac{\partial \xi_2}{\partial x}, \frac{\partial \xi_2}{\partial y}. \tag{2.8.7}$$

The matrix of correlations for  $\xi_1, ..., \xi_6$  is

$$(\Xi_{ij}) = \begin{pmatrix} \mu_{00} & 0 & 0 & 0 & \mu_{10} & \mu_{01} \\ 0 & \mu_{00} & -\mu_{10} & -\mu_{01} & 0 & 0 \\ 0 & -\mu_{10} & \mu_{20} & \mu_{11} & 0 & 0 \\ 0 & -\mu_{01} & \mu_{11} & \mu_{02} & 0 & 0 \\ \mu_{10} & 0 & 0 & 0 & \mu_{20} & \mu_{11} \\ \mu_{01} & 0 & 0 & 0 & \mu_{11} & \mu_{02} \end{pmatrix},$$

$$(2.8.8)$$

where  $\mu_{pq}$  is the (p, q)th moment of E(u, v) about  $(\overline{u}, \overline{v})$ . But since  $(\overline{u}, \overline{v})$  is the centroid of E, the first-order moments  $\mu_{10}$ ,  $\mu_{01}$  vanish. Hence  $\xi_3$ ,  $\xi_4$ ,  $\xi_5$ ,  $\xi_6$  are independent of  $\xi_1$ ,  $\xi_2$ , and

$$p(\xi_1,...,\xi_6) = p(\xi_1,\xi_2) \, p(\xi_3,...,\xi_6), \tag{2.8.9}$$

where

$$p(\xi_3, \dots, \xi_6) = \frac{1}{(2\pi)^2 \,\delta} \exp\left\{-\left[\mu_{02}(\xi_3^2 + \xi_5^2) - 2\mu_{11}(\xi_3\xi_4 + \xi_5\xi_6) + \mu_{20}(\xi_4^2 + \xi_6^2)\right]/2\delta\right\}, \quad (2.8.10)$$

$$\delta = \begin{vmatrix} \mu_{20} & \mu_{11} \\ \mu_{11} & \mu_{02} \end{vmatrix}.$$
(2.8.11)

$$= \frac{\partial}{\partial x} (\rho \cos \phi) = \rho_x \cos \phi - \rho \phi_x \sin \phi, \qquad (2.8.12)$$

and we have written

$$\xi_3 = \frac{\partial}{\partial x} (\rho \cos \phi) = \rho_x \cos \phi - \rho \phi_x \sin \phi, \qquad (2.8.12)$$

we have

$$rac{\partial(\xi_1,\xi_2,\xi_3,\xi_4,\xi_5,\xi_6)}{\partial(
ho,\phi,
ho_x,\phi_x,
ho_y,\phi_y)}=-
ho^3,$$
 (2.8.13)

46-2

and hence

$$p(\rho, \phi, \rho_x, \phi_x, \rho_y, \phi_y) = \frac{1}{(2\pi)^3 m_{00} \delta} \rho^3 \exp\{-\rho^2/2m_{00}\}$$
  
 
$$\times \exp\{-(\mu_{02}\rho_x^2 - 2\mu_{11}\rho_x\rho_y + \mu_{20}\rho_y^2)/2\delta\} \exp\{-\rho^2\{\mu_{02}\phi_x^2 - 2\mu_{11}\phi_x\phi_y + \mu_{20}\phi_y^2\}/2\delta\}.$$
 (2.8.14)

From this distribution some immediate conclusions may be drawn. First, by integrating with respect to  $\phi$  (from 0 to  $2\pi$ ) and  $\phi_x$ ,  $\phi_y$  (from  $-\infty$  to  $\infty$ ) we obtain the joint distribution of  $\rho$ ,  $\rho_x$  and  $\rho_y$ . Thus (a, a) = h(a) h(a)

$$p(\rho, \rho_x, \rho_y) = p(\rho) p(\rho_x, \rho_y), \qquad (2.8.15)$$

where  $p(\rho)$  is given by (2.8.6) and

....

$$p(\rho_x, \rho_y) = \frac{1}{2\pi\delta^{\frac{1}{2}}} \exp\left\{-(\mu_{02}\rho_x^2 - 2\mu_{11}\rho_x\rho_y + \mu_{20}\rho_y^2)/2\delta\right\}.$$
(2.8.16)

This shows that  $\rho_x$ ,  $\rho_y$  are statistically independent of  $\rho$ , just as  $\partial \zeta / \partial x$ ,  $\partial \zeta / \partial y$  are independent of  $\zeta$ . Further, the distributon (2.8.16) is formally identical with (2.1.12), if the moments  $\mu_{bq}$  about the centroid are substituted for the moments  $m_{bq}$  about the origin. We deduce immediately that

(1) the steepest r.m.s. gradient of the envelope is in the principal direction of the envelope, and the gentlest r.m.s. gradient of the envelope is in the direction at right angles;

(2) the most probable direction of contours of the envelope is perpendicular to the principal direction of the envelope, and the least probable direction is parallel to the principal direction.

We see from (2.8.14) that the mean values of  $\phi_x$  and  $\phi_y$  are zero. It follows that the phase angle  $\phi$  of the envelope has zero secular increase in any horizontal direction. Now the phase angle of  $\zeta$  is the sum of the phase angles of the envelope and of the carrier wave. Hence the phase angle of  $\zeta$  increases at the same average rate as that of the carrier wave, in any horizontal direction. This property is the result of our having chosen the centroid  $(\bar{u}, \bar{v})$  of the energy distribution as the wave-number of the carrier wave  $(\S 1 \cdot 5)$ .

By integrating (2.8.14) with respect to  $\rho_x$ ,  $\rho_y$  and  $\phi$  we obtain

$$p(\rho,\phi_x,\phi_y) = \frac{1}{2\pi m_{00}\delta^{\frac{1}{2}}}\rho^3 \exp\left\{-\rho^2/2m_{00}\right\} \exp\left\{-\rho^2(\mu_{02}\phi_x^2 - 2\mu_{11}\phi_x\phi_y + \mu_{20}\phi_y^2)/2\delta\right\}.$$
 (2.8.17)

This shows that  $\phi_x$  and  $\phi_y$  are not statistically independent of  $\rho$ . In fact the standard deviation of  $(\phi_x, \phi_y)$  (defined as the square root of the mean value of  $(\phi_x^2 + \phi_y^2)$ ) is

$$(\overline{\phi_x^2 + \phi_y^2})^{\frac{1}{2}} = \frac{1}{\rho} \left(\mu_{20} + \mu_{02}\right)^{\frac{1}{2}}, \tag{2.8.18}$$

which is inversely proportional to  $\rho$ . Roughly, this means that the higher waves are more regular than the lower waves (cf. §2.10). The joint distribution of  $\phi_x$  and  $\phi_y$  alone is found from  $(2 \cdot 8 \cdot 17)$  to be

$$p(\phi_x, \phi_y) = \frac{1}{\pi} \frac{m_{00}/\delta^{\frac{1}{2}}}{\left[1 + (\mu_{02}\phi_x^2 - 2\mu_{11}\phi_x\phi_y + \mu_{20}\phi_y^2) m_{00}/\delta\right]^2}.$$
 (2.8.19)

It will be useful to consider also the statistical properties of the envelope of the curve in which the surface is intersected by a vertical plane in a direction  $\theta$ . When  $\theta = 0$ , x = x', the

distribution of  $\rho$ ,  $\phi$ ,  $\rho_x$ ,  $\phi_x$  may be found from (2.8.14) by integration with respect to  $\rho_y$  and  $\phi_y$ . On replacing  $\mu_{20}$  by  $\mu_2(\theta)$  and x by x' we have in the general case

$$p(\rho,\phi,\rho_{x'},\phi_{x'}) = \frac{1}{(2\pi)^2 m_{00}\mu_2} \rho^2 \exp\left\{-\rho^2/2m_{00}\right\} \exp\left\{-(\rho_{x'}^2 + \rho^2 \phi_{x'}^2)/2\mu_2\right\}.$$
 (2.8.20)

Alternatively, the distribution may be derived from first principles by the method used to obtain  $(2\cdot8\cdot14)$ . The joint distribution of  $\rho$  and  $\rho_{x'}$  is found by further integration with respect to  $\phi$  and  $\phi_{x'}$ :

$$p(\rho, \rho_{x'}) = \frac{1}{(2\pi)^{\frac{1}{2}} m_{00} \mu_2^{\frac{1}{2}}} \rho \exp\left\{-\rho^2 / 2m_{00}\right\} \exp\left\{-\rho_{x'}^2 / 2\mu_2\right\}.$$
(2.8.21)

Similarly the joint distribution of  $\rho$ ,  $\phi$  and  $\phi_{x'}$  is

$$p(\rho,\phi,\phi_{x'}) = \frac{1}{(2\pi)^{\frac{3}{2}} m_{00} \mu_2^{\frac{1}{2}}} \rho^2 \exp\left\{-\rho^2/2m_{00}\right\} \exp\left\{-\rho^2 \phi_{x'}^2/2\mu_2\right\}, \qquad (2\cdot 8\cdot 22)$$

and the distribution of  $\phi$  and  $\phi_{x'}$  is

$$p(\phi, \phi_{x'}) = \frac{(m_{00}/\mu_2)^{\frac{1}{2}}}{4\pi (1 + \phi_{x'}^2 m_{00}/\mu_2)^{\frac{3}{2}}}.$$
(2.8.23)

From these distributions one can state immediately some general conclusions for the onedimensional envelope analogous to those for the two-dimensional envelope of surface. Thus  $\rho_{x'}$ , but not  $\phi_{x'}$ , is independent of  $\rho$ ; the mean secular increase of  $\phi$  with x' is zero; the standard deviation of  $\phi_{x'}$  is inversely proportional to  $\rho$ .

When the spectrum is fairly narrow, the envelope follows closely the crests of the waves. In any particular plane section the waves will appear in groups, and a rough measure of the average length of a group is given by 2/N, where N is the average number of times per unit distance that the envelope crosses an arbitrary level  $\rho$ . Now by the argument of §2.2,

$$\mathbf{N}(\rho) = \int_{-\infty}^{\infty} p(\rho, \rho_{x'}) \left| \rho_{x'} \right| \mathrm{d}\rho_{x'}. \tag{2.8.24}$$

On substituting from  $(2 \cdot 8 \cdot 23)$  and carrying out the integration we have

$$\mathbf{N}(\rho) = \left(\frac{2}{\pi}\right)^{\frac{1}{2}} \frac{\mu_2^{\frac{1}{2}}}{m_{00}} \rho \exp\left\{-\rho^2/2m_{00}\right\}.$$
 (2.8.25)

For definiteness we may take the largest possible value of N, which occurs when  $\rho = m_{00}^{\frac{1}{2}}$ , giving

$$\mathbf{N} = \left(\frac{2}{e\pi}\right)^{\frac{1}{2}} \left(\frac{\mu_2}{m_{00}}\right)^{\frac{1}{2}}.$$
 (2.8.26)

Now by  $\S 1 \cdot 5 \mu_2(\theta)$  is greatest when  $\theta$  defines the principal direction. It follows that the average length of a group of waves is least in the principal direction and greatest in the direction at right angles.

A rough measure of the number of waves in each group is given by  $N_0/N$ , where  $N_0$  is the number of zero-crossings of  $\zeta$  in the direction  $\theta$  (see § 2.2). When  $\rho = m_{00}^{\frac{1}{2}}$  we have

$$\frac{N_0}{N} = \left(\frac{e}{2\pi}\right)^{\frac{1}{2}} \left(\frac{m_2}{\mu_2}\right)^{\frac{1}{2}} = \left(\frac{e}{2\pi}\right)^{\frac{1}{2}} \left[\frac{m_{20}\cos^2\theta + 2m_{11}\cos\theta\sin\theta + m_{02}\sin^2\theta}{\mu_{20}\cos^2\theta + 2\mu_{11}\cos\theta\sin\theta + \mu_{02}\sin^2\theta}\right]^{\frac{1}{2}}.$$
 (2.8.27)

In general this number will vary with the direction  $\theta$ , but it may also be constant. The condition for constancy is (2.8.28)

$$\mu_{20}:\mu_{11}:\mu_{02}=m_{20}:m_{11}:m_{02}.$$
(2.8.28)

When this condition is satisfied the number of waves in a group is independent of the direction.

If we write 
$$\nu'(\theta) = \frac{m_{00}}{m_1(\theta)} \left(\frac{\mu_2(\theta)}{m_{00}}\right)^{\frac{1}{2}},$$
 (2.8.29)

so that (for a narrow spectrum)

$$\nu'(\theta) \doteq \left(\frac{\mu_2(\theta)}{m_2(\theta)}\right)^{\frac{1}{2}} = \left(\frac{2\pi}{e}\right)^{\frac{1}{2}} \frac{\mathbf{N}}{N_0}, \qquad (2\cdot8\cdot30)$$

it is clear that  $\nu'$  is inversely proportional to the number of waves in a group. In particular, when the section is taken in the direction  $\theta \doteq 0$  we have

$$\nu'(0) = \left(\frac{\mu_{20}}{\overline{u}^2 m_{00}}\right)^{\frac{1}{2}} = \nu,$$
(2.8.31)

where  $\nu$  is the parameter defined in § 1.6. Now  $\theta = 0$  was taken there to be the mean direction, and also the principal direction. It follows that  $\nu$  is inversely proportional to the number of waves in a group corresponding to a vertical section taken in the principal direction.

### 2.9. The heights of maxima

Throughout this and the following section it will be assumed that the spectrum is narrow. We shall see that from the properties of the envelope one can then derive some interesting statistical properties that are otherwise difficult to obtain.

Consider first the distribution of the heights  $\xi$  of the crests. A crest may be defined as the locus of the maxima of all vertical sections of the surface parallel to the mean direction  $\overline{\theta}$ . Now when the spectrum is narrow, the waves will be long-crested and regular, and the crests will lie almost on the envelope. Further, the crests will be spaced at more or less equal intervals in the x, y plane. It follows that the distribution of the crest heights is practically the same as the distribution of the envelope function  $\rho$ . So from (2.8.6)

$$p(\xi) = \frac{\xi}{m_{00}} \exp\{-\xi^2/2m_{00}\}.$$
(2.9.1)

In other words,  $\xi$  has a Rayleigh distribution.

Consider, on the other hand, the distribution of the heights of the maxima. A maximum of the surface is simultaneously a maximum perpendicular and parallel to the mean direction. The distribution of maxima of the surface therefore approximates to the distribution of the maxima of the envelope of a section at right angles to the mean direction.

Now the distribution of the maxima of the envelope of a single random variable has been studied by Rice (1944, 1945). Making the simplifying assumption that  $\mu_1 = \mu_3 = 0$  (in our case  $\mu_1 = 0$  anyway), he obtains for the joint distribution of x' and the height R of a maximum

$$p(x',R) = \frac{1}{4\pi^{\frac{1}{2}}\mu_0^{\frac{3}{2}}} (a^2 - 1)^{\frac{3}{2}} z^{\frac{3}{2}} e^{-a^2 z^2} \sum_{n=0}^{\infty} \frac{A_n z^n}{(\frac{1}{2}n + \frac{3}{4})!}, \qquad (2.9.2)$$

where

$$a^2 = \frac{\mu_0 \mu_4}{\mu_2^2}, \quad z = \frac{R}{[2(a^2 - 1) \mu_0]^{\frac{1}{2}}},$$
 (2.9.3)

and

$$A_n = \sum_{m=0}^n \frac{\left(\frac{1}{2}\right) \left(\frac{3}{2}\right) \dots \left(m - \frac{1}{2}\right)}{m!} \left(n - m + 1\right) \left(\frac{3}{2} - \frac{1}{2}a^2\right)^m$$
(2.9.4)

(the term corresponding to m = 0 in  $(2 \cdot 9 \cdot 4)$  is n+1). To obtain the probability density of R alone we must normalize  $(2 \cdot 9 \cdot 2)$  by dividing by the number N of maxima per unit distance x'. N is found by integrating with respect to R from 0 to  $\infty$ . We have

$$N = \frac{1}{4(2\pi)^{\frac{1}{2}}} \left(\frac{\mu_2}{\mu_0}\right)^{\frac{1}{2}} \frac{(a^2 - 1)^2}{a^{\frac{5}{2}}} \sum_{n=0}^{\infty} \frac{(\frac{1}{2}n + \frac{1}{4})! A_n}{(\frac{1}{2}n + \frac{3}{4})! a^n}$$
(2.9.5)

(Rice 1945, p. 83).

This may be checked immediately by comparison with our previous work. For, since the maxima occur only at the crests of the waves, which are more or less evenly spaced at distance  $2\pi/\bar{u}$  apart, it follows that the mean density of maxima per unit area is  $N\bar{u}/2\pi$  approximately. On replacing  $(\mu_2/\mu_0)^{\frac{1}{2}}$  by  $\gamma\bar{u}$ , where  $\gamma^{-1}$  is the long-crestedness, we have

$$D_{\rm ma.} = \frac{N\overline{u}}{2\pi} = \frac{1}{4(2\pi)^{\frac{3}{2}}} \frac{(a^2 - 1)^2}{a^{\frac{5}{2}}} \sum_{n=0}^{\infty} \frac{(\frac{1}{2}n + \frac{1}{4})! A_n}{(\frac{1}{2}n + \frac{3}{4})! a^n} \gamma \overline{u}^2.$$
(2.9.6)

This expression should agree with  $(2\cdot3\cdot57)$ . Rice summed the series in  $(2\cdot9\cdot6)$  for  $a^2 = 3$  (the normal distribution) and found it to be about  $3\cdot97$ . With this value  $(2\cdot9\cdot6)$  becomes

$$D_{
m ma.} = 0.0638 \gamma \overline{u}^2, \qquad (2.9.7)$$

which is in agreement with the more accurate value  $D_{\text{ma.}} = 0.0639 \gamma \overline{u}^2$  given by (2.3.57) and table 1.

The distribution of the heights of maxima may be stated in terms of the non-dimensional parameter  $\eta = R/\mu_0^{\frac{1}{2}} = R/m_{00}^{\frac{1}{2}}$ . On dividing p(x', R) by  $N\mu_0^{-\frac{1}{2}}$ , we find

$$p(\eta) = \frac{1}{2^{\frac{1}{4}}(1-a^{-2})^{\frac{5}{4}}} \eta^{\frac{3}{2}} \exp\left\{-\eta^2/2(1-a^{-2})\right\} \sum_{n=0}^{\infty} \frac{A_n \left[\eta^2/2(a^2-1)\right]^{\frac{1}{2}n}}{\left(\frac{1}{2}n+\frac{3}{4}\right)!} / \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}n+\frac{1}{4}\right)!A_n}{\left(\frac{1}{2}n+\frac{3}{4}\right)!a^n}.$$
 (2.9.8)

This distribution has been computed for  $a^2 = 2, 3, 5, 9$ , and the curves are shown in figure 13.

When  $a^2 \doteq 1$  the above series becomes unsuitable for computation, but we may obtain a solution by an independent method as follows.  $a^2 = 1$  corresponds to two narrow bands of energy of slightly different frequency. These form beats, and the maxima of the envelope occur when the two wave bands are in phase. The height of the envelope R is then the sum of the amplitudes  $\rho_1$ ,  $\rho_2$  of the two wave trains at that point. But the amplitude of each wave train has a Rayleigh distribution:

$$p(\rho_1) = \frac{2\rho_1}{m_{00}} \exp\left\{-\rho_1^2/m_{00}\right\}, \qquad p(\rho_2) = \frac{2\rho_2}{m_{00}} \exp\left\{-\rho_2^2/m_{00}\right\}$$
(2.9.9)

(the mean energy for each wave train being  $\frac{1}{2}m_{00}$ ). The distribution of the sum of these is

$$p(R) = \int_0^R p(\rho_1) \, p(\rho_2) \, \mathrm{d}\rho_1, \qquad (2 \cdot 9 \cdot 10)$$

where  $\rho_2 = R - \rho_1$ . In terms of non-dimensional variables we have

$$p(\eta) = 2 \int_0^{\eta} \xi \,\mathrm{e}^{-\xi^2} \,(\eta - \xi) \,\mathrm{e}^{-(\eta - \xi)^2} \,\mathrm{d}\xi. \tag{2.9.11}$$

On evaluating the integral we have

$$p(\eta) = e^{-\frac{1}{2}\eta^2} [\eta e^{-\frac{1}{2}\eta^2} + (\eta^2 - 1) \int_0^{\eta} e^{-\frac{1}{2}t^2} dt], \qquad (2.9.12)$$

which is the distribution shown in figure 13 for  $a^2 = 1$ .

## M. S. LONGUET-HIGGINS ON THE

It can be seen independently that this is the appropriate distribution for two long-crested systems of waves intersecting at an angle. For the maxima of the combine system occur at the points of intersection of the crests of the two long-crested systems. The height of a maximum is the sum of the heights of the crests of the two systems. Consequently  $\eta$  is the sum of two variables each having a Rayleigh distribution.



FIGURE 13. The probability distribution of the heights of maxima, for a narrow spectrum  $(a^2 = 1, 2, 3, 5, 9)$ .

### 2.10. The intervals between successive zeros

Finally, let us consider the distribution of the intervals l between successive zeros of the surface, along a line drawn in an arbitrary direction  $\theta$ . An approximate expression for the distribution of intervals for a one-dimensional function has been derived by Rice (1945, § 3·4) after a series of approximations assuming that the spectrum is narrow. It will now be shown how the same distribution can be derived very simply using the properties of the phase angle  $\phi$ . Further, the distribution will be derived for a section of the surface in an arbitrary direction  $\theta$ , and for waves of any particular amplitude  $\rho$ .

For simplicity we may take initially  $\theta = \theta_p = 0$  and x' = x, and we may generalize to arbitrary values of  $\theta$  at a suitable stage. The equation of the curve at an arbitrary time, say t = 0, may be written

$$\zeta = \mathscr{R}\rho \,\mathrm{e}^{\mathrm{i}\chi},\tag{2.10.1}$$

$$\chi = \overline{u}x + \phi, \quad \chi_r = \overline{u} + \phi_r, \quad (2.10.2)$$

where

and  $\overline{u} = m_1/m_0$ . Like  $\phi$ ,  $\chi$  is a multivalued function of x, having branches separated by multiples of  $2\pi$ . Since  $\partial(\chi, \chi)$ 

$$rac{\partial(\chi,\chi_x)}{\partial(\phi,\phi_x)}=1, \hspace{1.5cm} (2{\cdot}10{\cdot}3)$$

we have from  $(2 \cdot 8 \cdot 23)$ 

$$p(\chi,\chi_x) = p(\phi,\phi_x) = \frac{(m_0/\mu_2)^{\frac{1}{2}}}{4\pi \{1 + (\chi_x - \overline{u})^2 m_0/\mu_2\}^{\frac{3}{2}}}.$$
 (2.10.4)

Now  $\zeta$  has a zero -crossing when and only when  $\chi = n\pi$ , where *n* is an integer. By the same reasoning as in §2.2, the probability of  $\chi$  taking the value  $2r\pi$  in (x, x+dx) is

$$H(\chi) \, \mathrm{d}x = \int_{-\infty}^{\infty} p(\chi, \chi_x) \, | \, \chi_x \, | \, \mathrm{d}x \, \mathrm{d}\chi_x = \frac{1}{2\pi} \left(\frac{m_2}{m_0}\right)^{\frac{1}{2}} \mathrm{d}x. \tag{2.10.5}$$

The probability of  $\chi$  taking the value  $(2r+1)\pi$  is the same. Therefore the total probability of a zero in (x, x+dx) is twice this value, or

$$\frac{1}{\pi} \left(\frac{m_2}{m_0}\right)^{\frac{1}{2}} \mathrm{d}x, \qquad (2.10.6)$$

in agreement with  $(2 \cdot 3 \cdot 5)$ .

Let l denote the interval between successive zeros. The average value  $\overline{l}$  of the distribution of l may be written down immediately; for it is simply the reciprocal of the average number of zeros per unit distance, i.e.

$$\bar{l} = \pi \left(\frac{m_0}{m_2}\right)^{\frac{1}{2}} = \frac{\pi}{\bar{u}} \left(1 + \nu^2\right)^{\frac{1}{2}},$$
(2.10.7)

where v is defined by  $(2 \cdot 8 \cdot 31)$ . When the spectrum is narrow (v is small) we have

$$\bar{l} = \pi/\bar{u},$$
 (2.10.8)

approximately.

Let us now consider the whole distribution of l, on the assumption that  $\nu$  is small. In the first place we may note that where  $\chi$  crosses any level  $n\pi$  it nearly always has an up-crossing. For the probability of a down-crossing ( $\chi_x < 0$ ) in the interval (x, x+dx) is given by

$$\int_{-\infty}^{0} p(\chi, \chi_x) |\chi_x| dx d\chi_x = \frac{1}{4\pi} \left(\frac{m_2}{m_0}\right)^{\frac{1}{2}} \left[1 - (1 + \nu^2)^{-\frac{1}{2}}\right] dx, \qquad (2.10.9)$$

and the proportion of down-crossings is therefore

$$\frac{1}{2} \left[ 1 - (1 + \nu^2)^{-\frac{1}{2}} \right] \doteq \frac{1}{4} \nu^2, \qquad (2 \cdot 10 \cdot 10)$$

which is negligible. If each crossing of  $2r\pi$  or  $(2r+1)\pi$  is an up-crossing, it follows that between any two successive zeros  $x_1$  and  $x_2$ ,  $\chi$  must increase by  $\pi$ . Hence

$$\pi = \chi(x_2) - \chi(x_1) = [l\chi_x + \frac{1}{2}l^2\chi_{xx} + \dots], \qquad (2.10.11)$$

where we have expanded in a Taylor series about  $x = x_1$ . It can be shown that  $\chi_{xx}$  is of order  $\nu^2$ , and so to our present order of approximation

$$l = \pi/\chi_x. \tag{2.10.12}$$

Now the distribution of  $\chi_x$ , at points where  $\chi$  takes a particular value, is given by

$$p(\chi_x)_{\chi} = \frac{p(\chi, \chi_x) |\chi_x|}{H(\chi)}, \qquad (2.10.13)$$

Vol. 249. A.

where  $H(\chi)$  is given by (2.10.5). That is to say

$$p(\chi_x)_{\chi} = \frac{1}{2} \left( \frac{m_0}{m_2} \right)^{\frac{1}{2}} \frac{|\chi_x| (m_0/\mu_2)^{\frac{1}{2}}}{[1 + (\chi_x - \overline{u})^2 m_0/\mu_2)]^{\frac{3}{2}}}.$$
 (2.10.14)

On substituting from  $(2 \cdot 10 \cdot 12)$  we have, when l > 0,

$$p(l) = \frac{1}{2\pi} \left(\frac{m_0}{m_2}\right)^{\frac{1}{2}} \frac{\overline{u}^2/\nu}{\left[l^2/\overline{l}^2 + (l/\overline{l}-1)^2/\nu^2\right]^{\frac{3}{2}}},$$
(2.10.15)

where  $\bar{l}$  is given by the approximate relation (2.10.8). Clearly p(l) is appreciable only when l differs from  $\bar{l}$  by an amount of order  $\nu$ . Writing

$$\xi = (l - \bar{l})/\bar{l}$$
 (2.10.16)

for the relative departure of l from its mean value, we have finally for the approximate distribution of  $\xi$  in the neighbourhood of the mean

$$p(\xi) = \frac{1}{2\nu(1+\xi^2/\nu^2)^{\frac{3}{2}}}.$$
(2.10.17)

This is similar to the approximate distribution found by Rice (1945, p. 63) by a rather longer method.

In the general case when the line drawn on the surface is in an arbitrary direction  $\theta$ ,  $\nu$  may be replaced by  $\nu'(\theta)$ . Thus we have in general

$$p(\xi) = \frac{1}{2\nu'(1+\xi^2/\nu'^2)^{\frac{3}{2}}}.$$
(2.10.18)

The second moment of this distribution is divergent, but a convenient measure of its spread is the width of the interquartile range, given by

$$\frac{2}{\sqrt{3}}\nu'.$$
 (2.10.19)

So we may say that the width of the distribution of  $\xi$  is inversely proportional to the average number of waves in a group. The width of the distribution of l is given by the above expression multiplied by  $\overline{l}$ , that is to say

$$\frac{2\pi}{\sqrt{3}} \frac{m_0^{\sharp} \mu_2^{\sharp}}{m_1^2}.$$
 (2.10.20)

To find the distribution of intervals l for waves of a given amplitude (say with amplitude lying between  $\rho$  and  $\rho + d\rho$ ), we may start from the distribution of  $(\phi, \phi_x)$  for values of  $\rho$  lying between these limits. This is given by

$$p_{\rho}(\phi,\phi_x) = \frac{p(\rho,\phi,\phi_x)}{p(\rho)} = \frac{1}{(2\pi)^{\frac{3}{2}}\mu_2^{\frac{1}{2}}}\rho\exp\left\{-\rho^2\phi_x^2/2\mu_2\right\}$$
(2·10·21)

from  $(2 \cdot 8 \cdot 6)$  and  $(2 \cdot 8 \cdot 22)$ . Hence

$$p_{\rho}(\chi,\chi_x) = \frac{1}{(2\pi)^{\frac{3}{2}} \mu_2^{\frac{1}{2}}} \rho \exp\left\{-\rho^2 (\chi_x - \overline{u})^2 / 2\mu_2\right\}.$$
(2.10.22)

On carrying out the same calculation with  $p_{\rho}(\chi, \chi_x)$  in place of  $p(\chi, \chi_x)$  we find, with  $\xi$  given by (2.10.16), that

$$p_{\rho}(\xi) = \frac{1}{(2\pi)^{\frac{1}{2}}\nu' m_{00}^{\frac{1}{2}}} \rho \exp\{-\rho^2 \xi^2 / 2\nu'^2 m_{00}\}.$$
(2.10.23)

This is a normal distribution for  $\xi$ , with mean value zero and standard deviation

$$\frac{\nu' m_{00}^{\frac{1}{2}}}{\rho}.$$
 (2.10.24)

The mean and standard deviation of l are equal to  $\bar{l}$  and  $\nu' m_{00}^{\frac{1}{2}} l/\rho$  respectively. Hence we may say that the expectation of l is independent of the height of the waves (to the present order of approximation) and also the width of the distribution of l is inversely proportional to the wave height. If we take this width as a measure of the irregularity of the waves as regards their intervals, we may also say that the lower the waves, the less regular are their intervals.

### PART III. A METHOD OF DETERMINING THE ENERGY SPECTRUM

In the two previous parts of this paper we have derived some statistical properties of a random surface in terms of its energy spectrum E(u, v). In this part we solve the converse problem: given the statistical properties, to find the energy spectrum.

The best method of determining E depends to some extent upon which properties can be measured most conveniently. We assume that it is possible to obtain the height  $\zeta(x')$  of the surface along a line in an arbitrary direction  $\theta$ . (In the case of the sea surface one may imagine the observations to be made by an aircraft flying on a fixed course at high speed and constant altitude and recording by radar its height above the waves.) We also assume that  $\partial \zeta(x')/\partial t$  can be measured (by a pair of radar sets, or otherwise).

In §3·1 it is shown how from the statistical analysis of such measurements the moments  $m_n(\theta)$ , for each value of  $\theta$ , can be deduced. In §3·2 it is shown how to obtain the two-dimensional moments  $m_{bq}$  from the moments  $m_n(\theta)$ ; and in §3·3 how to obtain the energy spectrum from the moments  $m_{bq}$ .

3.1. To obtain  $m_n(\theta)$ 

We saw in § 2.2 that the number of zeros of  $\zeta(x')$  per unit horizontal distance x' is given by

$$N_0 = \frac{1}{\pi} \left( \frac{m_2(\theta)}{m_0(\theta)} \right)^{\frac{1}{2}},$$
 (3.1.1)

and in general the number of zeros of the rth derivative of  $\zeta$  is given by

$$N_r = \frac{1}{\pi} \left( \frac{m_{2r+2}(\theta)}{m_{2r}(\theta)} \right)^{\frac{1}{2}}.$$
 (3.1.2)

Now from the record of  $\zeta$ , the numbers  $N_0$ ,  $N_1$ , etc., may be determined by simple counting of zeros, maxima and minima, points of inflexion, and so on.  $m_0(\theta)$  can be determined as the r.m.s. value of  $\zeta$  along the curve. From (3.1.1) we have

$$m_2(\theta) = \pi^2 N_0^2 m_0(\theta) \tag{3.1.3}$$

and from (3·1·2) 
$$m_{2r+2}(\theta) = \pi^2 N_r^2 m_{2r}(\theta).$$
 (3·1·4)

So  $m_2, m_4, \ldots, m_{2r+2}$  can be determined in succession, or else directly from

$$m_{2r+2}(\theta) = \pi^{2r+2} N_0^2 N_1^2 \dots N_r^2 m_0(\theta).$$
(3.1.5)

To obtain the moments of odd order we have to use some property involving the motion of the surface. We take the distribution of the velocities of zeros of  $\zeta(x')$ , which was derived in §2.5. It was shown that the mean velocity of the zeros of  $\zeta$  is given by

$$ar{c} = -rac{m_1'( heta)}{m_2( heta)}, \hspace{1.5cm} (3\cdot 1\cdot 6)$$

47-2

and that the mean velocity of points where the rth derivative vanishes is

$$\bar{c}_r = -\frac{m'_{2r+1}(\theta)}{m_{2r+2}(\theta)}.$$
(3.1.7)

Now  $m'_{2r+2}(\theta)$  is already known, from equation (3.1.5), so from

$$m'_{2r+1}(\theta) = -\bar{c}_r m_{2r+2}(\theta)$$
 (3.1.8)

we may determine  $m'_{2r+1}(\theta)$ .

It is true that the odd moments  $m'_{2r+1}(\theta)$  correspond not to the original function E(u, v) but to  $\sigma(u, v) E(u, v)$ . However, we shall show in § 3.3 how this difficulty can be overcome.

3.2. To obtain  $m_{pq}$ 

In §1.4 we saw that  $m_n(\theta)$  is related to the moments  $m_{pq}$  (p+q=n) by the equation

$$m_{n}(\theta) = m_{n,0} \cos^{n}\theta + {\binom{n}{1}} m_{n-1,1} \cos^{n-1}\theta \sin\theta + \dots + m_{0,n} \sin^{n}\theta.$$
(3.2.1)

The expression on the right-hand side is a trigonometric polynomial of degree n, with coefficients which are linear combinations of the moments. Therefore we may expect to solve for  $m_{pq}$  by taking the Fourier components of  $m_n(\theta)$ , that is, by considering the quantities

$$a_{n,l} = \frac{1}{\pi} \int_0^{2\pi} m_n(\theta) \,\mathrm{e}^{\mathrm{i} l\theta} \,\mathrm{d}\theta. \tag{3.2.2}$$

Going back to equation (1.4.11), we have

$$a_{n,l} = \frac{1}{\pi} \int_0^{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \left( u \cos \theta_1 + v \sin \theta_1 \right)^n \mathrm{d}u \,\mathrm{d}v \,\mathrm{e}^{\mathrm{i}l\theta_1} \,\mathrm{d}\theta_1$$
  
=  $\frac{1}{\pi} \int_0^{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) \left\{ w \cos \left(\theta - \theta_1\right) \right\}^n \mathrm{d}u \,\mathrm{d}v \,\mathrm{e}^{\mathrm{i}l\theta_1} \,\mathrm{d}\theta_1,$  (3.2.3)

where  $(w \cos \theta, w \sin \theta) = (u, v)$ . On writing  $\theta_1 - \theta = \theta_2$  and reversing the order of integration we have

$$a_{n,l} = \frac{1}{\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{0}^{2\pi} E(u,v) w^{n} \cos^{n} \theta_{2} e^{il(\theta+\theta_{2})} d\theta_{2} du dv$$
$$= \gamma_{n,l} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} E(u,v) w^{n} e^{il\theta} du dv, \qquad (3.2.4)$$

where  $\gamma_{n,l}$  is a numerical constant:

$$\gamma_{n,l} = \frac{1}{\pi} \int_0^{2\pi} \cos^n \theta_2 e^{il\theta_2} d\theta_2 = \begin{cases} 2^{1-n} \binom{n}{r} & \text{when} \quad n-l = 2r > 0\\ 0 & \text{otherwise.} \end{cases}$$
(3.2.5)

Now

where 
$$\begin{pmatrix} p & q \\ r & \end{pmatrix}$$
 is the coefficient of  $x^r$  in the expansion of  $(1+x)^p (1-x)^q$ . In full,  
 $\begin{pmatrix} p & q \\ r & \end{pmatrix} = \begin{pmatrix} p \\ r & \end{pmatrix} - \begin{pmatrix} p \\ r-1 \end{pmatrix} \begin{pmatrix} q \\ 1 \end{pmatrix} + \begin{pmatrix} p \\ r-2 \end{pmatrix} \begin{pmatrix} q \\ 2 \end{pmatrix} - \dots + (-1)^r \begin{pmatrix} q \\ r \end{pmatrix},$  (3.2.7)

where  $\binom{p}{0} = 1$  and  $\binom{p}{r} = 0$  for r > p. From (3.2.4) and (3.2.6) we have

$$m_{pq} = \frac{1}{2i^{q}} \left[ a_{n,n} + \binom{p}{1} \frac{q}{n} \right] \binom{n}{1} a_{n,n-2} + \binom{p}{2} \binom{q}{2} \binom{n}{2} a_{n,n-4} + \dots + (-1)^{q} a_{n,n} \right]. \quad (3.2.8)$$

TABLE 2. FUNCTIONS  $C_{pq}(\theta)$ 

$$\begin{split} C_{00} &= \frac{1}{2} & \begin{cases} C_{10} &= \cos \theta \\ C_{01} &= \sin \theta \end{cases} \\ \begin{cases} C_{20} &= & \cos 2\theta + \frac{1}{2} \\ C_{11} &= & \sin 2\theta \\ C_{02} &= & -\cos 2\theta + \frac{1}{2} \end{cases} & \begin{cases} C_{30} &= & \cos 3\theta + \cos \theta \\ C_{21} &= & \sin 3\theta + \frac{1}{3} \sin \theta \\ C_{12} &= & -\cos 3\theta + \frac{1}{3} \cos \theta \\ C_{03} &= & -\sin 3\theta + \sin \theta \end{cases} \\ \begin{cases} C_{40} &= & \cos 4\theta + \cos 2\theta + \frac{1}{2} \\ C_{31} &= & \sin 4\theta + \frac{1}{2} \sin 2\theta \\ C_{22} &= & -\cos 4\theta + \frac{1}{6} \\ C_{13} &= & -\sin 4\theta + \frac{1}{2} \sin 2\theta \\ C_{04} &= & \cos 4\theta - \cos 2\theta + \frac{1}{2} \end{cases} & \begin{cases} C_{50} &= & \cos 5\theta + \cos 3\theta + \cos \theta \\ C_{41} &= & \sin 5\theta + \frac{3}{5} \sin 3\theta + \frac{1}{5} \sin \theta \\ C_{32} &= & -\cos 5\theta - \frac{1}{5} \cos 3\theta + \frac{1}{5} \cos \theta \\ C_{23} &= & -\sin 5\theta + \frac{1}{5} \sin 3\theta + \frac{1}{5} \sin \theta \\ C_{14} &= & \cos 5\theta - \frac{3}{5} \cos 3\theta + \frac{1}{5} \cos \theta \\ C_{05} &= & \sin 5\theta - \sin 3\theta + \sin \theta \end{cases} \end{split}$$

So on substitution from  $(3 \cdot 2 \cdot 2)$ 

$$m_{pq} = \frac{1}{\pi} \int_0^{2\pi} m_n(\theta) C_{pq}(\theta) \,\mathrm{d}\theta, \qquad (3.2.9)$$

where

$$C_{pq}(\theta) = \frac{1}{2\mathbf{i}^q} \left[ e^{\mathbf{i}n\theta} + {\binom{p}{1}} {\binom{q}{1}} / {\binom{n}{1}} e^{\mathbf{i}(n-2)\theta} + \dots + (-1)^n e^{-\mathbf{i}n\theta} \right].$$
(3.2.10)

The quantities  $m_n(\theta)$  being known, this determines the moments  $m_{pq}$ . The first few functions  $C_{bq}(\theta)$  are listed in table 2.

Incidentally, when the spectrum E(u, v) has circular symmetry,  $m_n(\theta)$  is independent of  $\theta$  and so from (3.2.9)

$$m_{pq}(\theta) = \begin{cases} (-1)^{\frac{1}{2}q} \binom{p}{\frac{1}{2}n} / \binom{n}{\frac{1}{2}n} m_n & \text{when } p, q \text{ are both even} \\ 0 & \text{otherwise.} \end{cases}$$
(3.2.11)

In particular

$$m_{20} = m_{02} = m_2, \qquad m_{40} = m_{04} = m_4, \qquad m_{22} = \frac{1}{3}m_4, \qquad (3.2.12)$$

and so the condition  $(1\cdot3\cdot11)$  for a narrow ring spectrum reduces to

$$\frac{8}{3}m_4m_0 - 4m_2^2 = 0 \tag{3.2.13}$$

or

$$\frac{m_4 m_0}{m_2^2} = \frac{3}{2}.$$
 (3.2.14)

As a corollary, we see that  $m_4 m_0/m_2^2$  is never less than  $\frac{3}{2}$ .

47**-**3

#### 3.3. To obtain E(u, v)

We have so far obtained the even moments  $m_{pq}$  of E(u, v) and the odd moments  $m'_{pq}$  of  $\sigma(u, v) E(u, v)$ . Now consider the function

$$F(u,v) = \frac{1}{2} [E(u,v) + E(-u,-v)].$$
(3.3.1)

This is clearly an even function of (u, v), since F(-u, -v) = F(u, v). Therefore its odd moments vanish. But its even moments are the same as those of E(u, v). Therefore both the odd and even moments of F(u, v) are known. Similarly

$$G(u,v) = \frac{1}{2} [\sigma(u,v) E(u,v) - \sigma(-u,-v) E(-u,-v)]$$
(3.3.2)

is clearly an odd function of (u, v), since G(-u, -v) = -G(u, v), and so its even moments vanish. But since  $\sigma(-u, -v) = \sigma(u, v)$  (equation (1·1·5)) the odd moments of G are equal to those of  $\sigma(u, v) E(u, v)$ . Therefore both the odd and even moments of G are known. If F and G can both be determined from their moments we may then determine E, from the identity  $E(u, v) = F(u, v) + G(u, v)/\sigma(u, v)$  (3·3·3)

$$E(u, v) = F(u, v) + G(u, v) / \sigma(u, v).$$
(3.3.3)

We have then simply to consider how to determine F and G from their moments.\*

Formally, if the moments were known to all orders, the problem would be solved. For since the even moments  $m_{pq}$  are equivalent to the derivatives of the correlation function  $\psi(x, y, 0)$  (equation  $(1 \cdot 2 \cdot 10)$ ) we have

$$\psi(x,y,0) = \sum_{p+q=2r} (-1)^r \frac{m_{pq}}{p! \, q!} x^p y^q.$$
(3.3.4)

But by  $(1 \cdot 2 \cdot 9)$ ,  $\psi(x, y, 0)$  is the cosine transform of E(u, v) and so of F(u, v). Hence

$$F(u,v) = \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \psi(x,y,0) \cos(ux+vy) \, \mathrm{d}x \, \mathrm{d}y.$$
(3.3.5)

Similarly, if we define a function

G

$$\psi'(x,y,0) = \sum_{p+q=2r} (-1)^{r+1} \frac{m_{pq}}{p! \, q!} x^p y^q, \qquad (3\cdot3\cdot6)$$

we have

$$(u,v) = \frac{1}{(2\pi)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \psi'(x,y,0) \sin(ux+vy) \, \mathrm{d}x \, \mathrm{d}y.$$
(3.3.7)

In practice, however, only a finite number of moments can be obtained, and if  $(3\cdot3\cdot4)$  is replaced by only a finite number of terms of the series,  $(3\cdot3\cdot5)$  does not converge. The problem then is to find a convergent sequence of approximations to F and G, each approximation depending on the moments of the function up to a finite order.

It was shown by Weierstrass (1885) that a function of a single variable may be approximated over a finite range by a polynomial, and that this may be done in a variety of different ways. A simple method is given by Courant & Hilbert (1953, §4), which we generalize to two dimensions as follows. Consider the function<sup>†</sup>

$$g_n(u,v) = (1 - u^2 - v^2)^n.$$
(3.3.8)

\* For a discussion of whether a function is uniquely determined by its moments, see Kendall (1952, chap. 4).

<sup>†</sup> This is different from the generalization suggested in Courant & Hilbert (1953, p. 68), and leads to a more homogeneous approximation. As *n* tends to infinity,  $g_n$  tends to zero for all values of (u, v) inside the circle  $u^2 + v^2 = 1$  except the origin. Further, if S is any smaller circle of radius  $\delta < 1$ ,

$$\iint_{S} g_{n}(u,v) \, \mathrm{d}u \, \mathrm{d}v = \int_{0}^{\delta} \int_{0}^{2\pi} (1-w^{2})^{n} \, w \, \mathrm{d}w \, \mathrm{d}\theta = \frac{\pi}{n+1} \left[ 1 - (1-\delta^{2})^{n+1} \right], \qquad (3\cdot3\cdot9)$$

which also tends to zero. However, the dominant part of the above integral is contributed by the neighbourhood of the origin, that is, if S' is any interior circle of fixed radius  $\delta'$ , however small, almost the entire contribution to the integral comes from S':

$$\lim_{n \to \infty} \frac{\iint_{s'} g_n(u, v) \, \mathrm{d}u \, \mathrm{d}v}{\iint_{s} g_n(u, v) \, \mathrm{d}u \, \mathrm{d}v} = \lim_{n \to \infty} \frac{1 - (1 - \delta'^2)^{n+1}}{1 - (1 - \delta^2)^{n+1}} = 1.$$
(3.3.10)

Now suppose that f(u, v) is any continuous function of two variables that we wish to approximate in the region S,  $(u^2+v^2)^{\frac{1}{2}} < \frac{1}{2}$ . Then if (u, v) is any interior point of S, the function

$$f_n(u,v) = \frac{\iint_{S} f(u_1,v_1) \left[1 - (u - u_1)^2 - (v - v_1)^2\right]^n \mathrm{d}u_1 \mathrm{d}v_1}{\iint_{S} [1 - u_1^2 - v_1^2]^n \mathrm{d}u_1 \mathrm{d}v_1}$$
(3.3.11)

is a weighted mean of f(u, v), with weighting function  $g_n(u-u_1, v-v_1)$  centred on (u, v). And since the neighbourhood of (u, v) contributes almost all the weight when n is large we see that  $\lim_{n\to\infty} f_n(u, v) = f(u, v)$ . (3.3.12)

The convenience of this approximation lies in the fact that  $f_n(u, v)$  is a polynomial in (u, v) of degree 2n, and with coefficients that are definite integrals taken over S. Further, if we assume that f(u, v) is negligible or zero outside S the coefficients in  $f_n(u, v)$  are simply combinations of the moments of f of order not greater than 2n.

To apply the representation in the present case let us assume that E(u, v) is negligible when  $(u^2+v^2)^{\frac{1}{2}} > \frac{1}{2}w_0$ , say. In other words, we assume a cut-off at high wave-numbers (some such assumption is in any case necessary in order to ensure the uniqueness of the solution.) Then we take as an approximation to F(u, v)

$$F_n(u,v) = \frac{n+1}{\pi w_0^2} \iint_{S_1} F(u_1,v_1) \left[ 1 - (u-u_1)^2 / w_0^2 - (v-v)^2 / w_0^2 \right]^n \mathrm{d}u_1 \mathrm{d}v_1, \qquad (3\cdot 3\cdot 13)$$

where  $S_1$  is the region  $(u_1^2 + v_1^2)^{\frac{1}{2}} \leq \frac{1}{2}w_0$ . Similarly we take

$$G_n(u,v) = \frac{n+1}{\pi w_0^2} \iint_{S_1} G(u_1,v_1) \left[ 1 - (u-u_1)^2 / w_0^2 - (v-v_1)^2 / w_0^2 \right]^n \mathrm{d}u_1 \mathrm{d}v_1, \qquad (3\cdot 3\cdot 14)$$

and finally

$$E_n(u,v) = F_n(u,v) + G_n(u,v) / \sigma(u,v).$$
 (3.3.15)

On expanding the polynomial expressions in  $(3\cdot3\cdot13)$  and  $(3\cdot3\cdot14)$  and carrying out the integrations we find, say, for n = 2,

$$F_{2}(u,v) = \frac{3}{\pi} \left[ m_{00}(w_{0}^{2} - u^{2} - v^{2}) - 2(m_{20} + m_{02}) (w_{0}^{2} - u^{2} - v^{2}) w_{0}^{2} + (m_{40} + m_{04} + 2m_{22}) w_{0}^{4} \right] \quad (3\cdot3\cdot16)$$

and

$$G_{2}(u,v) = \frac{3}{\pi} \left[ 4(m'_{10}u + m'_{01}v) (w_{0}^{2} - u^{2} - v^{2}) w_{0} - 4 \left\{ (m'_{30} + m'_{12}) u + (m'_{21} + m'_{03}) v \right\} w_{0}^{3} \right], \quad (3\cdot3\cdot17)$$

so that  $F_2$  and  $G_2$  are expressible as polynomials in (u, v) having as coefficients the moments of E up to degree 4. Approximations of higher order may be written down at will.



FIGURE 14



FIGURE 14. Successive approximations  $E_n(u, v) = W_n(u, v; w_0/4, 0)$ .

We have seen that  $F_n$  and  $G_n$  are essentially weighted averages of F and G by a weighting function proportional to  $g_n[(u-u_1)/w_0, (v-v_1)/w_0]$ . The weighting function corresponding to  $E_n$  is somewhat different owing to the presence of the factor  $\sigma(u, v)$  in (3·3·15). In fact we have from (3·3·1) and (3·3·2)

$$\begin{split} E_n(u,v) &= \frac{n+1}{2\pi w_0^2} \iint_{S_1} E(u_1,v_1) \left[ 1 + \frac{\sigma(u_1,v_1)}{\sigma(u,v)} \right] \left[ 1 - \left(\frac{u-u_1}{w_0}\right)^2 - \left(\frac{v-v_1}{w_0}\right)^2 \right]^n \mathrm{d}u_1 \mathrm{d}v_1 \\ &+ \frac{n+1}{2\pi w_0^2} \iint_{S_1} E(-u_1,-v_1) \left[ 1 - \frac{\sigma(u_1,v_1)}{\sigma(u,v)} \right] \left[ 1 - \left(\frac{u+u_1}{w_0}\right)^2 - \left(\frac{v+v_1}{w_0}\right)^2 \right]^n \mathrm{d}u_1 \mathrm{d}v_1. \end{split}$$
(3.3.18)

On changing the sign of  $(u_1, v_1)$  in the second integral we have

$$E(u,v) = \frac{1}{w_0^2} \iint_{S_1} E(u_1,v_1) W_n(u,v;u_1,v_1) \,\mathrm{d}u_1 \,\mathrm{d}v_1, \qquad (3\cdot 3\cdot 19)$$

where

$$\begin{split} W_n(u,v;u_1,v_1) &= \frac{n+1}{2\pi} \bigg[ 1 + \frac{\sigma(u_1,v_1)}{\sigma(u,v)} \bigg] \bigg[ 1 - \left(\frac{u-u_1}{w_0}\right)^2 - \left(\frac{v-v_1}{w_0}\right)^2 \bigg]^n \\ &+ \frac{n+1}{2\pi} \bigg[ 1 - \frac{\sigma(u_1,v_1)}{\sigma(u,v)} \bigg] \bigg[ 1 - \left(\frac{u+u_1}{w_0}\right)^2 - \left(\frac{v+v_1}{w_0}\right)^2 \bigg]^n. \end{split}$$
(3·3·20)

 $W_n$  is not a function of  $(u-u_1)$  and  $(v-v_1)$  alone. However, the second half of  $(3\cdot 3\cdot 20)$  only gives a contribution when  $(u_1, v_1) \doteq (u, v)$ , and then this is small owing to the presence of the factor  $[1 - \sigma(u_1, v_1)/\sigma(u, v)]$ .

To obtain an idea of the accuracy of successive approximations we may consider the case of a narrow spectrum, when E(u, v) is appreciably large only in the neighbourhood of a single point, say  $(-\frac{1}{4}w_0, 0)$ . Then

$$E_n(u,v) = W_n(u,v; -\frac{1}{4}w_0, 0).$$
(3.3.21)

 $W_n$  has been computed for n = 2, 4, 8 assuming that, as for gravity waves on deep water,

$$\sigma(u, v) \propto (u^2 + v^2)^{\frac{1}{4}}.$$
 (3.3.22)

The results are shown in figure 14 *a*, *b* and *c*. It will be seen how the functions become progressively more peaked as the degree of the approximation is raised. When n = 8 the area in which  $W_n$  exceeds half its maximum value has a radius of about  $0.3w_0$ . For large values of *n* we have, in the neighbourhood of  $(u_1, v_1)$ ,

$$W_n \doteq \frac{n}{\pi} \exp\left\{-n[(u-u_1)^2 + (v-v_1)^2]/w_0^2\right\}, \qquad (3\cdot3\cdot23)$$

and so the 'radius' of  $W_n$  is proportional to  $n^{-\frac{1}{2}}$ . It will be seen then that  $E_n$  converges to E rather slowly. In order to distinguish parts of the spectrum separated by a distance  $\delta$ , it is necessary to take n to be of order  $(w_0/\delta)^2$ .

I am indebted to Mr D. E. Cartwright for advice in evaluating the elliptic integral  $(2\cdot 4\cdot 47)$  and to Mr E. A. Steer and Miss S. A. Yeo for assistance with the computation for figures 12 and 13.

#### References

- Barber, N. F., Ursell, F., Darbyshire, J. & Tucker, M. J. 1946 A frequency analyser used in the study of ocean waves. *Nature, Lond.*, 158, 329–332.
- Barber, N. F. & Ursell, F. 1948 The generation and propagation of ocean waves and swell. I. Wave periods and velocities. *Phil. Trans.* A, 140, 527–560.
- Barber, N. F. 1950 Ocean waves and swell. Lecture, Institution of Civil Engineers, London, 22 pp.
- Cartwright, D. E. 1956 On determining the directions of waves from a ship at sea. Proc. Roy. Soc. A, 234, 382-387.
- Courant, R. & Hilbert, D. 1953 Methods of mathematical physics. 1st English edition. New York: Interscience Publishers, Inc.
- Cox, C. & Munk, W. 1954*a* Measurement of the roughness of the sea surface from photographs of the sun's glitter. J. Opt. Soc. Amer. 44, 838–850.
- Cox, C. & Munk, W. 1954b Statistics of the sea surface derived from the sun's glitter. J. Mar. Res. 13, 198–227.
- Cramér, H. 1937 Random variables and probability distributions. Cambridge Mathematical Tract, no. 36. Cambridge University Press.
- Duntley, S. Q. 1950 The visibility of submerged objects. Part I. Optical effects of water waves. Report, Visibility Lab., Mass. Inst. Techn.
- Eckart, C. 1946 The sea surface and its effect on the reflection of sound and light. University of California, Division of War Research, Report M407 (unpublished).
- Eckart, C. 1953 a The scattering of sound from the sea surface. J. Acoust. Soc. Amer. 25, 566-570.

Eckart, C. 1953 b The generation of wind waves on a water surface. J. Appl. Phys. 24, 1485–1494. Eckart, C. 1953 c Relation between time averages and ensemble averages in the statistical dynamics

of continuous media. Phys. Rev. 91, 784–790.

- Euler, L. 1752-3 Elementa doctrinae solidorum. Nov. Comment. Acad. Sci. Imp. Petropol. 4, 109-160.
- Kendall, M. G. 1952 The advanced theory of statistics, vol. 1. London: Griffin. 457 pp.
- Khintchine, A. 1934 Korrelationstheorie der stationären stochastischen Prozesse. Math. Ann. 109, 604–615.
- Legendre, A. M. 1811 Exercises de calcul integrale, t. 1. Paris: Courcier. 386 pp.
- Longuet-Higgins, M. S. & Barber, N. F. 1946 Four theoretical notes on the estimation of sea conditions. Admiralty Research Laboratory, Teddington, Report N1–N4 103.30/W (unpublished).
- Longuet-Higgins, M. S. 1950 A theory of the origin of microseisms. Phil. Trans. A, 243, 1-35.
- Longuet-Higgins, M. S. 1952 On the statistical distribution of the heights of sea waves. J. Mar. Res. 9, 245-266.
- Marks, W. 1954 The use of a filter to sort out directions in a short-crested gaussian sea surface. Trans. Amer. Geophys. Un. 35, 758-766.
- Pierson, W. J. 1952 A unified mathematical theory for the analysis, propagation and refraction of storm generated ocean surface waves, Part I. Report, Dept. of Meteorology, N.Y. Univ., 336 pp.
- Rayleigh, Lord 1880 On the resultant of a large number of vibrations of the same pitch and of arbitrary phase. *Phil. Mag.* 10, 73–78.
- Rayleigh, Lord 1945 The theory of sound, 1st American edition. New York: Dover Publications.
- Rice, S. O. 1944 The mathematical analysis of random noise. Bell Syst. Tech. J. 23, 282-332.
- Rice, S. O. 1945 The mathematical analysis of random noise. Bell. Syst. Tech. J. 24, 46-156.
- Rudnick, P. 1950 Correlograms for Pacific Ocean waves. Proc. 2nd Berkeley Symposium on Mathematical Statistics and Probability, pp. 627–638, University of California Press.
- St Denis, M. & Pierson, W. J. 1953 On the motions of ships in confused seas. Trans. Soc. Nav. Archit., N.Y., 61, 1–65.
- Schumacher, A. 1952 Results of exact wave measurements (by stereophotogrammetry) with special reference to more recent theoretical investigations. Symposium on Gravity Waves, Washington, June 1951. Nat. Bur. Stand. Circ. no. 521, U.S. Govt. Printing Office.
- Schooley, A. H. 1954 A simple optical method for measuring the statistical distribution of water surface slopes. J. Opt. Soc. Amer. 44, 37-40.
- Sommerville, D. M. Y. 1929 An introduction to the geometry of n dimensions. London: Methuen. 196 pp.
- Taylor, G. I. 1922 Diffusion by continuous movements. Proc. Lond. Math. Soc. (2), 20, 196-212.
- Tukey, J. W. 1949 The sampling theory of power spectrum estimates. Symposium on application of autocorrelation analysis to physical problems, Woods Hole, Mass., June 1949. Office of Naval Research, Washington, D.C.
- Tukey, J. W. & Hamming, R. W. 1949 *Measuring noise color*. Bell Telephone Laboratories Internal Memorandum, MM 49.
- Watters, J. K. 1953 Distribution of height in ocean waves. N.Z. J. Sci. Tech. B, 34, 409-422.
- Weierstrass, K. T. W. 1885 Über die analytische Darstellbarkeit sogenannter willkürlicher Funktionen reeller Argumente. S.B. Akad. Wiss. Berl. 633–639, 789–805.
- Weierstrass, K. T. W. 1903 Werke, vol. 3, 1-37. Berlin.

Whittaker, E. T. & Watson, G. N. 1952 Modern analysis. 4th edition. Cambridge University Press.