

Wave reflection from a gently sloping beach

By JOHN MILES

Institute of Geophysics and Planetary Physics, University of California, San Diego,
 La Jolla, CA 92093, USA

(Received 10 July 1989)

The linear reflection of an obliquely incident gravity wave of frequency ω from a gently sloping beach of shoreline slope σ and characteristic length l is determined for $\sigma \ll 1 \ll \omega^2 l/g$. An asymptotic ($\sigma \downarrow 0$), inviscid approximation that is uniformly valid in the shallow-water domain is matched to Keller's (1958) geometrical-optics approximation for non-shallow water. An exact solution is obtained for the profile $h = \sigma l[1 - \exp(-x/l)]$ in the shallow-water domain and used to test the asymptotic approximation. The absence of viscosity implies perfect reflection. A model that incorporates both small viscosity and small capillarity predicts a fixed contact line and the reflection coefficient $|R| = \exp[-\pi\sigma^{-2}g^{-1}(2\nu\omega^3)^{\frac{1}{2}}]$, where ν is the kinematic viscosity. These predictions are in qualitative agreement with the experimental results of Mahony & Pritchard (1980).

1. Introduction

I consider here the linear reflection of a gravity wave from a gently sloping beach of depth

$$h(x) = \sigma l H(x/l) \sim \begin{cases} \sigma x & (x/l \downarrow 0) \\ h_\infty & (x/l \uparrow \infty) \end{cases}, \quad (1.1)$$

where σ is the shoreline ($x = 0$) slope, $l = O(h_\infty/\sigma)$ is a characteristic length of the beach, H is a smooth, monotonically increasing function, and h_∞ is the offshore depth. The characteristic length for a wave of frequency ω ranges from $(h_\infty/K)^{\frac{1}{2}}$ to $1/K$, where

$$K \equiv \omega^2/g, \quad (1.2)$$

and the parameters of the inviscid problem are σ and Kh_∞ . The free-surface displacement has the form

$$\zeta(x, y, t) = a_0 Z(x) \cos(\omega t - k_y y) \quad (0 \leq x < \infty, \quad -\infty < y < \infty), \quad (1.3)$$

where a_0 is the shoreline ($x = 0$) amplitude, k_y is the longshore wavenumber, and $Z(x)$ is to be determined subject to the boundary conditions ($\zeta_{\max} \equiv a_0$ and mass flux = 0)

$$Z = 1, \quad hZ' = 0 \quad (x = 0). \quad (1.4a, b)$$

The absence of dissipation implies perfect reflection, by virtue of which

$$Z \sim A \cos[(k_\infty^2 - k_y^2)^{\frac{1}{2}}x + \psi] \quad (k_\infty x \uparrow \infty), \quad (1.5)$$

where k_∞ is determined by the gravity-wave dispersion relation

$$k_\infty \tanh(k_\infty h_\infty) = K, \quad (1.6)$$

and the offshore amplitude $a_\infty \equiv Aa_0$ and phase shift ψ are to be obtained as part of the solution (although in practice a_∞ is specified and a_0 is inferred from $A = a_\infty/a_0$).

The earliest solutions of this linear, inviscid reflection problem (see Stoker 1957, §5.1) are for uniform slope with $\sigma = \tan(\pi/2n)$, $n = 1, 2, \dots$, but these solutions are cumbersome for the most important (for oceanography) case of small slope. Matched asymptotic approximations for normal incidence and $\sigma \ll 1$ have been determined for uniform slope by Friedrichs (1948) and for non-uniform slope by Keller (1961), who matched his (1958) geometrical-optics approximation for $Kl \gg 1$ to the shallow-water approximation (Lamb 1932, §186)

$$Z = J_0[2(Kx/\sigma)^{1/2}] \quad (h \sim \sigma x \ll 1/K), \quad (1.7)$$

where J_0 is a Bessel function. Carrier & Greenspan (1958) have solved the nonlinear problem for non-breaking reflection from a uniform slope and shown that $Ka_0 < \sigma^2$ is necessary for a smooth solution. Their solution reduces to (1.7) for $Ka_0 \ll \sigma^2$.

The approximation (1.7) is often adopted in the investigation of edge waves (see Guza 1985 for a review) and other shore processes on the hypothesis that the disturbances associated with these processes are essentially confined to the domain of uniform slope; however, this hypothesis fails in some problems, and, in any event, its testing requires the development of approximations that incorporate non-uniform slope. Towards that end, I develop (in §2) an approximation that is uniformly valid in $x = O(l)$ for $\sigma \ll 1 \ll Kl \ll 1/\sigma$ and (in §4) match that approximation to a geometrical-optics approximation for arbitrary Kh_∞ . In §3, I obtain an exact solution of the shallow-water equations for an exponential profile through an extension of Ball's (1967) solution for edge waves. This exact solution provides a test of the asymptotic approximations of §2.

Viscosity is almost always significant in laboratory experiments (although it may be negligible for non-breaking waves of geophysical scale). In §5, I consider its effects on normally incident waves on the assumption of no slip at the bottom, $K\delta_* \ll \sigma^2$, and $\lambda = O(\delta_*/\sigma)$, where $\delta_* \equiv (\nu/2\omega)^{1/2}$ is a viscous lengthscale (ν is the kinematic viscosity), and λ is the capillary length (2.8 mm for clean water). The corresponding extension of the shallow-water equation (Miles 1990) predicts total absorption (zero reflection) of the incident wave if capillarity is neglected ($\lambda = 0$). If $0 < \lambda = O(\delta_*/\sigma)$ capillarity is significant only in an inner approximation, which may be matched to an outer, boundary-layer approximation (which assumes $h \gg \delta_*$ and manifestly fails as $h \downarrow 0$). This matching determines the reflection coefficient. If h is given by (1.1) for $x \leq x_1$ (so that $h_\infty = \sigma x_1$), as in a typical laboratory experiment, the reflection coefficient referred to $x = x_1$ is given by

$$|R_1| = \exp[-2\pi(K\delta_*/\sigma^2)] \quad (1.8)$$

for a clean free surface. The exponent is doubled for a fully contaminated surface. (Note that dissipation implies an exponential increase/decrease of the incident/reflected wave as $x \uparrow \infty$, in consequence of which $|R|$ is intrinsically sensitive to the location of the reference station.) It should be emphasized that the assumption of no slip at the bottom implies a fixed contact line. This prediction is supported by Mahony & Pritchard's (1980) laboratory experiments, but not by oceanographic observation.

Mahony & Pritchard (1980) measured $|R_1| = 0.114$ for $\sigma = 0.09$ and $2\pi/\omega = 0.70$ s, for which (1.8) yields $|R_1| = 0.22$ on the assumption of a clean surface or 0.048 for a fully contaminated surface. Guza & Bowen (1976) report almost perfect reflection for $\sigma = 0.1$ and $2\pi/\omega = 2.4$ – 3.4 s, for which (1.8) predicts $|R_1| = 0.90$ – 0.82 ; the difference $1 - |R_1|$ is within the accuracy of their measurements, but their contact line was moving (Guza, private communication).

2. Shallow-water domain

The assumptions of incompressible, irrotational fluid motion with

$$Ka_0 \ll \sigma^2, \quad Kh \ll 1, \quad (2.1a, b)$$

lead to the shallow-water equation (Lamb 1932, §193)

$$g\nabla \cdot (h\nabla \zeta) = \partial^2 \zeta / \partial t^2. \quad (2.2)$$

Substituting (1.3) into (2.2) and invoking $\omega^2/g \equiv K$, we obtain the Sturm–Liouville equation

$$(hZ')' + (K - k_y^2 h)Z = 0. \quad (2.3)$$

We seek the solution of (2.3), subject to (1.4), for

$$\epsilon \equiv \sigma/Kl \ll 1. \quad (2.4)$$

2.1. Asymptotic solution

Guided by (1.7) and by Erdélyi's (1955, §4.1) treatment of Liouville's problem, we posit

$$Z(x) = [h(K - k_y^2 h)]^{-\frac{1}{2}} (\frac{1}{2}\sigma\chi)^{\frac{1}{2}} f(\chi), \quad (2.5a)$$

where
$$\chi = \int_0^x \left(\frac{K - k_y^2 h}{h} \right)^{\frac{1}{2}} dx = \epsilon^{-\frac{1}{2}} \int_0^\xi (H^{-1} - \beta)^{\frac{1}{2}} d\xi \quad (\xi \equiv x/l), \quad (2.5b)$$

and
$$\beta \equiv \sigma k_y^2 l / K = \epsilon (k_y l)^2 \quad (2.6)$$

is a measure of obliquity. Transforming (2.3) and (1.4), we obtain

$$f'' + \chi^{-1} f' + f = r f \quad (2.7)$$

and
$$f = 1, \quad \chi f' = 0 \quad (\chi = 0), \quad (2.8a, b)$$

where
$$r = \frac{1}{4} \left\{ \frac{(K - 2k_y^2 h) h''}{(K - k_y^2 h)^2} - \frac{(K^2 + 4k_y^2 h^2) h'^2}{4(K - k_y^2 h)^3 h} + \chi^{-2} \right\} \equiv r(\chi), \quad (2.9)$$

$f' \equiv df/d\chi$, and $h' \equiv dh/dx$. We remark that $r(\chi)$ is regular and has the limiting values

$$r(0) = \frac{1}{6}\epsilon(H_0'' - \beta), \quad r \sim \frac{1}{4}\chi^{-2} (\epsilon\chi^2 \uparrow \infty). \quad (2.10a, b)$$

A first approximation to the solution of (2.7) and (2.8) is given by

$$f = J_0(\chi) + O(\epsilon). \quad (2.11)$$

It follows that
$$Z(x) = [h(K - k_y^2 h)]^{-\frac{1}{2}} (\frac{1}{2}\sigma\chi)^{\frac{1}{2}} J_0(\chi) + O(\epsilon), \quad (2.12)$$

where χ is given by (2.5b), provides a uniformly valid approximation to the solution of (1.4) and (2.3). Letting $\sigma\chi \downarrow 0$, we recover the inner approximation (1.7). Letting $\chi \uparrow \infty$, we obtain the outer approximation

$$Z(x) \sim (\sigma/\pi)^{\frac{1}{2}} [h(K - k_y^2 h)]^{-\frac{1}{2}} [\cos(\chi - \frac{1}{4}\pi) + O(\chi^{-1})] \quad (\chi \uparrow \infty). \quad (2.13)$$

Letting $h \sim h_\infty$ in (2.13) with $Kh_\infty \ll 1$, comparing with (1.5), and invoking

$$k_y^2 h_\infty / K = (k_y/k_\infty)^2 = \sin^2 \theta_1 \quad (Kh_\infty \ll 1), \quad (2.14)$$

where $k_\infty = \omega/(gh_\infty)^{\frac{1}{2}}$ is the offshore wavenumber and θ_1 is the angle of incidence, we obtain

$$A = (\sigma/\pi)^{\frac{1}{2}} (Kh_\infty)^{-\frac{1}{2}} (\sec \theta_1)^{\frac{1}{2}} = \sigma^{\frac{1}{2}} (\pi k_\infty h_\infty \cos \theta_1)^{-\frac{1}{2}} \quad (2.15a)$$

and
$$\psi = \int_0^\infty [(Kh^{-1} - k_y^2)^{\frac{1}{2}} - k_\infty \cos \theta_1] dx - \frac{1}{4}\pi. \quad (2.15b)$$

2.2. Higher approximations

To construct higher approximations, we regard (2.7) as an inhomogeneous Bessel equation, solve by variation of parameters, and invoke (2.8) to obtain the Volterra integral equation

$$f(\chi) = J_0(\chi) + \int_0^\chi G(\chi, \hat{\chi}) r(\hat{\chi}) f(\hat{\chi}) d\hat{\chi}, \quad (2.16a)$$

where
$$G(\chi, \hat{\chi}) = \frac{1}{2}\pi \hat{\chi} [J_0(\hat{\chi}) Y_0(\chi) - J_0(\chi) Y_0(\hat{\chi})]. \quad (2.16b)$$

The solution of (2.16) may be obtained by iteration. Substituting the first approximation $f(\hat{\chi}) = J_0(\hat{\chi})$ into (2.16a) and integrating by parts, we obtain the second approximation

$$f(\chi) = J_0(\chi) + \int_0^\chi G(\chi, \hat{\chi}) J_0(\hat{\chi}) r(\hat{\chi}) d\hat{\chi} + O(\epsilon^2) \quad (2.17a)$$

$$= J_0(\chi) + \epsilon r(\chi) \int_0^\chi G(\chi, \hat{\chi}) J_0(\hat{\chi}) d\hat{\chi} + O(\epsilon^2) \quad (2.17b)$$

$$= J_0(\chi) + \frac{1}{2}r(\chi) J_1(\chi) + O(\epsilon^2) = J_0[(1-r)^{\frac{1}{2}}\chi] + O(\epsilon^2) \quad (2.17c)$$

((2.17c) may be derived heuristically by regarding r as slowly varying in (2.7)). The outer approximation (2.13) remains unchanged in these higher approximations and is valid to any algebraic order in ϵ (but it may exhibit an exponentially small error as $\epsilon \downarrow 0$; see §3).

3. Exponential profile

The shallow-water equation (2.3) admits exact solutions for (cf. Ball 1967)

$$h = \sigma l (1 - e^{-x/l}) \quad (H = 1 - e^{-\xi}, \quad h_\infty = \sigma l). \quad (3.1)$$

Adopting H as the independent variable, we obtain

$$Z = \text{Re}\{e^{i\tau\xi} F(\tfrac{1}{2} + \rho - i\tau, \tfrac{1}{2} - \rho - i\tau; 1; 1 - e^{-\xi})\} \quad (3.2a)$$

$$= \text{Re}\{\mathcal{A} e^{i\tau\xi} F(\tfrac{1}{2} + \rho - i\tau, \tfrac{1}{2} - \rho - i\tau; 1 - 2i\tau; e^{-\xi})\}, \quad (3.2b)$$

where Re signifies the real part of, F is Gauss's hypergeometric function,

$$\rho \equiv (\tfrac{1}{4} + \epsilon^{-1} \sin^2 \theta_1)^{\frac{1}{2}}, \quad \tau \equiv \epsilon^{-\frac{1}{2}} \cos \theta_1, \quad (3.3a, b)$$

$$\mathcal{A} = \frac{2\Gamma(2i\tau)}{\Gamma(\tfrac{1}{2} + \rho + i\tau) \Gamma(\tfrac{1}{2} - \rho + i\tau)}, \quad (3.4)$$

and θ_1 is the angle of incidence given by (2.14) with $h_\infty = \sigma l$ therein.

Letting $\xi \uparrow \infty$ in (3.2b), comparing the result with (1.5), and invoking $\Gamma(z) \times \Gamma(1-z) = \pi \csc \pi z$ and $|\Gamma(iy)| = (\pi/y \sinh \pi y)^{\frac{1}{2}}$, we obtain

$$A = |\mathcal{A}| = (\pi\tau)^{-\frac{1}{2}} (\cos^2 \pi\rho \coth \pi\tau + \sin^2 \pi\rho \tanh \pi\tau)^{\frac{1}{2}} \quad (3.5a)$$

and
$$\psi = \arg \mathcal{A} = \arg \Gamma(2i\tau) - 2 \arg \Gamma(\tfrac{1}{2} + \rho + i\tau) + \tan^{-1} (\tan \pi\rho \tanh \pi\tau). \quad (3.5b)$$

The approximation implied by (2.15a) with $h_\infty = \sigma l$ therein is $A = (\pi\tau)^{-\frac{1}{2}}$, which is

exponentially close to (3.5a) in the limit $\epsilon \downarrow 0$ and differs therefrom by less than 0.01 % for $\epsilon \leq \frac{1}{2}$.

It does not appear possible to reduce (3.5b) to elementary transcendents, but asymptotic ($\epsilon \downarrow 0$) approximations may be obtained from the asymptotic expansion of the gamma function. For normal incidence ($\rho = \frac{1}{2}, \tau = \epsilon^{-\frac{1}{2}}$),

$$\psi \sim \epsilon^{-\frac{1}{2}} \ln 4 - \frac{1}{4}\pi + \frac{1}{8}\epsilon^{\frac{1}{2}} + O(\epsilon^{\frac{3}{2}}). \quad (3.6)$$

The corresponding approximation implied by (2.15b) is

$$\psi = \epsilon^{-\frac{1}{2}} \int_0^\infty [(1 - e^{-\xi})^{-\frac{1}{2}} - 1] d\xi - \frac{1}{4}\pi = \epsilon^{-\frac{1}{2}} \ln 4 - \frac{1}{4}\pi, \quad (3.7)$$

which differs from (3.6) by less than 7.5 % for $\epsilon < \frac{1}{2}$.

4. Transition to non-shallow water

We now suppose that $Kh \ll 1$ is satisfied out to a depth for which (2.13) is a valid approximation, but that h continues to increase (with x) to non-shallow values. It then follows from Keller's (1958) geometrical-optics approximation that $Z(x)$ has the form

$$Z = A(x) \cos [\Psi(x)] + O(1/Kl), \quad (4.1)$$

where A and Ψ satisfy

$$\Psi'^2 + k_y^2 = k^2, \quad [A^2 \operatorname{sech}^2 kh (\sinh^2 kh + Kh) \Psi']' = 0, \quad (4.2a, b)$$

$$\text{and} \quad k \tanh kh = K. \quad (4.3)$$

Integrating (4.2), matching (4.1) to (2.13) to determine the constants of integration, and eliminating the hyperbolic functions with the aid of (4.3), we obtain

$$\Psi = \int_0^x (k^2 - k_y^2)^{\frac{1}{2}} dx - \frac{1}{4}\pi, \quad (4.4a)$$

$$A = (2\sigma\pi)^{\frac{1}{2}} k (k^2 - k_y^2)^{-\frac{1}{4}} [K + h(k^2 - K^2)]^{-\frac{1}{2}}. \quad (4.4b)$$

Letting $k_y = 0$, we recover Keller's (1961) results.

Letting $h \rightarrow h_\infty \rightarrow \infty$ in (4.3) and (4.4), we obtain $k \rightarrow k_\infty \rightarrow K$ and

$$A \sim (2\sigma/\pi)^{\frac{1}{2}} (\sec \theta_i)^{\frac{1}{2}} \quad (Kh_\infty \uparrow \infty) \quad (4.5)$$

as the deep-water counterpart of the shallow-water approximation (2.15a).

5. Viscous and capillary effects

We now admit viscosity and capillarity in the two-dimensional, shallow-water domain. Assuming normal incidence and replacing (1.3) by

$$\zeta = a \operatorname{Re} \{Z(x) e^{-i\omega t}\}, \quad (5.1)$$

where a is an amplitude scale, Re implies the real part of, and Z is a dimensionless, complex amplitude, we find that (2.3) is replaced by (Miles 1990)

$$[p(Z - \lambda^2 Z'')]' + KZ = 0, \quad p = h - \delta \tanh(h/\delta), \quad (5.2a, b)$$

where

$$\delta \equiv (\nu/\omega)^{\frac{1}{2}} e^{i\frac{1}{2}\pi} \equiv (1+i) \delta_*, \quad (5.3)$$

$\delta_* = (\nu/2\omega)^{\frac{1}{2}}$ is a viscous lengthscale, and $\lambda \equiv (T/\rho g)^{\frac{1}{2}}$ is the capillary length (T is the surface tension). The corresponding boundary conditions (for $h \sim \sigma x$ as $x \downarrow 0$) are

$$Z = 0, \quad p(Z - \lambda^2 Z'')' = 0 \quad (x = 0). \quad (5.4a, b)$$

The preceding formulation is for a clean surface. If the surface is fully contaminated (inextensible) δ is replaced by 2δ in (5.2b).

5.1. Inner and outer approximations

We proceed on the assumptions that $|\alpha| \ll 1$ and $\gamma = O(1)$, where

$$\alpha = \frac{K\delta}{\sigma^2}, \quad \gamma \equiv \frac{\lambda\sigma}{\delta}. \quad (5.5a, b)$$

Capillarity then is negligible in $h \gg \delta_*$, and (5.2) admits an outer approximation of the form (cf. (2.12) with $k_y = 0$ therein)

$$Z = [K(h - \delta)]^{-\frac{1}{2}} (\frac{1}{2}\sigma\chi)^{\frac{1}{2}} [J_0(\chi) \cos \chi_0 - Y_0(\chi) \sin \chi_0], \quad (5.6a)$$

where
$$\chi = \int_{\delta/\sigma}^x \left(\frac{K}{h - \delta} \right)^{\frac{1}{2}} dx, \quad (5.6b)$$

and, by hypothesis, $\chi_0 = O(\alpha)$.

Assuming that $h \approx \sigma x$ for $\sigma x = O(\delta)$ and introducing the inner variables

$$\xi = \frac{h}{\delta} = \frac{\sigma x}{\delta}, \quad Z = Z_1(\xi), \quad (5.7a, b)$$

we transform (5.2) to

$$[\varpi(Z_1 - \gamma^2 Z_1'')] + \alpha Z_1 = 0, \quad \varpi = \xi - \tanh \xi. \quad (5.8a, b)$$

Integrating (5.8a) from 0 to ξ , so that (5.4b) is satisfied, dividing by ϖ , and integrating again, we obtain

$$Z_1 - \gamma^2 Z_1'' = C_1 + \alpha F(\xi), \quad F(\xi) \equiv - \int_{\xi_1}^{\xi} \frac{d\eta}{\varpi(\eta)} \int_0^{\eta} Z_1(\zeta) d\zeta, \quad (5.9a, b)$$

where C_1 and ξ_1 are interdependent constants of integration. Integrating (5.9a) and invoking (5.4a) and $Z_1 = O(1)$ for $\xi \rightarrow \infty$, we obtain

$$Z_1 = C_1 (1 - e^{-\xi/\gamma}) + \frac{\alpha}{2\gamma} \int_0^{\infty} [e^{-|\xi-\eta|/\gamma} - e^{-(\xi+\eta)/\gamma}] F(\eta) d\eta. \quad (5.10)$$

The integral equation (5.10) may be solved by iteration, starting from the first approximation $C_1[1 - \exp(-\xi/\gamma)]$, the substitution of which into (5.9b) yields

$$F = -C_1 \int_{\xi_1}^{\xi} \left[\frac{\eta - \gamma(1 - e^{-\eta/\gamma})}{\varpi(\eta)} \right] d\eta, \quad (5.11)$$

where, here and subsequently, error factors of $1 + O(\alpha)$ are implicit. Letting $\xi \rightarrow \infty$, anticipating that $|\xi_1| \gg 1$ so that $\varpi(\xi) \sim \xi - 1$ holds throughout the range of integration in (5.11), and substituting the resulting approximation to F into (5.9a), we obtain

$$Z_1 \sim C_1 \left\{ 1 - \alpha \left[\xi - \xi_1 + (1 - \gamma) \log \left(\frac{\xi - 1}{\xi_1 - 1} \right) \right] + O \left[\frac{\alpha \gamma^2 (1 - \gamma)}{(\xi - 1)^2} \right] \right\}. \quad (5.12)$$

Returning to the outer approximation (5.6) and letting $h \sim \sigma x$ and $\chi \rightarrow 0$, we obtain

$$\chi \sim 2\sigma^{-1} [K(h - \delta)]^{\frac{1}{2}} = 2[\alpha(\xi - 1)]^{\frac{1}{2}} \quad (5.13a)$$

and
$$Z \sim [1 - \alpha(\xi - 1)] \cos \chi_0 - \pi^{-1} \sin \chi_0 \log [C^2 \alpha(\xi - 1)], \quad (5.13b)$$

where $C = 1.78 \dots$ is Euler's constant. We match (5.12) and (5.13b) by choosing

$$C_1 = \cos \chi_0, \quad \tan \chi_0 = \pi \alpha(1 - \gamma), \quad \xi_1 + (1 - \gamma) \log (\xi_1 - 1) = 1 + (1 - \gamma) \log (1/C\alpha^2), \quad (5.14a-c)$$

wherein error factors of $1 + O(\alpha^2)$ are implicit.

Finally, we let $\chi \rightarrow \infty$ in (5.6a) to obtain (cf. (2.13) with $k_y = 0$ therein)

$$Z \sim (\sigma/\pi)^{\frac{1}{2}} [K(h - \delta)]^{-\frac{1}{2}} \cos (\chi + \chi_0 - \frac{1}{4}\pi), \quad \chi \sim k_\infty x + \chi_*, \quad (5.15a, b)$$

where
$$k_\infty = \left(\frac{K}{h_\infty - \delta} \right)^{\frac{1}{2}}, \quad \chi_* = \frac{-k_\infty \delta}{\sigma} + K^{\frac{1}{2}} \int_{\delta/\sigma}^{\infty} [(h - \delta)^{-\frac{1}{2}} - (h_\infty - \delta)^{-\frac{1}{2}}] dx. \quad (5.16a, b)$$

Combining (5.15) and (5.16), we place the result in the form

$$Z = \frac{1}{2}(\sigma/\pi)^{\frac{1}{2}} R_0^{-\frac{1}{2}} [K(h - \delta)]^{-\frac{1}{2}} (e^{-ik_\infty x} + R_0 e^{ik_\infty x}), \quad (5.17a)$$

where
$$R_0 = \exp [2i(\chi_0 + \chi_* - \frac{1}{4}\pi)] \quad (5.17b)$$

is the reflection coefficient referred to $x = 0$.

We are interested primarily in the magnitude of R_0 . Letting

$$|R_0| \equiv e^{-\rho}, \quad (5.18)$$

and approximating $\tan \chi_0$ by χ_0 in (5.14b) (note that $\alpha\gamma$ is real), we obtain

$$\rho = 2\pi \operatorname{Im} \alpha = 2\pi \sigma^{-2} K \delta_*. \quad (5.19)$$

5.2. Wedge profile

The development in (5.15)–(5.17) assumes smooth $h(x)$, but in the typical wave tank

$$h = \frac{\sigma x}{h_1} \left(x \leq x_1 \equiv \frac{h_1}{\sigma} \right). \quad (5.20)$$

and h' is discontinuous at $x = x_1$. The outer approximation (5.6) then reduces to

$$Z = J_0(\chi) \cos \chi_0 - Y_0(\chi) \sin \chi_0, \quad \chi = 2\sigma^{-1} [K(h - \delta)]^{\frac{1}{2}} \quad (x < x_1), \quad (5.21a, b)$$

while the solution in $x > x_1$ has the form

$$Z = \left(\frac{A_1}{1 + R_1} \right) [e^{-ik_1(x-x_1)} + R_1 e^{ik_1(x-x_1)}], \quad k_1 = \left(\frac{K}{h_1 - \delta} \right)^{\frac{1}{2}} \quad (x > x_1), \quad (5.22a, b)$$

where R_1 is the reflection coefficient referred to $x = x_1$ (note that $h_1 \equiv h_\infty$ and $k_1 \equiv k_\infty$). Requiring Z and Z' to be continuous at $x = x_1$, we obtain

$$A_1 = J_0(\chi_1) \cos \chi_0 - Y_0(\chi_1) \sin \chi_0, \quad R_1 = \exp \left\{ 2i \tan^{-1} \left[\frac{J_1(\chi_1) \cos \chi_0 - Y_1(\chi_1) \sin \chi_0}{J_0(\chi_1) \cos \chi_0 - Y_0(\chi_1) \sin \chi_0} \right] \right\}. \quad (5.23a, b)$$

Letting $\chi_1 \rightarrow \infty$ in (5.23b), we obtain

$$R_1 \sim \exp \{ 2i(\chi_0 + \chi_1 - \frac{1}{4}\pi) [1 + O(\chi_1^{-1})] \}. \quad (5.24)$$

Comparing (5.24) and (5.17b) and invoking (5.16a, b), $h_1 = h_\infty$ and $k_1 = k_\infty$, we obtain

$$R_1 = R_0 \rho^{2ik_1 x_1}, \quad |R_1| = e^{-\rho} [1 + O(\chi_1^{-1})]. \quad (5.25a, b)$$

Considering, for example, the experiments of Mahony & Pritchard (1980), for which $\omega = 9.06 \text{ s}^{-1}$, $\sigma = 0.090$, $h_1 = 3 \text{ cm}$, $g = 980 \text{ cm/s}^2$, and $\lambda = 0.28 \text{ cm}$ (for clean

water), we obtain $K = 0.084 \text{ cm}^{-1}$, $|\delta| = 0.033 \text{ cm}$, $|\alpha| = 0.34$, $|\gamma| = 0.76$, $|\chi_1| = 11.2$, $\rho = 1.52$ and $|R_1| = 0.22$. The observed value was $|R_1| = 0.114$ ($\rho = 2.17$). If the surface were fully contaminated δ would be doubled, which would yield $\rho = 3.0$ and $|R_1| = 0.048$. Mahony & Pritchard state that 'the surface of the water was skimmed with a vacuum pump' before the start of each experiment, but this may not have been sufficient to avoid at least partial contamination before the completion of a particular measurement. Allowing for this possibility, we conclude that the agreement between the present calculation of $|R_1|$ and Mahony & Pritchard's observed value is within the uncertainties associated with the error factor $1 + O(\alpha)$ in the calculation and the free-surface condition in the experiments.

Guza & Bowen (1976) observed almost perfect reflection for $\sigma = 0.1$ and $2\pi/\omega = 2.4\text{--}3.4 \text{ s}$ ($|\alpha| = 0.043\text{--}0.025$), for which (5.26) predicts $|R_1| = 0.90\text{--}0.82$ for clean water. However, their estimate of complete reflection was based on the fit of the observed profile to that predicted by the inviscid theory and is as consistent with $|R_1| = 0.8$ as with $|R_1| = 1$ (Guza, private communication).

This work was supported in part by the Physical Oceanography, Applied Mathematics and Fluid Dynamics/Hydraulics programmes of the National Science Foundation, NSF Grant OCE-85-18763, by the Office of Naval Research, Contract N00014-84-K-0137, 4322318 (430), and by the DARPA Univ. Res. Init. under Appl. and Comp. Math. Programme Contract N00014-86-K-0758 administered by the Office of Naval Research.

REFERENCES

- BALL, F. K. 1967 Edge waves in an ocean of finite depth. *Deep-Sea Res.* **14**, 79–88.
- CARRIER, G. F. & GREENSPAN, H. P. 1958 Water waves of finite amplitude on a sloping beach. *J. Fluid Mech.* **4**, 97–109.
- ERDÉLYI, A. 1955 *Asymptotic Expansions*. Dover.
- FRIEDRICHS, K. O. 1948 Water waves on a sloping beach. *Commun. Appl. Maths* **1**, 109–134.
- GUZA, R. T. 1985 Laboratory modelling of edge waves. *Physical Modelling in Coastal Engineering* (ed. R. A. Dalrymple), pp. 105–116. Balkema.
- GUZA, R. T. & BOWEN, A. J. 1976 Resonant interactions for waves breaking on a beach. *Proc. 15th Coastal Engng Conf. Honolulu*, pp. 560–579.
- KELLER, J. B. 1958 Surface waves on water of non-uniform depth. *J. Fluid Mech.* **4**, 607–614.
- KELLER, J. B. 1961 Tsunamis – Water waves produced by earthquakes. *Tsunami Hydrodynamics Conference* (IUGG Monograph no. 24), pp. 154–166.
- LAMB, H. 1932 *Hydrodynamics*. Cambridge University Press.
- MILES, J. 1990 Wave motion in a viscous fluid of variable depth. *J. Fluid Mech.* **212**, 365–372.
- MAHONY, J. J. & PRITCHARD, W. G. 1980 Wave reflexion from beaches. *J. Fluid Mech.* **101**, 809–832.
- STOKER, J. J. 1957 *Water Waves*. Interscience.