





Automated Grid Generation for WAVEWATCH III

Prepared by Arun Chawla Research Physicist, EMC/NCEP

Jose-Henrique Alves Senior Research Scientist, SRG at EMC/NCEP NOAA Center for Weather & Climate Prediction 5830 University Research Ct College Park, MD 20740 Henrique.Alves@NOAA.gov







Outline

Covered in this lecture:

- The grids necessary for running WAVEWATCH III codes
- The reference data used for creating the grids
- The different modules of a grid generation software to develop the grids
- Additional changes needed to the grids to allow two way communications between grids for multi-grid WAVEWATCH III



Motivation



WAVEWATCH III requires 3 grids (1 necessary and two optional)

- A bathymetry grid (necessary)
- A land –sea mask grid (optional in WW III v 2.22, needed in multi-grid version of WW III)
- Obstruction grid to account for energy decay due to sub-grid blocking (optional)
- Development of these grids can be a fairly arduous task. This is specially true for multi-grid version of WW III where grid consistency across overlapping grids is necessary for accurate two – way coupling.
- To facilitate this we developed a set of algorithms that can automatically create accurate grids with minimal input from the user.
- The necessary tools have been developed using MATLAB.

Step 1: Build a bathymetric grid from a high resolution base bathymetry

NCEP

Step 2: Create an appropriate land / sea mask to accurately depict coast lines

Step 3: Mask out un-necessary water bodies

Step 4: Generate obstruction grids for unresolved islands

Step 5: Identify nodes for exchanging boundary information (multiple-grid only)









Reference Data



RS FOR ENVIRONMENT

NCEP

Subroutines



A grid generation routine

Generates a grid from base bathymetric data set. We use global ETOPO2/ETOPO1 bathymetry netcdf files. (NOTE: To use this routine your matlab should be able to handle netcdf files. Capability is available as open source)

A boundary extraction routine

 Extracts a subset of boundaries from a global set of polygon boundaries (GSHHS)

A land-mask routine

 Blends the bathymetric data with the coastal boundaries to develop an accurate land-sea mask

A water body routine

- Groups the wet cells into different water bodies (each having a unique id)
- A sub-grid obstruction routine
 - Develops sub-grid obstruction matrices in x and y direction for wet cells



Fall 2013 Ifremer Short Course on Ocean Waves

S FOR ENVIRONMENT



Putting it together



Reference_data/ (base bathymetry and coastal polygons)

GRIDGEN/ (Parent directory)

bin/ (individual matlab subroutines)

examples/ (Templates of master scripts used to generate grids) S FOR ENVIRONMENT

NCEP

Grid Generation Routine



- Routine uses 2D averaging to interpolate the higher resolution (reference) grid at the lower resolution
- Averaging carried out over all the reference "wet cells" that lie within a grid cell
- If the proportion of reference wet cells is less than user specified cut-off (ranging between 0 and 1) then the grid cell is marked dry.
- Averaging filters out higher spatial frequencies and prevents aliasing





Bathymetry (Hawaiian islands) using different cutoffs for proportion of wet cells







Grid generation 10/32

Fall 2013 Ifremer Short Course on Ocean Waves

Boundary extraction routine



Uses the GSHHS (Global Self-consistent Hierarchical High-resolution Shoreline) polygon database.

High resolution data is available as mat files at 5 resolutions

≻ Full

NCEP

- high (0.2 km)
- intermediate (1 km)
- > low (5 km)

coarse (25 km)

- Only accounts for land sea boundaries (ignores lakes)
- Properly splits boundaries that are intersecting grid domain boundary

Important to properly close boundaries for land masks and sub grid obstructions



OT CENTERS FOR ENVIRONME

NCEP

Splitting a boundary properly







Fall 2013 Ifremer Short Course on Ocean Waves

Grid generation 13/32



Fall 2013 Ifremer Short Course on Ocean Waves

Grid generation 14/32

ON CENTERS FOR EA

NCEP



Full Resolution Global Bound



Fall 2013 Ifremer Short Course on Ocean Waves

Grid generation 15/32

NTERS FOR ENVIRONMENTAL ARE

Land-sea mask routine



- A first order land mask (based on a depth cutoff) is determined
- The land mask routine then
 - > Checks all wet cells near the boundaries
 - Switches wet cells to dry if a proportion of cell within the boundary > user specified cutoff (currently set at 0.5)
- Land mask routine needed to
 - > Account for land masses not present in base bathymetry grid
 - Resolve discrepancies between shoreline data base and base bathymetry shorelines
 - > Account for additional polygons

FOR ENVIRONMENT

NCEP

To compute proportion of cell inside boundary



 Equally spaced points spread throughout the cell
Determine number of points enclosed within boundary
Number of points inside boundary/Total number of



points

OTHERN FOR EN

NCEP

Land-Sea Mask (Bahamas 15 min grid)



Red – Wet Cell Blue – Dry Cell White – Wet cell switched to dry cell



Fall 2013 Ifremer Short Course on Ocean Waves

Grid generation 18/32

out centers.

NCEP



Depth (0.1)

Depth + Mask



-5000-4000-3000-2000-1000 0

Fall 2013 Ifremer Short Course on Ocean Waves







Boundary Splitting

- Mask cleaning requires checking parts of cell being inside/outside a boundary
- For very large boundaries this can be a very cumbersome process
- An optional algorithm splits up the larger boundaries into more manageable smaller boundaries.
- Significantly speeds up the mask cleaning algorithm

OT CENTERS FOR ENVIRONME

NCFP

Boundary Splitting





Fall 2013 Ifremer Short Course on Ocean Waves

Grid generation 21/32

Wet cell clean up routine



- Cycle through all the wet cells and flag all connected cells with the same id
- Independent water bodies have different ids
- Function returns an id map that allows users to switch cells of a particular water body from wet to dry
- Switching of cells can either be done inside the routine with a flag option, or outside by the user

NCEP

S FOR ENVIRONMENT

NCEP

Wet Cell clean up (contd.)



- Initialize all wet cells as unmarked
- Starting from first unmarked cell with marker value at 1, mark all connected wet cells with the same marker
- If more unmarked cells then increment marker by 1 and repeat step 2.
- Keep repeating steps 3 and 2 till no longer unmarked wet cells
- End result is a mask map with the wet cells grouped into independent water bodies





CENTERS FOR ENVIRONMENTAL ARE A

Obstruction grid algorithm



Start from the final land – sea mask and boundary polygons (not the split polygons)

- Following the algorithm outlined in obstruction grid presentation build the obstruction grids for all the wet cells
- Assign obstructions next to dry cells to zero (to prevent spurious attenuation of wave energy)





At this point



The grid generation software has created

- A bathymetric grid
- A final land sea mask
- A pair of obstruction grids associated with the land sea mask and (unresolved) boundaries within the grid
- When using WW-III in single grid mode this is all the information needed to create model definition files
- When using WW-III in multiple grid mode with two way nesting between the grids further considerations have to be made that will be outlined next
- If building multiple grids, repeat steps till now for each of the individual grids before proceeding to the next step

Multi-grid algorithms



Additional changes to support two – way nesting

- Two way nesting in WW-III occurs as follows
 - Boundary information from low resolution grids to high resolution grids
 - Grid cell averaging from high resolution back to low resolution

> Grid resolution across overlapping grids of similar resolution

- In multi-grid WW-III, masks can have a value of 0 (land), 1 (ocean), 2 (boundary ; for data transfer from low resolution grids) and 3 (undefined ; outside the domain).
- Two additional algorithms to modify the masks for multi-grid WW-III
 - A mask modifying algorithm that identifies boundary positions
 - > A mask reconciliation algorithm

NCEP

Modifying mask









Fall 2013 Ifremer Short Course on Ocean Waves

OT CENTERS FOR ENVIRONMENTS

NCEP

Grid generation 27/32

Modifying masks



- Once grid information for all the individual grids have been developed, the modify mask routine can be used with pairs of base and higher resolution grids to modify masks for exchange of information from low resolution to high resolution
- Exchange of information from high to low resolution does not occur at a boundary so mask does not need to be modified
- Lowest resolution grid also does not need the mask to be modified
- If using the entire higher resolution grid (instead of carving out a domain inside the grid) then the polygon is defined by the corners of the grid
- For overlapping grids with similar resolution (where data is reconciled) a common overlap area has to be defined



NCEP

Overlapping grids principle



Overlaping length = stencil_width * global_time_step/CFL_time_step + stencil_width*global_time_step/CFL_time_step+1











Grid 1 extends from 65N to 82 N

Grid 2 extends from 77.5S to 77.5 N



Putting it together ...



Grid generation 30/32

Overlapping grids



- If there is not enough overlapping domain in overlapping grids of similar resolution the WW-III model will fail
- Since the minimum required domain width is a function of the global and CFL time steps in the respective grids, changing any can have an important impact on required overlapping domain width
- Land sea masks in the two grids with overlapping domains has to be similar
- An optional land sea mask reconciliation algorithm has been developed
 - > Compares two sets of masks in an overlapping domain
 - If the mask at any location are different then they are both set to land values (0)

NCEP



The end



End of lecture

Fall 2013 Ifremer Short Course on Ocean Waves

Grid generation 32/32